

BOTTLENOSE DOLPHIN (*Tursiops truncatus*): Western North Atlantic Coastal ~~Stock~~Morphotype Stocks

STOCK DEFINITION AND GEOGRAPHIC RANGE

Stock Structure of the Coastal Morphotype

A. Latitudinal distribution and structure along the coast

The coastal morphotype of bottlenose dolphin is continuously distributed along the Atlantic coast south of Long Island, around the Florida peninsula ~~Florida~~ and along the Gulf of Mexico coast. On the basis of differences in mtDNA haplotype frequencies, ~~however~~, Curry (1997) concluded that the nearshore animals in the northern Gulf of Mexico and the western North Atlantic ~~were significantly different and~~ represent separate stocks.

~~Within the western North Atlantic, the stock structure of coastal bottlenose dolphins is complex.~~ Scott et al. (1988) hypothesized a single coastal migratory stock ranging seasonally from as far north as Long Island, NY, to as far south as central Florida, citing stranding patterns during a high mortality event in 1987-88 and observed density patterns along the US Atlantic coast. ~~The continuous distribution of dolphins along the coast seemed to support this hypothesis. It was recognized that bottlenose dolphins were resident in some estuaries; these were considered to be separate from the coastal migratory animals.~~ More recent studies suggest that the single coastal migratory stock hypothesis is incorrect and that there is likely a complex mosaic of stocks. ~~For example, year-round resident populations have been reported at a variety of sites in the southern part of the range, from Charleston, South Carolina (Zolman 1996) to central Florida (Odell and Asper 1990); seasonal residents and migratory or transient animals also occur in these areas (summarized in Hohn 1997). In the northern part of the range the patterns reported include seasonal residency, year-round residency with large home ranges, and migratory or transient movements (Bareo and Swingle 1996, Sayigh et al. 1997). Communities of dolphins have been recognized in embayments and coastal areas of the Gulf of Mexico (Wells et al. 1996; Scott et al. 1990; Weller 1998) so it is not surprising to find similar situations along the Atlantic coast (NMFS 2001; McLellan et al. 2003).~~

Recent genetic analyses of samples from Jacksonville, FL, ~~southern~~Georgia, central South Carolina (primarily the estuaries around Charleston), southern North Carolina, and coastal Virginia, using both mitochondrial DNA and nuclear microsatellite markers, indicate that a significant amount of the overall genetic variation can be explained by differences between ~~the groups~~ (NMFS 2001). ~~The degree of population subdivision, estimated using the parameter F_{ST} , between each of the groups was statistically significant~~these areas (NMFS 2001). These results indicate a minimum of ~~four populations~~five stocks of coastal bottlenose dolphins ~~in~~along the ~~Northwest~~U.S. Atlantic ~~coast~~ and reject the null hypothesis of one homogeneous population of bottlenose dolphins.

~~Another potential population has been identified from stable isotope ratios of oxygen (NMFS 2001). Animals sampled along the beaches of North Carolina between Cape Hatteras and Bogue Inlet during the months of February and March show very low stable isotope ratios of ^{18}O relative to ^{16}O (referred to as depleted ^{18}O or depleted oxygen) (Cortese 2000). One possible explanation for the depleted oxygen signature is that there is a resident group of dolphins in Pamlico Sound. Alternatively, these animals may represent a component of the migratory animals that spend their summers at the northernmost end of the range of bottlenose dolphins and winter in North Carolina. Either possibility suggests they represent a separate stock. Stable isotope ratios of ^{18}O from samples taken in estuarine waters around Charleston, SC, showed little variation and none were at depleted levels.~~

Photo-identification studies also support the existence of multiple stocks (NMFS 2001). A coastwide photographic catalogue has been established using contributions from 15 sites from Cape May, NJ, to Cape Canaveral, FL (Urian et al. 1999). No matches have been found between the northernmost and southernmost sites. However, there appears to be a high rate of exchange among northern field sites, where dolphins occur only seasonally, and central North Carolina including the Beaufort area. Other areas of frequent exchange include Beaufort and Wilmington, NC. In contrast to the patterns found in the northern end of the range, there appears to be less movement between southern field sites – there are only two confirmed matches between the relatively large catalogs of Jacksonville, FL, and Hilton Head, SC, for example, and no matches between the Charleston, SC site and other sites.

Satellite-linked radio transmitters have been deployed on dolphins in Virginia Beach, VA, Beaufort, NC, ~~and~~ Charleston, SC and New Jersey. The movement patterns of animals with satellite tags provided additional

information that was complementary to ~~the photo-identification, genetic, and stable isotope studies~~ other stock identification approaches. The results, along with photo-identification of freeze-branded animals, indicate that a significant number of dolphins reside in NC in summer and do not migrate. ~~Satellite telemetry results reinforced the photo-identification results from Charleston, SC, indicating a resident population there.~~ Finally, a dolphin tagged in Virginia Beach, VA, spent the winter between Cape Hatteras and Cape Lookout, NC, ~~rather than migrating to Florida as would have been expected in the single coastal migratory stock hypothesis (NMFS 2001).~~


~~Stable~~ Stable, indicating seasonal migration between North Carolina and areas further north (NMFS 2001).

~~Another potential stock has been identified from stable~~ isotope ratios of oxygen ~~suggested the possibility of a resident group of bottlenose dolphins in Pamlico Sound, NC~~ (NMFS 2001). Animals sampled along the beaches of North Carolina between Cape Hatteras and Bogue Inlet during the months of February and March show very low stable isotope ratios of ^{18}O relative to ^{16}O (referred to as depleted ^{18}O or depleted oxygen) (Cortese 2000). One possible explanation for the depleted oxygen signature is that there is a resident group of dolphins in Pamlico Sound that move into nearby nearshore areas in the winter ~~when Pamlico Sound may have an inadequate resource base~~. The possibility of a resident group of bottlenose dolphins in Pamlico Sound is supported by the results from satellite telemetry and photo-identification results. Alternatively, ~~however~~, these animals may represent a component of the migratory animals that spend their summers at the northernmost end of the range of bottlenose dolphins and winter in North Carolina. Either possibility suggests they represent a separate stock. ~~Stable isotope ratios of ^{18}O from samples taken in estuarine waters around Charleston, SC, showed little variation and none were at depleted levels.~~

There are additional resident estuarine stocks that are likely demographically distinct from coastal stocks, but they are currently included in the coastal management unit definitions. For example, year-round resident populations have been reported at a variety of sites from Charleston, South Carolina (Zolman 1996) to central Florida (Odell and Asper 1990). Seasonal residents and migratory or transient animals also occur in these areas (summarized in Hohn 1997). In the northern part of the range, the patterns reported include seasonal residency, year-round residency with large home ranges, and migratory or transient movements (Barco and Swingle 1996, Sayigh et al. 1997). Communities of dolphins have been recognized in embayments and coastal areas of the Gulf of Mexico (Wells et al. 1996; Scott et al. 1990; Weller 1998), and it is not surprising to find similar situations along the Atlantic coast.

In summary, integration of the ~~preliminary~~ results from genetics, photo-identification, satellite telemetry, and stable isotope studies confirms a complex mosaic of ~~stocks of coastal bottlenose dolphins in the western North Atlantic stocks~~. As an interim measure, pending additional results, seven management units within the range of the ~~“coastal migratory stock”~~ coastal morphotype of western North Atlantic bottlenose dolphin have been defined (Figure 1). The true population structure is likely more complex than the seven units identified in this report, ~~and~~ research efforts continue ~~in an attempt~~ to identify that structure.

Figure 1. Management units of the coastal morphotype of bottlenose dolphins along the Atlantic coast of the U.S. as defined from recent results from genetic, stable isotope ratio, photo-identification, and telemetry studies (~~per Hohn 1997~~; NMFS 2001).



B. Longitudinal distribution

Earlier aerial surveys conducted between 1978-1982 (CETAP 1982) and shipboard (NMFS unpublished data) surveys north of Cape Hatteras, North Carolina identified two concentrations of bottlenose dolphins, one inshore of the 25 m isobath and the other offshore of the 2550 m isobath. The lowest density of bottlenose dolphins was observed over the continental shelf, with higher densities along the coast and near the continental shelf edge. It was suggested, therefore, that the coastal morphotype is restricted to waters < 25 m in depth north of Cape Hatteras (Kenney 1990). There was no apparent similar patterns were observed during summer months north of Cape Lookout, North Carolina in more recent aerial surveys (Garrison and Yeung 2001; Garrison et al. 2003). However, south of Cape Lookout during both winter and summer months, there was no clear longitudinal discontinuity in bottlenose dolphin herd sightings sightings (Garrison and Yeung 2001; Garrison et al. 2003). However, dolphin groups observed during aerial surveys south of Cape Hatteras in the winter (Blaylock and Hoggard 1994). NMFS surveys conducted from 1992-1998 show a clustering of bottlenose dolphins nearshore and then additional bottlenose dolphins in the offshore areas. However, the morphotype of bottlenose dolphins (WNA offshore or WNA coastal) can not be determined from the air so attributing each sighting cannot be attributed to a specific morphotype cannot be done. There is also a potential for confusing immature spotted dolphins, with few or no spots dorsally, with based on sighting information alone.

Genetic analysis of tissue samples can be used to identify animals to a specific morphotype (Hoelzel et al., P. Rosel SEFSC unpublished results). An analysis of tissue samples from large vessel surveys during the summers of 1998 and 1999 indicated that bottlenose dolphins where the two species are sympatric.

In 1995, NMFS conducted two aerial surveys along the Atlantic coast (Blaylock 1995; Garrison and Yeung 2001). One survey was conducted during summer 1995 between within 7.5 km from shore were most likely of the coastal morphotype, and there was an extensive region of overlap between the coastal and offshore morphotypes between 7.5 and 34 km from shore south of Cape Hatteras, NC, and Sandy Hook, NJ, and included three replicate surveys. The second survey was conducted during winter 1995 between Cape Hatteras, NC, and Ft. Pierce, FL. A

distributional analysis identified a significant spatial pattern in (Torres et al. 2003). However, relatively few samples were available from the region of overlap, and therefore the longitudinal boundaries based on these initial analyses are uncertain (Torres et al. 2003). Extensive systematic biopsy sampling efforts were conducted using small vessels in the summers of 2001 and 2002 to supplement collections from large vessel surveys. During the winters of 2002 and 2003, additional biopsy collection efforts were conducted in nearshore continental shelf waters of North Carolina and Georgia. A small number of additional biopsy samples were collected in deeper continental shelf waters south of Cape Hatteras during winter 2002. Genetic analyses of these biopsies identified individual animals to the coastal or offshore morphotype. Based upon the genetic results from all surveys combined, a logistic regression approach was used to model the probability that a particular bottlenose dolphin sightings as a function of distance from shore (Garrison 2001a). During the northern (summer) surveys, the significant spatial boundary occurred at 12 km from shore. During the southern (winter) survey, the significant spatial boundary occurred at 27 km from shore. The gap in sightings best defines, for the time being, the eastern extent group is of the coastal morphotype for purposes of habitat definition and abundance estimates. NMFS continues to collect as a function of environmental variables including depth, sea surface temperature, and distance from shore. These models were used to partition the bottlenose dolphin groups observed during aerial surveys between the two overlapping morphotypes (Garrison et al. 2003).

The genetic results and spatial patterns observed in aerial surveys indicate both regional and seasonal differences in the longitudinal distribution of the two morphotypes in coastal Atlantic waters. North of Cape Lookout, North Carolina (i.e., northern migratory and northern North Carolina management units) during summer months, the previously observed pattern of strong nearshore aggregation of bottlenose dolphins was again observed. All biopsy samples collected from Tursiops throughout the possible range nearshore waters (<20 m depth) were of the coastal morphotype so that stock boundaries can be confirmed or modified on the basis of a more comprehensive data set.

and all offshore samples (> 40m depth) were of the offshore morphotype. The genetic results confirm separation of the two populations in this region during summer months. South of Cape Lookout, NC, the probability of an observed bottlenose dolphin group being of the coastal morphotype declined with increasing depth; however, there was significant spatial overlap between the two morphotypes. Offshore morphotype bottlenose dolphins were observed at depths as shallow as 13m, and coastal morphotype dolphins were observed at depths of 31m and 75 km from shore (Garrison et al. 2003). These results indicate significant overlap between the two morphotypes in the southern management units during summer months.

Winter samples were collected primarily from nearshore waters in North Carolina and Georgia. The vast majority of samples collected in nearshore waters of North Carolina during winter were of the coastal morphotype; however, one offshore morphotype group was sampled during November just south of Cape Lookout, North Carolina only 7.3 km from shore. Coastal morphotype samples were also collected further away from shore at 33 m depth and 39 km from shore. The logistic regression model for this region indicated a decline in the probability of a coastal morphotype group with increasing distance from shore; however, the model predictions are highly uncertain due to limited sample sizes and high overlap between the two morphotypes. Samples collected in Georgia waters also indicated significant overlap between the two morphotypes with a declining probability of the coastal morphotype with increasing depth. A coastal morphotype sample was collected well offshore at a distance of 112 km from shore and a depth of 38 m. An offshore sample was collected in 22 m depth at 40 km from shore. As with the North Carolina model, the Georgia logistic regression predictions are uncertain due to limited sample size and high overlap between the two morphotypes (Garrison et al. 2003). The logistic regression models were used to predict the probability that an observed bottlenose group is of the coastal morphotype as a function of habitat variables and spatial location. There remain significant sampling gaps in the biopsy collections, particularly during winter months, that increase the uncertainty of model predictions. Both the predicted probability of a coastal morphotype occurring and the associated uncertainty in that prediction are incorporated into the abundance estimates for coastal morphotype bottlenose dolphin management units.

POPULATION SIZE

The 1995 Previous abundance estimates for the coastal morphotype of WNA bottlenose dolphin were based primarily upon aerial surveys were conducted to estimate population size of the hypothesized conducted during the

summer and winter of 1995. The surveys were designed based upon the previous assumption of a single coastal migratory stock, and therefore they did not provide complete seasonal and spatial coverage for the more recently defined management units. Previous abundance estimates were also not corrected for visibility bias (Blaylock 1995; Garrison and Yeung 2001). The summer aerial survey was conducted between July 1 and August 14, 1995, covering Cape Hatteras, NC, to Sandy Hook, NJ, (35.23°N 40.5°N), and from the mainland shore to the 25 m isobath. This survey provided coverage and abundance. Aerial surveys to update the abundance estimates were conducted during winter (January-February) and summer (July-August) of 2002. Survey tracklines were set perpendicular to the shoreline and included coastal waters to depths of 40m. The surveys employed a stratified design so that most effort was expended in waters shallower than 20m depth where a high proportion of observed bottlenose dolphins were expected to be of the coastal morphotype. Survey effort was also stratified to optimize coverage in seasonal management units. The surveys employed two observer teams operating independently on the same aircraft as an approach to estimate and account for sources of visibility bias.

The winter survey included the region from the Georgia/Florida state line to the southern edge of Delaware Bay. A total of 6,411 km of trackline was completed during the survey, and 185 bottlenose dolphin groups were sighted including 2,114 individual animals. No bottlenose dolphins were sighted north of Chesapeake Bay corresponding to water temperatures <9.5 °C. During the summer survey, 6,734 km of trackline were completed between Sandy Hook, NJ to Ft. Pierce, FL. All tracklines in the 0-20 m stratum, were completed throughout the survey range while offshore lines were completed only as far south as the Georgia-Florida state line. A total of 185 bottlenose dolphin groups were sighted during summer including 2,544 individual animals.

Abundance estimates for the Northern Migratory (NM) and Northern North Carolina (NNC) management units. However, coverage of the NNC unit was incomplete as the surveys did not cover the region south of Cape Hatteras, NC, to Cape Lookout, NC. Abundance was estimated for each stratum pooling across the three replicate surveys. The winter survey was conducted between January 27 and March 6, covering from Fort Pierce, FL, to Cape Hatteras, NC, (27.30°N 35.23°N), from the mainland shore to 9.25 km (5 Nautical Miles) beyond the inshore edge of the Gulf Stream or <200 km offshore. This survey included coverage of the NNC, Southern North Carolina (SNC), South Carolina (SC), Georgia (GA), Northern Florida (NFL) and Central Florida (CFL) management units. However, the coverage of the NNC management unit was incomplete and did not include the region north of Cape Hatteras, NC. These abundance estimates also include NM unit animals that have migrated south of the NC/VA border during winter. Abundance for bottlenose dolphins in each management unit was estimated were calculated using line transect methods and the program DISTANCE (Buckland *et al.* 1993) for both the winter and summer surveys (Table 1). There was no significant difference between distance analysis (Buckland *et al.* 1993). The independent and joint estimates from the two survey teams were used to quantify the probability that animals available to the survey on the trackline were missed by the observer teams, or perception bias, using the direct duplicate estimator (Palka, 1995). These estimates were further partitioned between the coastal and offshore morphotypes based upon the results of the logistic regression models and spatial analyses described above. A parametric bootstrap approach was used to incorporate the uncertainty in the logistic regression models into the overall uncertainty in the abundance estimates for the combined NM and NNC each management units in summer and the combined NM, NNC, and SNC stocks in winter.

Another set of unit (Garrison *et al.* 2003).

The aerial surveys was conducted parallel to the coastline from the North Carolina/South Carolina border to the Maryland/Delaware border during 1998 and 1999 to document the distribution of dolphins and fishing gear in nearshore waters (Hohn *et al.* unpubl. data). These strip transect surveys were conducted weekly, weather permitting, over 12 months in most of North Carolina and for six months (May to December) in Virginia and Maryland. In retrospect, they provide seasonal coverage of the Southern North Carolina, Northern North Carolina, and Northern Migratory management units (Figure 1; Hohn *et al.* unpubl. data). The strip transect surveys cannot be used directly for abundance estimation because they did not follow the design constraints of line transect survey methods and covered only a small proportion of the habitat of coastal bottlenose dolphin. The density of dolphins near the coastline is high relative to habitats further offshore, and the use of density estimates in this region to calculate overall abundance would likely result in significant positive bias. However, these surveys do provide information on the relative abundance of dolphins between regions that may be used to supplement the included only animals in coastal waters, and the resulting abundance estimates from the line transect surveys conducted in 1995 (Garrison and Hohn 2001). Both sets of aerial surveys covered ocean coasts only therefore do not include animals inside estuaries that are currently included in the defined management units. An abundance estimate was

generated for bottlenose dolphins in estuarine waters of Pamlico Sound, North Carolina using mark-recapture methodology (Read et al. 2004³). It is possible to post-stratify the mark-recapture estimates, and these estimates were post-stratified to be consistent with management unit definitions (Palka et al. 2001) (Table 1).

2001a; Table 1). Since abundance estimates do not exist for all estuarine waters, the overall estimates and PBRs for these management units are negatively biased.

Bottlenose dolphins in the northern migratory stock migrate south during winter months and overlap with those from the northern North Carolina and southern North Carolina management units. It is not possible at this time to apportion the incidental mortality occurring during winter months in North Carolina waters among animals from these three management units. Therefore, a half-year PBR value is applied for each management unit in the summer based upon abundance estimates from summer aerial surveys. During winter months, these three stocks overlap spatially and a half-year PBR is applied to the North Carolina mixed management unit based upon winter aerial survey abundance estimates. For the South Carolina and Georgia management units, the abundance estimates, minimum population size values, and the resulting PBR values are derived using a weighted average of abundance estimates from the winter and summer 2002 aerial surveys. The northern Florida management unit was only surveyed during the summer of 2002 and the winter of 1995. The resulting abundance estimate is therefore the inverse variance weighted average of the seasonal estimates from the available surveys. Finally, the central Florida management unit was only covered during the 1995 surveys. Due to the age of the available abundance estimate, the PBR of the central Florida management unit was set to "undefined".

Table 1. Estimates of abundance and the associated CV, n_{min} , and PBR for each management unit of WNA coastal bottlenose dolphins (from Palka-Garrison et al. 2004³). The PBR for the Northern Migratory, Northern NC, and Southern NC management units are applied biannually semi-annually. For management units south of NC, the PBR is applied annually.

Management Unit		Best Abundance		N _{min}	PBR	
		Estimate	CV		Annual	½ Yr
SUMMER (May - October)						
Northern migratory		568 <u>17,466</u>	<u>19.1</u>	24,441 <u>14,640</u> <u>1</u>	(461) <u>46.2</u>	<u>2373.1</u>
Northern NC						
	oceanic	<u>6,160</u>	<u>51.9</u>	3,383 <u>41,824</u> <u>3255</u>	(24) <u>32.6</u>	<u>126.1</u>
	estuary <u>estuary</u> ⁴	919	12.5	828	(8.3)	4.2
	BOTH	<u>7,079</u>	<u>45.2</u>	4,302 <u>33,083</u>	3,281 <u>(33,40.8)</u>)	<u>1620.3</u>
Southern NC						
	oceanic	<u>3,645</u>	<u>111.0</u>	1,457 <u>507</u> <u>7786</u> <u>3</u>	(718.8)	<u>9.3</u> 9
	estuary <u>estuary</u> ⁴	141	15.2	124	(1.2)	0.6
	BOTH	<u>3,786</u>	<u>106.9</u>	1,298 <u>44,690</u> <u>987</u>	(919.4)	<u>49.5</u> 9
WINTER (November - April)						
NC mixed <u>mixed</u> ¹		616,474 <u>39,74,691</u> (47) <u>913</u>	<u>23.0</u>	<u>13,558</u>	<u>(135.6)</u>	<u>67.8</u>
<u>ALL YEAR</u>						
South Carolina		<u>2,351</u> <u>325</u>	<u>20.3</u>	<u>471,963</u>	<u>2,412</u> <u>2419.6</u>	na
Georgia		<u>767</u> <u>782,195</u>	<u>29.9</u>	<u>1,716</u>	<u>17,442</u> <u>84.32</u>	na
Northern Florida <u>Florida</u> ²		<u>354</u> <u>448</u>	<u>38.4</u>	<u>562</u> <u>28328</u>	<u>23.3</u>	na

Central Florida <u>Florida</u> ³	10,652	45.8	7,377 <u>na</u>	74 <u>na</u>	na
--	--------	------	----------------------------	-------------------------	----

~~NC~~NC mixed⁴ = northern migratory, Northern NC, and Southern NC

~~Abundance estimates for each management unit are the sum of estimates, where appropriate, from the recent analyses. Estimated overall abundance was 9,206 from summer surveys and 19,459 from winter surveys. However, for consistency with achieving the goals of the MMPA, such as maintaining marine mammals as functioning components of their ecosystems, it is more appropriate to establish abundance estimates for each management unit. Abundance for each management unit was estimated by post-stratifying sightings and effort data consistent with geographic and seasonal management unit boundaries (Table 1) (Garrison and Yeung 2001; Palka *et al.* 2001). Although these estimates are better than previous abundance estimates for coastal bottlenose dolphins, there remain potential biases. The aerial survey estimates are not corrected for g(0), the probability of detecting a group on the track line as a function of perception bias and availability bias. The exclusion of g(0) from the abundance estimate results in a negative bias of unknown magnitude. The relatively large herd sizes in the summer surveys north of Cape Hatteras likely reduce this bias; however, herd sizes were smaller south of Cape Hatteras during winter, likely resulting in greater negative bias (Palka, unpub. data). A positive bias may occur if the longitudinal boundaries have been extended too far offshore resulting in offshore dolphins being included in the abundance estimates for the coastal morphotype or if estuarine dolphins were overrepresented in coastal waters during the time of the survey. Further uncertainties in the abundance estimates result from incomplete coverage of some seasonal management units during the line transect surveys. While the strip transect surveys were used to supplement the survey coverage, uncertainties associated with that analysis also introduce uncertainty in the overall abundance estimate (Garrison and Hohn, 2001). The SEFSC intends to conduct both winter and summer coastwide aerial surveys during 2002 to obtain more robust abundance estimates.~~

²Northern Florida estimates are the inverse variance weighted mean of abundance
estimates from the winter 1995 survey and the summer 2002 survey.

³Central Florida estimates are from the winter 1995 survey and cannot be used to determine PBR due to

their age.

⁴Read et al., 2003

Minimum Population Estimate

The minimum population size (~~NMIN~~Nmin) for each management was calculated ~~according to Equation 1 from the PBR Guidelines as the lower bound of the 60% confidence interval for a lognormally distributed mean~~ (Wade and Angliss 1997): $NMIN = N / \exp(0.842 \times [\ln(1 + [CV(N)]^2)]/2)$ (Table 1). ~~It is recognized that these~~ For the estimates derived from bootstrap resampling, the appropriate Nmin was taken directly from the bootstrap distribution of abundance estimates. These estimates may be negatively biased because they do not include ~~corrections for g(0) and, for some of the managements units, do not include the entire spatial range of the unit during that season. The strip transect surveys compensate for some of the abundance omitted during line transect survey; nonetheless, for some management units the entire range was not covered.~~ estuarine animals and do not fully account for visibility bias.

Current Population Trend

There are insufficient data to determine the population trend for ~~this~~these stocks~~s~~.

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Current and maximum net productivity rates are not known for the WNA coastal morphotype. The maximum net productivity rate was assumed to be 0.04. This value is based on theoretical modeling showing that cetacean populations may not grow at rates much greater than 4% given the constraints of their reproductive life history (Barlow *et al.* 1995).

POTENTIAL BIOLOGICAL REMOVAL

Potential Biological Removal (PBR) is the product of the minimum population size, one-half the maximum productivity rate, and a “recovery” factor (Wade and Angliss 1997). The “recovery” factor is assumed to be 0.50, the default for depleted stocks and stocks of unknown status. ~~At least part of the range-wide stock complex is depleted; for the remainder, status is unknown. For consistency with achieving~~ This complex of management units

incorporates the range of the former “WNA coastal migratory stock” that has been defined as depleted under MMPA guidelines. At least some of these management units are likely depleted relative to their optimum sustainable population (OSP) size due both to mortality during the 1987-1988 die-off and historically high incidental mortality in fisheries relative to PBR. The status of these stocks relative to OSP is best described as unknown, and therefore the recovery factor of 0.5 is appropriate for the PBR calculation. Given the known population structure within the coastal morphotype bottlenose dolphins, it is appropriate to apply PBR separately to each management unit so as to achieve the goals of the MMPA, such as maintaining marine mammals as functioning components of their ecosystems, it is more appropriate to establish separate PBRs for each management unit (Table 1). (Wade and Angliss 1997).

ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

Total estimated average annual fishery-related mortality or serious injury resulting from observed fishing trips during 1996-2000 was 233 bottlenose dolphins (CV=0.16) in the mid-Atlantic coastal gillnet fishery. The management units affected by this fishery ~~would be the NM, NNC, and SC~~ are the northern migratory, northern North Carolina, and southern North Carolina management units. An estimated ~~246~~ (CV= 0.89) ~~were taken mortalities occurred annually~~ in the shark drift gillnet fishery off the coast of Florida during 1999-2000~~2~~, affecting the Central and Northern Florida management units. ~~No estimates of mortality from observed trips observer data~~ are available for ~~any of~~ the other fisheries that may interact with WNA coastal bottlenose dolphins. Therefore, the total average annual mortality estimate is considered to be a lower bound of the actual annual human-caused mortality and serious injury.

Fishery Information

Bottlenose dolphins are known to interact with commercial fisheries and occasionally are taken in various kinds of fishing gear including gillnets, seines, long-lines, shrimp trawls, and crab pots (Read 1994; Wang et al. 1994) especially in near-shore areas where dolphin densities and fishery efforts are greatest. There are nine Category II commercial fisheries that interact with WNA coastal bottlenose dolphins in the 2001~~3~~ MMPA List Of Fisheries (LOF), six of which occur in North Carolina waters. Category II fisheries include the mid-Atlantic coastal gillnet, NC inshore gillnet, mid-Atlantic haul/beach seine, NC long haul seine, NC stop net, Atlantic blue crab trap/pot, Southeast Atlantic gillnet, Southeastern U.S. Atlantic shark gillnet and the Virginia pound net ~~(Table 1.1)~~ (see 2001~~3~~ List of Fisheries, 66 FR 42780~~4~~1725, August~~July~~ 15, 2001~~3~~). The mid-Atlantic haul/beach seine fishery also includes the haul seine and swipe net fisheries. The term mid-Atlantic refers to the geographic area south of Long Island, landward to the 72° 30' W. line, and north of the line extending due east from the North Carolina/South Carolina border (66 FR 6545, January 22, 2001).

There are five Category III fisheries that may interact with WNA coastal bottlenose dolphins. Three of these are inshore gillnet fisheries: the Delaware Bay inshore gillnet, the Long Island Sound inshore gillnet, and the Rhode Island, southern Massachusetts, and New York Bight inshore gillnet. The remaining two are the shrimp trawl and mid-Atlantic menhaden purse seine fisheries. There have been no takes observed ~~by the NMFS observer programs~~ in any of these fisheries.

Mid-Atlantic Coastal Gillnet

~~The mid-Atlantic coastal gillnet fishery is actually a combination of small-vessel fisheries that target a variety of fish species, including bluefish, croaker, spiny and smooth dogfish, kingfish, Spanish mackerel, spot, striped bass, and weakfish (Steve et al. 2001). It operates in different seasons targeting different species in different states throughout the range of the coastal morphotype. Most nets are set gillnets without anchors and are fished close to shore. Anchored set gillnets or drift gillnets are used in some fisheries (e.g., monkfish or dogfish). A comprehensive description of coastal gillnet gear and fishing effort in North Carolina is available in Steve et al. (2001).~~ This fishery has the highest documented level of mortality of WNA coastal morphotype bottlenose dolphins, and the North Carolina sink gillnet fishery is its largest component in terms of fishing effort and observed takes. ~~Bycatch estimates are available for the past five years, 1996-2000 (Table 2).~~ Of 12 observed mortalities ~~from between~~ 1995-2000, 5 occurred in sets targeting spiny or smooth dogfish and another in a set targeting “shark” species, 2 occurred in striped bass sets, 2 occurred in Spanish mackerel sets, and the remainder were in sets targeting kingfish, weakfish, or finfish generically (Rossman and Palka 2001). Only two bottlenose dolphin mortalities were observed in 2001-2002, both occurring during in the winter mixed North Carolina unit. The overall estimated level of

mortality has declined during the past two years associated with reductions in fishery effort, reduced levels of observer coverage, and reduced bycatch rates (Rossman and Palka, in review). Due to these significant changes in the behavior of the fishery, bycatch estimates for these fisheries are separated into two periods covering from 1996-2000 and 2001-2002 (Table 2).

Table 2. Summary of the 1996-2000⁹² incidental mortality of bottlenose dolphins (*Tursiops truncatus*) by management unit in the commercial mid-Atlantic coastal gillnet fisheries. Data include the years sampled (Years), the number of vessels active within the fishery (Vessels), type of data used (Data Type), observer coverage (Observer Coverage), mortalities recorded by on-board observers (Observed Mortality), estimated annual mortality (Estimated Mortality), estimated CV of the annual mortality (Estimated CVs), and mean annual mortality (CV in parentheses).

Seasonal Management Unit	Years	Vessels	Data Type ¹	Observer Coverage ²	Observed Serious Injury	Observed Mortality	Estimated Mortality	Estimated CVs ³	Mean Annual Mortality
Summer Northern Migratory	1996-2000	NA	Obs. Data, NER Dealer Data	.05, .03, .02, .03, .03 ₂	0, 0, 0, 0, 0	0, 0, 1, 1, 1 ₂	33, 30, 37, 19, 30 ₂	<u>0.48,</u> <u>0.48,</u> <u>0.48,</u> <u>0.48,</u> <u>0.48</u>	30 (0.22) ₂
	<u>2001-2002</u>			<u>.02, .01</u>	<u>0, 0</u>	<u>0, 0</u>	<u>11, 11</u>	<u>0.35,</u> <u>0.35</u>	<u>11 (0.25)</u>
Summer Northern NC	1996-2000	NA	Obs. Data, NCDMF Dealer Data	.01, .00, <.01, .01, .03 ₂	0, 0, 0, 0, 0	1, 0, 0, 0, 0 ₂	27, 33, 17, 13, 26 ₂	<u>0.61,</u> <u>0.61,</u> <u>0.61,</u> <u>0.61,</u> <u>0.61</u>	23 (0.29) ₂
	<u>2001-2002</u>			<u>.01, <.01</u>	<u>0, 0</u>	<u>0, 0</u>	<u>8, 8</u>	<u>1.06,</u> <u>1.06</u>	<u>8 (0.75)</u>
Summer Southern NC	1996-2000	NA	Obs. Data, NCDMF Dealer Data	.00, .00, .01, .03, .03 ₂	0, 0, 0, 0, 0	0, 0, 0, 0, 0	0, 0, 0, 0, 0	NA	0 (NA)
	<u>2001-2002</u>			<u>.02, <.01</u>	<u>0, 0</u>	<u>0, 0</u>	<u>0, 0</u>	<u>NA</u>	<u>0 (NA)</u>
Winter NC mixed	1996-2000	NA	Obs. Data, NCDMF Dealer Data	.01, .01, .02, .02, .02 ₂	0, 0, 0, 0, 0	1, 0, 1, 2, 2 ₂	173, 211, 175, 196, 146 ₂	<u>0.46,</u> <u>0.46,</u> <u>0.46,</u> <u>0.46,</u> <u>0.46</u>	180 (0.21)
	<u>2001-2002</u>			<u>.01, .01</u>	<u>0, 0</u>	<u>0, 2</u>	<u>67, 50</u>	<u>0.45,</u> <u>0.45</u>	<u>58 (0.32)</u>
Total	<u>2001-2002 Only</u>								<u>77 (0.4626)</u>

NA=Not Available

¹ Observer data (Obs. data) are used to measure bycatch rates; the USA data are collected within the Northeast Fisheries Science Center (NEFSC) Sea Sampling Program. The NEFSC collects weighout (~~Weighout~~) landings data that are used as a measure of total effort for the USA sink gillnet fisheries.

² The observer coverage for the mid-Atlantic coastal sink gillnet fishery is measured in tons of fish landed.

³ The annual estimates of mortality from 1998-2000 were generated by applying one bycatch rate per management unit as estimated by a GLM (Palka and Rossman 2001). The CV does not account for variability that may exist in the unit of total landings (mt) from each year that are used to expand the

bycatch rate. Therefore, the CV is the same for all five annual estimates.

~~A sink gillnet fishery for American shad operates seasonally from Connecticut to Georgia, with nets being moved from coastal ocean waters into fresh water with the shad spawning migration. It has been considered likely that a few bottlenose dolphins are taken in this fishery each year (Read 1994) but no takes have been observed (NEFSC observer data). The portion of the fishery which operates along the South Carolina coast was sampled by observers during 1994 and 1995, and no fishery interactions were observed (McFee *et al.* 1996).⁴~~

The annual estimates of mortality from 2001-2002 were generated by applying the same method used in Palka and Rossman (2001). An new factor variable was added to the model to separate the time series of historical data (1996-2000) from data collected during the recent time period (2001-2002) (Rossman 2004, in review).

South Atlantic Shark Gillnet

~~The shark gillnet fishery operates in federal waters from off of southern Florida to southern Georgia. The fishery is defined by vessels using relatively large mesh nets (> 10 inches) and net lengths typically greater than 1500 feet. The fishery primarily uses drifting nets that are set overnight, however recently it has been employing a small number of shorter duration "strike" sets that encircle targeted schools of sharks. Since 1999, the Atlantic Large Whale Take Reduction Plan restricted the activities of the fishery to waters south of 27° 51' N latitude during the critical right whale season from 15 November – 31 March and mandated 100% observer coverage during this period. During the remainder of the year, these vessels generally operate north of Cape Canaveral, FL and there is little fishery observer coverage of the fleet.~~

~~The fishery potentially interacts with the Georgia, Northern Florida, and outside of this time and area has increased significantly in the last 2 years, and there was one observed mortality during summer months in fishing operations off of Cape Canaveral. All observed fishery takes are restricted to the Central Florida management unit* of coastal bottlenose dolphin. During an observer program in 1993 and 1994 and limited observer coverage during summer 1998, no takes of bottlenose dolphin were observed (Trent *et al.* 1997; Carlson and Lee, 2000). However, takes resulting in mortality were observed in the central Florida management unit during 1999 and 2000. Total bycatch mortality for this management unit has been estimated for 1999 and 2000 (Table 3) 1999-2002 following methods described in (Garrison 2001b2003) (Table 3).~~

Table 3. Summary of the 1999-2000² incidental mortality of bottlenose dolphins (*Tursiops truncatus*) by management unit in the driftnet fishery in federal waters off the coast of Florida. Data include years sampled (Years), number of vessels active within the fishery (Vessels), type of data used (Data Type), annual observer coverage (Observer Coverage), mortalities recorded by on-board observers (Observed Mortality), estimated annual mortality (Estimated Mortality), estimated CV of the annual mortality (Estimated CVs), and mean

annual mortality (CV in parentheses).

Seasonal Management Unit	Years	Vessels	Data Type ¹	Observer Coverage ²	Observed Serious Injury	Observed Mortality	Estimated Mortality	Estimated CVs	Mean Annual Mortality
Northern Florida	1999-2006	6	Obs. Data, SEFSC FVL	0.34, 0.29, 0.23, 0.07, 0.20	0, 0, 0, 0	0, 0, 0, 0	0, 0, 0, 0	NA	0
Central Florida	1999-2006	6	Obs. Data, SEFSC FVL	0.09, 0.24, 0.15, 0.42, 0.25	0, 0, 0, 0	4, 1, 4, 4, 1	12, 2, 4, 7	0.78, 1, 0, 1	246 (0.89)

NA=Not Available

¹ Observer data are used to estimate bycatch rates. The SEFSC Fishing Vessel Logbook (FVL) is used to estimate effort as total number of vessel trips per bottlenose dolphin management unit.

² Observer coverage in the central Florida management unit is largely restricted to the period between January - March south of 27° 51' N.

Beach Haul Seine

~~A beach seine fishery operates along northern North Carolina beaches targeting striped bass, mullet, spot, weakfish, sea trout, and bluefish. The fishery operates on the Outer Banks of North Carolina primarily in the spring (April through June) and fall (October through December). It uses two primary gear types: a "beach anchored gill net" and a "beach seine". Both systems utilize a small net anchored to the beach. The beach seine system also uses a bunt and a wash net that are attached to the beach and are in the surf (Steve *et al.* 2001). The North Carolina beach seine fishery has been observed since April 7, 1998 by the NMFS fisheries sampling program (observer program) based at the Northeast Fisheries Science Center. Through 2001, there were 101 sets observed during the winter season (Nov-Apr) and 65 sets observed during the summer season (May-Oct). There were no sets observed during the summer of 2001. A total of 2 coastal bottlenose dolphin takes were observed, in the mid-Atlantic beach haul seine fishery: 1 in May 1998 and 1 in December 2000. The beach seine observer data are currently being reviewed but estimates of mortality are not yet available.~~

Crab Pots

Between 1994 and 1998, 22 bottlenose dolphin carcasses (4.4 dolphins per year on average) recovered by the Stranding Network between North Carolina and Florida's Atlantic coast displayed evidence of possible interaction with a trap/pot fishery (i.e., rope and/or pots attached, or rope marks). Additionally, at least 5 dolphins were reported to be released alive (condition unknown) from blue crab traps/pots during this time period. ~~In recent years, reports of strandings with evidence of interactions between~~ During 2003, two bottlenose dolphins ~~and both recreational and commercial crab pot fisheries have been increasing in the Southeast Region (McFee and Brooks 1998). The increased reporting may result from increased effort towards documenting these marks or increases in mortality. were observed entangled in crab pot lines in South Carolina.~~

Virginia Pound Nets

~~Data from the Chesapeake Bay suggest that the likelihood of bottlenose dolphin entanglement in pound net leads may be affected by the mesh size of the lead net (Bellmund *et al.* 1997), but the information is not conclusive.~~ Stranding data for 1993-1997 document interactions between WNA coastal bottlenose dolphins and pound nets in Virginia. Two bottlenose dolphin carcasses were found entangled in the leads of pound nets in Virginia during 1993-1997, for an average of 0.4 bottlenose dolphin strandings per year. A third record of an entangled bottlenose dolphin in Virginia in 1997 may have been applicable to this fishery. This entanglement involved a bottlenose dolphin carcass found near a pound net with twisted line marks consistent with the twine in the nearby pound net lead rather than with monofilament gillnet gear. ~~Given that other sources of annual serious injury and mortality estimates (e.g., observer data) are not available, the stranding data (0.4 bottlenose dolphins per year) were used as a minimum estimate of annual serious injury and mortality and this fishery was classified as a Category II fishery in the 2001 List of Fisheries.~~

Shrimp Trawl

~~The shrimp trawl fishery operates from North Carolina through northern Florida virtually year around,~~

~~moving seasonally up and down the coast.~~ One bottlenose dolphin was recovered dead from a shrimp trawl in Georgia in 1995 (Southeast USA Marine Mammal Stranding Network unpublished data), and another was taken in 1996 near the mouth of Winyah Bay, SC, during a research survey. No other bottlenose dolphin mortality or serious injury has been previously reported to NMFS.

Menhaden Purse Seine

The Atlantic menhaden purse seine fishery ~~targets the Atlantic menhaden in Atlantic coastal waters. Smith (1999) summarized menhaden fishing patterns by the Virginia-North Carolina vessels from 1985-1996. Most of the catch and sets during that time occurred within three miles of the shore. Between 1994 and 1997, menhaden were processed at only three facilities, two in Reedville Beach, VA, and one in Beaufort, NC. Each of the Virginia facilities had a fleet of 9-10 vessels while the Beaufort facility is supported by 2-6 vessels. Since 1998, only one plant has operated in Virginia and the number of vessels has been reduced to ten in Virginia and two in North Carolina (Vaughan *et al.* 2001). The fishery moves seasonally, with most effort occurring off of North Carolina from November-January and moving northward to southern New England during warmer months. Menhaden purse seiners have~~historically reported an annual incidental take of 1 to 5 bottlenose dolphins (NMFS 1991, pp. 5-73). However, although no observer data are ~~not~~ available, and this information has not been updated for some time.

Other Mortality

From ~~1997-1999~~1997-2000, ~~995~~1,382 bottlenose dolphins were reported stranded along the Atlantic coast from New York to Florida (Table 4) (Hohn and Martone 2001; Hohn *et al.* 2001; Palka ~~*et al.* 2001~~, *et al.* 2001b, Northeast Regional Stranding Program, Southeast Regional Stranding Program). Between 2001-2003, 977 bottlenose dolphins stranded along the Atlantic coast from New York to Florida (Table 4) Of these, it was possible to determine whether or not a human interaction had occurred for ~~44~~59 (~~45~~7%) for the remainder it was not possible to make that determination. The proportion of carcasses~~Of those cases where a cause could be determined, 37% of the carcasses were~~ determined to have been involved in a human interaction ~~averaged 34%, but ranged widely from 11-12% in Delaware and Georgia to 49% and 53% in Virginia on average coastwide, and the majority of these were classified as fisheries interactions. However, this proportion ranged widely and was highest Virginia (71%) and North Carolina, respectively (43%).~~

The nearshore habitat occupied by the coastal morphotype is adjacent to areas of high human population and in the northern portion of its range is highly industrialized. The blubber of stranded dolphins examined during the 1987-88 mortality event contained anthropogenic contaminants in levels among the highest recorded for a cetacean (Geraci 1989). There are no estimates of indirect human-caused mortality resulting from pollution or habitat degradation.

Table 4. Summary of bottlenose dolphins stranded along the Atlantic Coast of the US. Total Stranded is further stratified into carcasses with signs of human interaction, those without any signs, and those where human interaction could not be determined (CBD). Human Interaction is stratified into stranded animals with line or nets marks or gear attached (Fishery Interaction), cleanly removed (cut off) appendages or cuts on the body (Mutilation), and other indications of human interactions such as propellor wounds. Florida strandings include only the Atlantic coast of Florida ~~but~~ extending to Key West.

STATE	1997 <u>2001</u>	1998 <u>2002</u>	1999 <u>2003</u>
New York Total Stranded	<u>1</u>	<u>1</u>	2
Human Interaction			
---- Fishery Interaction	<u>+0</u>	0	0
---- Mutilation	0	0	0
---- Other	0	0	0
No Human Interaction	<u>0</u>	0	<u>21</u>
CBD	1	1	<u>01</u>
New Jersey Total Stranded	<u>101</u>	11	<u>157</u>
Human Interaction			
---- Fishery Interaction	<u>01</u>	1	<u>31</u>
---- Mutilation	0	0	0
---- Other	0	<u>1</u>	<u>00</u>
No Human Interaction	<u>27</u>	<u>34</u>	<u>25</u>
CBD	<u>83</u>	<u>75</u>	10
Delaware Total Stranded	<u>146</u>	<u>13</u>	<u>818</u>
Human Interaction			
---- Fishery Interaction	<u>+0</u>	1	1
---- Mutilation	0	0	0
---- Other	<u>20</u>	<u>+0</u>	0
No Human Interaction	<u>43</u>	<u>08</u>	<u>413</u>
CBD	<u>73</u>	<u>64</u>	<u>134</u>
Maryland Total Stranded	<u>23</u>	<u>25</u>	<u>5 10</u>
Human Interaction			
---- Fishery Interaction	<u>0</u>	0	<u>0 1</u>
---- Mutilation	0	0	0
---- Other	0	0	0
No Human Interaction	1	<u>0 2</u>	<u>+8</u>
CBD	<u>+ 2</u>	3	<u>1</u>
Virginia Total Stranded	<u>4471</u>	<u>4268</u>	<u>5060</u>
Human Interaction			
---- Fishery Interaction	<u>+17</u>	<u>15</u>	<u>818 25</u>
---- Mutilation	<u>1</u>	<u>2</u>	<u>0</u>
---- Other	<u>0 1</u>	<u>4</u>	<u>0</u>
No Human Interaction	<u>+58</u>	<u>7</u>	<u>126</u>
CBD	<u>+844</u>	<u>+939</u>	<u>23</u>
STATE	1997 <u>2001</u>	1998 <u>2002</u>	1999 <u>2003</u>
N. Carolina Total Stranded	<u>12387</u>	<u>10494</u>	<u>69</u>
Human Interaction			
---- Fishery Interaction	<u>289</u>	<u>2313</u>	<u>2411</u>
---- Mutilation	<u>50</u>	<u>32</u>	<u>+0</u>
---- Other	+0	<u>2</u>	<u>0</u>
No Human Interaction	<u>21 16</u>	<u>+9 15</u>	<u>16</u>
CBD	<u>6862</u>	62	<u>50 42</u>
S. Carolina Total Stranded	<u>4169</u>	<u>4128</u>	<u>-345</u>
Human Interaction			
---- Fishery Interaction	<u>83</u>	4	<u>+ 3</u>
---- Mutilation	<u>20</u>	0	<u>+ 0</u>
---- Other	<u>3</u>	0	<u>+2 0</u>
No Human Interaction	<u>+523</u>	<u>13</u>	<u>+010 17</u>
CBD	<u>+640</u>	<u>11</u>	<u>2620 15</u>
Georgia Total Stranded	<u>23</u>	<u>181</u>	<u>26147</u>
Human Interaction			
---- Fishery Interaction	1	<u>+0</u>	<u>+0</u>
---- Mutilation	0	0	0
---- Other	<u>01</u>	0	0
No Human Interaction	<u>85</u>	<u>60</u>	<u>82</u>
CBD	<u>9 16</u>	<u>191</u>	<u>515</u>
Florida Total Stranded	<u>1048087</u> <u>101</u>	<u>82</u>	<u>74</u>
Human Interaction			
---- Fishery Interaction	<u>79</u>	<u>38</u>	<u>411</u>
---- Mutilation	0	0	0
---- Other	<u>01</u>	<u>2</u>	0
No Human Interaction	<u>3446</u>	<u>292850</u>	<u>21</u>
CBD	<u>6345</u>	<u>475522</u>	<u>42</u>
Total	<u>358372</u>	<u>317313</u>	<u>32029</u> <u>2</u>

STATUS OF STOCKS

The coastal migratory stock ~~is was~~ designated as depleted under the MMPA. From 1995-2001, NMFS recognized only a single migratory stock of coastal bottlenose dolphins in the WNA, and, ~~therefore~~, the entire stock was listed as depleted. The management units in this report now replace the single coastal migratory stock. A re-analysis of the depletion designation on a management unit basis needs to be undertaken. In the interim, because one or more of the management units may be depleted, all management units retain the depleted designation. In addition, mortality ~~in multiple units exceed PBR (Table 1). There are no rigorous results that would provide reliable information on current abundance relative to historical abundance exceeded PBR in the North Carolina winter mixed stocks during the period from 1996-2000 (Table 1).~~ All prior estimates cover only part of the range of management units spatially or temporally, include the offshore morphotype, or are otherwise compromised—~~P, therefore~~ population trends cannot be determined due to insufficient data. -

~~Over the past five years, estimated average annual mortality exceeded PBR in the mid-Atlantic gillnet fisheries for the northern migratory and northern NC management units during summer and for the NC mixed management units in winter (Tables 1 and 2).~~

~~The species is not listed as threatened or endangered under the Endangered Species Act, but because, as noted above, the stock is listed as depleted under the MMPA it is a the management units are strategic stock. This stock is also considered strategic under the MMPA because fishery-related mortality and serious injury exceed the potential biological removal level stocks due to the depleted listing under the MMPA.~~

REFERENCES

- Barco, S. G. and W. M. Swingle. 1996. Sighting patterns of coastal migratory bottlenose dolphins (*Tursiops truncatus*) in the nearshore waters of Virginia and North Carolina. Final Report to the Virginia Dept. of Environmental Quality, Coastal Resources Management Program through Grant #NA47OZ0287-01 from NOAA, Office of Ocean and Coastal Resource Management. 32 pp.
- Barlow, J., S. L. Swartz, T. C. Eagle, and P. R. Wade. 1995. U.S. Marine Mammal Stock Assessments: Guidelines for Preparation, Background and a Summary of the 1995 Assessments. ~~U.S. Dep. Commer., NOAA Tech. Memo. Technical Memorandum~~ NMFS-OPR-6, 73 pp.
- ~~Bellman~~ Bellman, S.A., Musick, J.A., Klinger, R.C., Byles, R.A., Keinath, J.A., and Barnard, D.E. 1997. Ecology of Sea Turtles in Virginia. Final Report to National Marine Fisheries Service Northeast Region. Contract number NA80FAC-00004. Virginia Institute of Marine Science, Gloucester Point, Virginia.
- ~~Blaylock, R. A. 1995. A pilot study to estimate abundance of the U.S. Atlantic coastal migratory bottlenose dolphin stock. NOAA Tech. Memo. NMFS-SEFSC-362, 9 pp. NMFS, Southeast Fisheries Science Center, Miami, FL.~~
- ~~Blaylock, R. A. and W. Hoggard. 1994. Preliminary estimates of bottlenose dolphin abundance in southern U.S. Atlantic and Gulf of Mexico continental shelf waters. NOAA Tech. Mem. NMFS-SEFSC-356, 10 pp. NMFS, Southeast Fisheries Science Center, Miami, FL.~~
- Buckland, S. T., D. R. Anderson, K. P. Burnham, and J. L. Laake. 1993. Distance sampling: estimating abundance of biological populations. Chapman and Hall, London. 446 pp.
- Carlson, J.K. and Lee, D. 2000. The directed shark drift gillnet fishery: Catch and bycatch 1998-1999. NMFS/SEFSC/Sustainable Fisheries Division #SFD-99/00-87.
- Cortese, N.A. 2000. Delineation of bottlenose dolphin populations in the western Atlantic Ocean using stable isotopes. Master's thesis, University of Virginia, 118 pp.
- Curry, B. E. 1997. Phylogenetic relationships among bottlenose dolphins (genus *Tursiops*) in a world-wide context. Ph.D. dissertation, Texas A&M University, Texas, USA. 138 pp.
- CETAP (Cetacean and Turtle Assessment Program). 1982. A Characterization of Marine Mammals and Turtles in the Mid- and North Atlantic Areas of the U.S. Outer Continental Shelf, Final Report, Contract AA551-CT8- 48, U.S. NTIS PB83-215855, Bureau of Land Mgmt, Wash., DC, 576 pp.
- ~~Garrison, L. 2001a. Seeking a hiatus in sightings for bottlenose dolphin during summer and winter aerial surveys. NMFS/SEFSC report prepared for the Bottlenose Dolphin Take Reduction Team.~~
- ~~Garrison, L.P. 2001b. Mortality estimate for Atlantic bottlenose dolphin in Protected species interactions with the directed shark gillnet fishery off Florida and Georgia: from 1999-2002. NMFS/SEFSC Internal report. Available from: Southeast Fisheries Science Center, 75 Virginia Beach Dr., Miami, FL 33149.~~

- Garrison, L.P., Rosel, P.E., Hohn, A.A., Baird, R. and W. Hoggard. 2003. Abundance of the coastal morphotype of bottlenose dolphin *Tursiops truncatus*, in U.S. continental shelf waters between New Jersey and Florida during winter and summer 2002. NMFS/SEFSC report prepared and reviewed for the Bottlenose Dolphin Take Reduction Team. NMFS, Available from: Southeast Fisheries Science Center, Miami, FL.
- ~~Garrison, L. and A. Hohn. 2001. Abundance Estimates for Atlantic Bottlenose Dolphin: Combining strip transect data and line transect abundance estimation. NMFS/SEFSC report prepared for the Bottlenose Dolphin Take Reduction Team. NMFS, Southeast Fisheries Science Center, Miami, FL. 75 Virginia Beach Dr., Miami, FL 33149.~~
- Garrison, L. P. and C. Yeung. 2001. Abundance Estimates for Atlantic Bottlenose Dolphin Stocks During Summer and Winter, 1995. NMFS/SEFSC report prepared and reviewed for the Bottlenose Dolphin Take Reduction Team. NMFS, Available from: Southeast Fisheries Science Center, 75 Virginia Beach Dr., Miami, FL 33149.
- Geraci, J. R. 1989. Clinical investigation of the 1987-88 mass mortality of bottlenose dolphins along the U.S. central and south Atlantic coast. Final Report to National Marine Fisheries Service, U.S. Navy, Office of Naval Research, and Marine Mammal Commission, 63 pp.
- Hoelzel, A.R., Potter, C.W., Best, P.B. 1998. Genetic differentiation between parapatric nearshore and offshore populations of the bottlenose dolphin. Proceedings of the Royal Society of London. 265: 1177-1183.
- Hohn, A. A. 1997. Design for a multiple-method approach to determine stock structure of bottlenose dolphins in the mid-Atlantic. NOAA Technical Memorandum NMFS-SEFSC-401, 22 pp. NMFS, Southeast Fisheries Science Center, Miami, FL.
- Hohn, A.A. and P. Martone. 2001. Characterization of bottlenose dolphin strandings in North Carolina, 1997-2000. NMFS/SEFSC ~~R~~report prepared and reviewed for the Bottlenose Dolphin Take Reduction Team. NMFS, Available from: Southeast Fisheries Science Center, 75 Virginia Beach Dr., Miami, FL 33149.
- Hohn, A.A., B. Mase, J Litz, W. McFee, and B. Zoodsma. 2001. Characterization of human-caused strandings of bottlenose dolphins along the Atlantic coast from South Carolina to Southern Florida, 1997-2000. NMFS/SEFSC ~~R~~report prepared and reviewed for the Bottlenose Dolphin Take Reduction Team. NMFS, Available from: Southeast Fisheries Science Center, 75 Virginia Beach Dr., Miami, FL 33149.
- Kenney, R. D. 1990. Bottlenose dolphins off the northeastern United States. Pages 369-386 in S. Leatherwood and R. R. Reeves (eds), The bottlenose dolphin, Academic Press, San Diego, 653 pp.
- McFee, W. E., D. L. Wolf, D. E. Parshley and P. A. Fair. 1996. Investigation of marine mammal entanglement associated with a seasonal coastal net fishery. NOAA Technical Memo. NMFS-SEFSC-38, 22 pp. + 8 Tables, 5 Figures, 6 Appendices. NMFS, Southeast Fisheries Science Center, Miami, FL.
- McFee, W. E. and W. Brooks, Jr. 1998. Fact finding meeting of marine mammal entanglement in the crab pot fishery: a summary. U.S. Fish and Wildlife Service. Unpublished Report.
- McLellan, W.M., A.S. Friedlaender, J.G. Mead, C.W. Potter, and D.A. Pabst. 2003. Analysing 25 years of bottlenose dolphin (*Tursiops truncatus*) strandings along the Atlantic coast of the USA: do historic records support the coastal migratory stock hypothesis ? Journal of Cetacean Research and Management 4: 297-304.
- NMFS. 1991. Proposed regime to govern the interactions between marine mammals and commercial fishing operations after October 1, 1993. Draft Environmental Impact Statement, June 1991.
- NMFS. 2001. Stock structure of coastal bottlenose dolphins along the Atlantic coast of the US. NMFS/SEFSC Report prepared for the Bottlenose Dolphin Take Reduction Team. NMFS, Available from: Southeast Fisheries Science Center, 75 Virginia Beach Dr., Miami, FL 33149.
- Odell, D. K. and E. D. Asper. 1990. Distribution and movements of freeze-branded bottlenose dolphins in the Indian and Banana Rivers, Florida. Pp. 515-540 in: S. Leatherwood and R. R. Reeves (eds), The bottlenose dolphin, Academic Press, San Diego, 653 pp.
- Palka, D., L. Garrison, A. Hohn, and C. Yeung. 2001a. Summary of abundance estimates and PBR for coastal *Tursiops* for waters between New York and Florida during 1995 to 2000. NMFS report.
- ~~Palka D., M. Rossman. 2001. Bycatch estimates of coastal bottlenose dolphin (*Tursiops truncatus*) in U.S. mid-Atlantic gillnet fisheries for 1996-2000. NOAA-NMFS-NEFSC Ref. Doc. 01-15; 77 p. NMFS/NEFSC Report prepared for the Bottlenose Dolphin Take Reduction Team. Available from: National Marine NMFS-Northeast Fisheries Service Science Center, 166 Water Street St., Woods Hole, MA 02543-1026 02543.~~

- Palka, D., F. Wenzel, D. Hartley, and M. Rossman. 2001^b. Summary of bottlenose dolphin strandings from New York to Virginia. NMFS/NEFSC Report prepared for the Bottlenose Dolphin Take Reduction Team. Available from: NMFS-Northeast Fisheries Science Center, 166 Water St., Woods Hole, MA 02543.
- Read, A. J. 1994. Interactions between cetaceans and gillnet and trap fisheries in the northwest Atlantic. Rep. int Whal. Commn. Special Issue 15: 133-147.
- Read, A.J., B. Foster, K. Urian, ~~D. Waples~~, B. Wilson, ~~A. and D. Pierce~~ Waples. 2001³. Mark-recapture survey Abundance of bottlenose dolphins ~~on~~ in the bays ~~and~~ sounds, and estuaries of ~~NC. Report to Sea Grant~~ North Carolina. Marine Mammal Science 19: 59-73.
- Rossman, M.C. and D.L. Palka. 2001. Bycatch estimates of coastal bottlenose dolphin (*Tursiops truncatus*) in the U.S. mid-Atlantic gillnet fisheries for 1996 to 2000.- Northeast Fisheries Science Center Reference Document 01-15. 77 pp.
- Sayigh, L., K. W. Urian, A. Bocconcelli, G. Jones, D. Koster, K. Halbrook and A. J. Read. 1997. Photo-identification of dolphins near Wilmington, NC. Abstract. Fifth Annual Atlantic Coastal Dolphin Conference.
- Scott, G. P., D. M. Burn, and L. J. Hansen. 1988. The dolphin die off: long term effects and recovery of the population. Proceedings: Oceans '88, IEEE Cat. No. 88-CH2585-8, Vol. 3: 819-823.
- Scott, M. D., R. S. Wells and A. B. Irvine. 1990. A long-term study of bottlenose dolphins on the west coast of Florida. Pp. 235-244 In: S. Leatherwood and R. R. Reeves. The Bottlenose Dolphin. Academic Press, San Diego. 653 pp.
- Smith, J.W. 1999. Distribution of Atlantic menhaden, *Brevoortia tyrannus*, purse-seine sets and catches from southern New England to North Carolina, 1985-96. U.S. Dept. of Commerce, ~~NOAA~~ Technical NOAA Technical Memorandum, NMFS 144. 22p.
- Steve, C., J. Gearhart, D. Borggaard, L. Sabo and A.A. Hohn. 2001. Characterization of North Carolina Commercial Fisheries with Occasional Interactions with Marine Mammals. NOAA Technical Memorandum NMFS-SEFSC-458. 60 pp.
- Torres, L.G., P.E. Rosel, C. D'Agrosa, and A.J. Read. 2003. Improving management of overlapping bottlenose dolphin ecotypes through spatial analysis and genetics. Marine Mammal Science 19:502-514.
- Trent, L., Parshley, D.E., and Carlson, J.K. 1997. Catch and bycatch in the shark drift gillnet fishery off Georgia and East Florida. Marine Fisheries Review 59(1): 19-28.
- Urian, K., A.A. Hohn, and L.J. Hansen. 1999. Status of the photo-identification catalogue of coastal bottlenose dolphins of the western north Atlantic: report of a workshop of catalogue contributors. NOAA Tech. Memo. NMFS-SEFSC-425.
- Vaughan, D.S., J.W. Smith, E.H. Williams, and M.H. Prager. 2001. Analyses on the status of the Atlantic menhaden stock. Report prepared for the Atlantic States Marine Fisheries Commission. 57 pp-
- Wade, P. R. and R. P. Angliss. 1997. Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop April 3-5, Seattle, Washington. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-12, 93 pp.
- Wang, K. R., P. M. Payne and V. G. Thayer. 1994. Coastal stock(s) of Atlantic bottlenose dolphin: status review and management: Proceedings and recommendations from a workshop held in Beaufort, North Carolina, 13-14 September 1993. ~~U.S. Department of Commerce, NOAA, NMFS Tech. Mem~~ Technical Memorandum NMFS-OPR-4, 120 pp.
- Weller, D. W. 1998. Global and regional variation in the biology and behavior of bottlenose dolphins. Ph.D. dissertation, Texas A&M University, Galveston, TX. 142 pp.
- Wells, R. S., K. W. Urian, A. J. Read, M. K. Bassos, W. J. Carr and M. D. Scott. 1996. Low-level monitoring of bottlenose dolphins (*Tursiops truncatus*), in Tampa Bay, Florida: 1998-1993. NOAA ~~Tech. Mem.~~ Technical Memorandum NMFS- SEFSC-385, 25 pp. + 6 Tables, 8 Figures, and 4 Appendices.
- Zolman, E. S. 1996. Residency patterns, relative abundance and population ecology of bottlenose dolphins (*Tursiops truncatus*) in the Stono River Estuary, Charleston County, South Carolina. Master of Science thesis. University of Charleston, South Carolina, USA. 128 pp.

BOTTLENOSE DOLPHIN (*Tursiops truncatus*): Western North Atlantic Offshore Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

There are two hematologically morphologically and morphologically genetically distinct bottlenose dolphin ecotypes morphotypes (Duffield *et al.* 1983; Duffield 1986) which correspond to a shallow water ecotype and a deep water ecotype; both ecotypes have been shown to described as the coastal and offshore forms. Both inhabit waters in the western North Atlantic Ocean (Hersh and Duffield 1990; Mead and Potter 1995; Curry and Smith 1997)– along the U.S. Atlantic coast. The offshore and nearshore ecotypes are genetically distinct using both mitochondrial and nuclear markers (Hoelzel *et al.* 1998). The offshore form is distributed primarily along the outer continental shelf and continental slope in the Northwest Atlantic Ocean.

Bottlenose dolphins which had stranded alive in the western North Atlantic in areas with direct access to deep oceanic waters had hemoglobin profiles which matched that of the deep, cold water ecotype offshore morphotype (Hersh and Duffield 1990). Hersh and Duffield (1990) also described morphological differences between the deep, cold water ecotype offshore morphotype dolphins and dolphins with hematological profiles matching the shallow, warm water ecotype coastal morphotype which had stranded in the Indian/Banana River in Florida. Based on the distribution of sightings during ship-based surveys (Figure 1) and survey personnel observations (NMFS unpublished data); the western North Atlantic offshore stock is believed to consist North of Cape Hatteras, there is clear separation of the two morphotypes along the bathymetry gradient during summer months. Aerial surveys flown during 1979-1981 indicated a concentration of bottlenose dolphins in waters < 25 m corresponding to the hematologically and morphologically distinct deep, cold water ecotype.

Extensive aerial surveys in 1979-1981 indicated that the stock extended along the entire coastal morphotype, and an area of high abundance along the shelf break corresponding to the offshore stock (CETAP 1982; Kenney 1990). Biopsy tissue sampling and genetic analysis demonstrated that bottlenose dolphins concentrated close to shore were of the coastal morphotype, while those in waters > 40 m depth were from the coastal morphotype (Garrison *et al.* 2003). However, during winter months and south of Cape Hatteras, NC the range of the coastal and offshore morphotypes overlap to some degree. Torres *et al.* (2003) found a statistically significant break in the distribution of the ecotypes at 34 km from shore based upon the genetic analysis of tissue samples collected in nearshore and offshore waters. The offshore morphotype was found exclusively seaward of 34 km and in waters deeper than 34 m. Within 7.5 km of shore, all animals were of the coastal morphotype. Systematic biopsy collection surveys were conducted coastwide during the summer and winter between 2001-2003 to evaluate the degree of spatial overlap between the two morphotypes. Over the continental shelf south of Cape Hatteras, NC the two




Figure 1. Distribution of bottlenose dolphin sightings from NEFSC and SEFSC shipboard and aerial surveys during the summer in +1990- +1998 1998, 1999, and 2004. Isobaths are at 100

morphotypes overlap spatially, and the probability of a sampled group being from the offshore morphotype increased with increasing depth based upon a logistic regression analysis. Offshore morphotype animals have been sampled as close as 7.3 km from shore in water depths of 13 m (Garrison *et al.* 2003).

Seasonally, bottlenose dolphins occur over outer continental shelf and inner slope waters as far north as Georges Bank (Figure 1; CETAP 1982; Kenney 1990). Sightings occurred along the continental shelf break from Georges Bank to Cape Hatteras during spring and summer (CETAP 1982; Kenney 1990). The distribution of sightings contracted towards the south in the fall and the central portion of the survey area was almost devoid of sightings in the winter, although there were still sightings as far north as the southern edge of Georges Bank. The offshore stock is concentrated along the continental shelf break in waters of depths >25 m and extends beyond the continental shelf into continental slope waters in lower concentration (Figure 1) consistent with Kenney 1990. In Canadian waters, bottlenose dolphins have occasionally been sighted on the Scotian Shelf, particularly in the Gully (Gowans and Whitehead 1995; NMFS unpublished data). Recent information from Wells *et al.* (1999) indicates that the range of the offshore bottlenose dolphin may include waters beyond the continental slope and that offshore bottlenose dolphins may move between the Gulf of Mexico and the Atlantic. Dolphins with characteristics of the offshore type have been stranded as far south as the Florida Keys, but there are no abundance or distribution estimates available for this stock in USA Exclusive Economic Zone (EEZ) waters south of Cape Hatteras.

POPULATION SIZE

An abundance of 16,689 (CV=0.32) ~~for~~ bottlenose dolphins was estimated from a line-transect sighting survey conducted during July 6 to September 6, 1998, by a ship and plane that surveyed 15,900 km of track line in waters north of Maryland (38° N) (Figure 1; Palka *et al.*, in review). Shipboard data were analyzed using the modified direct duplicate method (Palka 1995) that accounts for school size bias and $g(0)$, the probability of detecting a group on the track line. Aerial data were not corrected for $g(0)$.

An abundance of 13,944 ~~085~~ (CV=0.3840) for bottlenose dolphins was estimated from a shipboard line transect sighting line-transect survey conducted between 8 July and 17 August 1998 that surveyed 54,570 ~~570163~~ km of track line in waters south of Maryland (38°N) (Figure Fig. 1; Mullin in review and Fulling 2003). Abundance estimates were made using the program DISTANCE (Buckland *et al.* ~~1993~~ 2001; ~~Laake Thomas~~ et al. ~~1993~~ 38) where school size bias and ship attraction were accounted for.

The best available During the summer (June - July) of 2002, aerial surveys were conducted along the U.S. Atlantic coast between Florida and New Jersey. A total of 6,734 km of trackline were completed during the summer survey between Sandy Hook, NJ to Ft. Pierce, FL. The abundance of bottlenose dolphins in survey strata were calculated using line transect methods and distance analysis, and the direct duplicate estimator was used to account for visibility bias (Buckland *et al.* 1993; Palka 1995). These estimates were further partitioned between the coastal and offshore morphotypes based upon the results of the logistic regression models and spatial analyses described above. A parametric bootstrap approach was used to incorporate the uncertainty in the logistic regression models into the overall uncertainty in the abundance estimate for offshore bottlenose dolphins (Garrison *et al.* 2003). The resulting coastwide abundance estimate for the offshore morphotype in waters < 40 m depth was 26,849 (CV = 0.193).

An abundance of 10,549 (CV = 0.56) for offshore morphotype bottlenose dolphins was estimated from a line transect sighting survey conducted during June 12 to August 4, 2004 by a ship and plane that surveyed 10,761 km of track line in waters north of 38° N Figure 1; Palka unpubl.). Shipboard data were collected using the two independent team line transect method and analyzed using the modified direct duplicate method (Palka 1995) accounting for biases due to school size and other potential covariates, reactive movements (Palka and Hammond 2001), and $g(0)$, the probability of detecting a group on the track line. Aerial data were collected using the Hiby circle-back line transect method (Hiby 1999) and analyzed accounting for $g(0)$ and biases due to school size and other potential covariates (Figure 1; Palka unpubl.).

A survey of the U.S. Atlantic outer continental shelf and continental slope (water depths ≥ 50 m) between 27.5 – 38°N latitude was conducted during June-August, 2004. The survey employed two independent visual teams searching with 50x bigeye binoculars. Survey effort was stratified to include increased effort along the continental shelf break and Gulf stream front in the mid-Atlantic. The survey included 5,659 km of trackline, and there were a total of 473 cetacean sightings. Sightings were most frequent in waters North of Cape Hatteras, North Carolina along the shelf break. Data were analyzed to correct for visibility bias ($g(0)$) and group-size bias employing line transect distance analysis and the direct duplicate estimator (Palka, 1995; Buckland *et al.*, 2001). The resulting abundance estimate for offshore morphotype bottlenose dolphins between Florida and Maryland was 44,953 (CV = 0.26).

The best available estimate for offshore morphotype bottlenose dolphins is the sum of the estimates from the ~~two 1998 USA Atlantic surveys, 30,633 (CV=0.25), where the estimate from the northern USA Atlantic is 16,689 (CV=0.32) and from the southern USA Atlantic is 13,944 (CV=0.38);~~ summer 2002 aerial survey covering the continental shelf, the summer 2004 vessel survey South of Maryland, and the summer 2004 vessel and aircraft surveys North of Maryland. This joint estimate ~~is considered best because together these two surveys have the most provides~~ complete coverage of the ~~species' habitat; offshore morphotype habitat from Florida to Georges Bank during summer months.~~ The combined abundance estimate from these surveys is 82,351 (CV = 0.17).

Minimum Population Estimate

The minimum population estimate is the lower limit of the two-tailed 60% confidence interval of the log-normally distributed best abundance estimate. This is equivalent to the 20th percentile of the log-normal distribution as specified by Wade and Angliss (1997). ~~The best estimate of abundance for offshore bottlenose dolphins is 30,633 (CV=0.25).~~ The minimum population estimate for ~~the~~ western North Atlantic offshore bottlenose dolphin is 24,89771,382.

Current Population Trend

The data are insufficient to determine population trends. Previous estimates cannot be applied to this process because previous survey coverage of the species' habitat was incomplete.

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Current and maximum net productivity rates are unknown for this stock. For purposes of this assessment, the maximum net productivity rate was assumed to be 0.04. This value is based on theoretical modeling showing that cetacean populations may not grow at rates much greater than 4% given the constraints of their reproductive life history (Barlow *et al.* 1995).

POTENTIAL BIOLOGICAL REMOVAL

Potential ~~B~~biological ~~R~~emoval (PBR) is the product of minimum population size, one-half the maximum productivity rate, and a "recovery" factor (MMPA Sec. 3. 16 U.S.C. 1362; Wade and Angliss 1997). The minimum population size for offshore bottlenose dolphins is 24,89771,382. The maximum productivity rate is 0.04, the default value for cetaceans. The "recovery" factor, which accounts for endangered, depleted, threatened stocks, or stocks of unknown status relative to optimum sustainable population (OSP) is assumed to be 0.5 because this stock is of unknown status. PBR for the western North Atlantic offshore bottlenose dolphin is 249714.

ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

Total estimated mean annual fishery-related mortality ~~to~~for this stock during ~~1996-2000~~1998-2002 was 27 (CV=1.12) bottlenose dolphins.

Fishery Information

~~There was no documentation of marine mammal mortality or serious injury in distant-water fleet (DWF) activities off the northeast coast of the USA prior to 1977. A fisheries observer program which recorded fishery data and information on incidental bycatch of marine mammals was established with implementation of the Magnuson Fisheries Conservation and Management Act (MFCMA) in 1977. DWF effort in the US Atlantic EEZ under MFCMA was directed primarily towards Atlantic mackerel and squid. An average of 120 different foreign vessels per year (range 102-161) operated within the US Atlantic EEZ from 1977 through 1982. In 1982, the first year that NMFS Northeast Regional Observer Program assumed responsibility for observer coverage of the longline vessels, there were 112 different foreign vessels, eighteen (16%) of which were Japanese tuna longline vessels operating along the USA east coast. Between 1983 and 1991, the number of foreign fishing vessels operating within the US Atlantic EEZ each year declined from 67 to 9. Between 1983 and 1988, the numbers of DWF vessels included 3, 5, 7, 6, 8, and 8, respectively, Japanese longline vessels. Observer coverage on DWF vessels was 25-35% during 1977-82, and increased to 58%, 86%, 95%, and 98%, respectively, in 1983-86. From 1987-91, 100% observer coverage was maintained. Foreign fishing operations for squid ceased at the end of the 1986 fishing season and for mackerel at the end of the 1991 season. Observers in this program recorded 9 bottlenose dolphin mortalities in foreign-fishing activities during 1977-1988 (Waring *et al.* 1990). Seven takes occurred in the mackerel fishery, and 1 bottlenose dolphin each was caught in both the squid and hake trawl fisheries.~~

~~—————Data on current incidental takes in USA fisheries are available from several sources. In 1986, NMFS established a mandatory self-reported fisheries information system for large pelagic fisheries. Data files are maintained at the Southeast Fisheries Science Center (SEFSC). The Northeast Fisheries Science Center (NEFSC) Sea Sampling Observer Program was initiated in 1989, and since that year several fisheries have been covered by the program. In late 1992 and in 1993, the SEFSC provided observer coverage of pelagic longline vessels fishing off the Grand Banks (Tail of the Banks) and provides observer coverage of vessels fishing south of Cape Hatteras.~~

Fisheries Information

Bycatch has been observed by NMFS Sea Samplers in the pelagic drift gillnet, pelagic pair trawl, New England multispecies sink gillnet, North Atlantic bottom trawl ~~fisheries~~, and pelagic longline fisheries.

Pelagic Longline

The pelagic longline fishery operates in the ~~USA~~U.S. Atlantic (including Caribbean) and Gulf of Mexico EEZ (SEFSC unpublished data). Interactions between the pelagic longline fishery and bottlenose dolphins have been reported; however, a vessel may fish in more than one statistical reporting area and it is not possible to separate estimates of fishing effort other than to subtract Gulf of Mexico effort from Atlantic fishing effort, which includes the Caribbean Sea. ~~Total effort, excluding the Gulf of Mexico, for the pelagic longline fishery, based on mandatory self-reported fisheries information, was 11,279 sets in 1991, 9,869 sets in 1992, 9,862 sets in 1993, 9,481 sets in 1994, 10,129 sets in 1995, 9,885 sets in 1996, 8,023 sets in 1997, 6,765 in 1998, 6770 sets in 1999, and 6,582 sets in 2000 (Cramer 1994; Scott and Brown 1997; Johnson *et al.* 1999; Yeung 1999a; and Yeung, 2001). Since 1992, this fishery has been monitored with about 5% observer coverage, in terms of trips observed, within every statistical reporting area within the US Atlantic EEZ and beyond. Off the USA Atlantic coast, the fishery has been observed from January to March off Cape Hatteras, in May and June in the entire mid-Atlantic, and in July through December in the mid-Atlantic Bight and off Nova Scotia. Yeung (1999a) provides estimates of mortality for 1998, based on a treatment which includes seriously injured animals as mortalities, following guidelines proposed by the NOAA Fisheries Office of Protected Resources (Federal Register Docket No., I.D. 051398C). Yeung (1999b) provides revised estimates of mortality for 1993-1997. These estimates, which treat seriously injured animals as mortalities, replace the 1993-1997 estimates which were based on a revised analysis of the observed incidental take and self-reported incidental take and effort data (Johnson *et al.* 1999), and which replaced previous estimates for the 1992-1993 (Cramer 1994) and 1994-1995 periods (Scott and Brown 1997). Most of the estimated marine mammal bycatch was from US Atlantic EEZ waters between South Carolina and Cape Cod (Johnson *et al.* 1999). During 1993-1998, in waters not including the Gulf of Mexico, 1 bottlenose dolphin was caught and released alive during 1993, and 1 was caught and released alive during 1998. No bottlenose dolphins have been observed taken in the pelagic longline fishery since 1998 (Garrison, 2003; Garrison and Richards, 2004; Yeung, 1999).~~

Pelagic Drift Gillnet

The estimated total number of hauls in the pelagic drift gillnet fishery increased from 714 in 1989 to 1,144 in 1990; thereafter, with the introduction of quotas, effort was severely reduced. The estimated number of hauls in 1991, 1992, 1993, 1994, 1995, 1996, and 1998 were 233, 243, 232, 197, 164, 149, and 113, respectively. In 1996 and 1997, NMFS issued management regulations which prohibited the operation of this fishery in 1997. Further, in January 1999 NMFS issued a Final Rule to prohibit the use of driftnets (*i.e.*, permanent closure) in the North Atlantic swordfish fishery (50 CFR Part 630). Fifty-nine different vessels participated in this fishery at one time or another between 1989 and 1993. Since 1994, between 10-13 vessels have participated in the fishery (Table 1). Observer coverage, expressed as percent of sets observed, was 8% in 1989, 6% in 1990, 20% in 1991, 40% in 1992, 42% in 1993, 87% in 1994, 99% in 1995, 64% in 1996, no fishery in 1997, and 99% in 1998.

~~—————Effort was concentrated along the southern edge of Georges Bank and off Cape Hatteras. Examination of the species composition of the catch and locations of the fishery throughout the year, suggested that the pelagic drift gillnet fishery be stratified into two strata, a southern or winter stratum, and a northern or summer stratum. Estimates of the total bycatch, from 1989 to 1993, were obtained using the aggregated (pooled 1989-1993) catch rates, by strata (Northridge 1996). Estimates of total annual bycatch for 1994 and 1995 were estimated from the sum of the observed caught and the product of the average bycatch per haul and the number of unobserved hauls as recorded in self-reported fisheries information. Variances were estimated using bootstrap re-sampling techniques. Sixty bottlenose dolphin mortalities have been observed between 1989 and 1998. Estimated bottlenose dolphin mortalities (CV in parentheses) extrapolated for each year were 72 in 1989 (0.18), 115 in 1990 (0.18), 26 in 1991 (0.15), 28 in~~

1992 (0.10), 22 in 1993 (0.13), 14 in 1994 (0.04), 5 in 1995 (0), 0 in 1996, and 3 in 1998 (0). Since this fishery no longer exists, it has been excluded from Table 1.

Pelagic Pair Trawl

Effort in the pelagic pair trawl fishery increased during the period 1989 to 1993, from zero hauls in 1989 and 1990, to an estimated 171 hauls in 1991, and then to an estimated 536 hauls in 1992, 586 in 1993, 407 in 1994, and 440 in 1995, respectively. This fishery ceased operations in 1996, when NMFS rejected a petition to consider pair trawl gear as an authorized gear type in Atlantic tunas fishery. The fishery operated from August-November in 1991, from June-November in 1992, from June-October in 1993 (Northridge 1996), and from mid-summer to November in 1994 and 1995. Sea sampling began in October 1992 (Gerrior *et al.* 1994), and 48 sets (9% of the total) were sampled in that season, 102 hauls (17% of the total) were sampled in 1993. In 1994 and 1995, 52% and 55%, respectively, of the sets were observed. Nineteen vessels have operated in this fishery. The fishery extends from 35°N to 41°N, and from 69°W to 72°W. Approximately 50% of the total effort was within a one degree square at 39°N, 72°W, around Hudson Canyon. Examination of the 1991-1993 locations and species composition of the bycatch, showed little seasonal change for the six months of operation and did not warrant any seasonal or areal stratification of this fishery (Northridge 1996). Thirty-two bottlenose dolphin mortalities were observed between 1991 and 1995. Estimated annual fishery-related mortality (CV in parentheses) was 13 dolphins in 1991 (0.52), 73 in 1992 (0.49), 85 in 1993 (0.41), 4 in 1994 (0.40) and 17 in 1995 (0.26). Since this fishery no longer exists, it has been excluded from Table 1. During the 1994 and 1995 experimental fishing seasons, fishing gear experiments were conducted to collect data on environmental parameters, gear behavior, and gear handling practices to evaluate factors affecting catch and bycatch (Goudey 1995, 1996). Results of these studies have been presented at Offshore Cetacean Take Reduction Team Meetings.

North Atlantic Bottom Trawl:

~~Vessels in the North Atlantic bottom trawl fishery, a Category III fishery under the MMPA, were observed in order to meet fishery management needs, rather than marine mammal management needs. An average of 970 (CV=0.04) vessels (full and part time) participated annually in the fishery during 1989-1993. The fishery is active in New England waters in all seasons.~~

One bottlenose dolphin mortality was documented in 1991 and the total estimated mortality in this fishery in 1991 was 91 (CV=0.97). Since 1992 there were no bottlenose dolphin mortalities observed in this fishery.

Squid, Mackerel and Butterfish:

~~The mid-Atlantic mackerel and squid trawl fisheries were combined into the Atlantic squid, mackerel and butterfish trawl fishery in 1996. These fisheries operate seasonally, principally in the USA mid-Atlantic and southern New England continental shelf region. The mackerel trawl fishery was classified as a Category H fishery since 1990 and the squid fishery was originally classified as a Category H fishery in 1990, but was reclassified as a Category III fishery in 1992. The combined fishery has been proposed for classification as a Category H fishery. In 1996, mackerel, squid, and butterfish trawl fisheries were combined into the Atlantic squid, mackerel, and butterfish trawl fishery, and maintained a Category H classification.~~

Although there were reports of bottlenose dolphin mortalities in the foreign fishery during 1977-1988, there were no fishery-related mortalities of bottlenose dolphins reported in the self-reported fisheries information from the mackerel trawl fishery ~~between~~during 1990-1992.

New England Multispecies Sink Gillnet

In 1993, there were approximately 349 full and part-time vessels in the Northeast multispecies sink gillnet fishery, which covered the Gulf of Maine and southern New England (Table 1). An additional 187 vessels were reported to occasionally fish with gillnets in the Gulf of Maine for bait or personal use; however these vessels were not covered by the observer program (Walden 1996) and their fishing effort was not used in estimating mortality. Observer coverage in terms of trips has been 1%, 6%, 7%, 5%, 7%, 5%, 4%, 6%, 5%, 6%, and 6% for 1990 to 2000 respectively. The fishery has been observed in the Gulf of Maine and in Southern New England. In 2000, ~~t~~The first observed mortality of bottlenose dolphins was recorded in 2000. This was genetically identified as an offshore, deep-water ecotype. The estimated annual fishery-related serious injury and mortality attributable to this fishery (CV in parentheses) was 0 from 1996-1999, and 132 (CV=1.16) in 2000. There have been no observed bottlenose dolphin mortalities since 2000 in this fishery (Table 1).-

Mid-Atlantic Coastal Gillnet

Observer coverage of the USA Atlantic coastal gillnet fishery was initiated by the NEFSC Sea Sampling program in July 1993, and from July to December 1993, 20 trips were observed. During 1994 and 1995, 221 and 382 trips were observed, respectively. This fishery, which extends from North Carolina to New York, is actually a combination of small vessel fisheries that target a variety of fish species, some of which operate right off the beach. The number of vessels in this fishery is unknown, because records which are held by both state and federal agencies have not been centralized and standardized. Observer coverage, expressed as percent of tons of fish landed, was 5%, 4%, 3%, 5%, 2%, and 2% for 1995, 1996, 1997, 1998, 1999 and 2000, respectively (Table 1).

Bottlenose dolphins were only reported during the trips in 1998, when 1 mortality was observed as a result of this fishery. Though this dolphin was not genetically identified, it is being treated as an offshore, deep-water ecotype because it was caught in the offshore habitat and statistical analyses of all biopsied bottlenose dolphins caught in this offshore habitat indicate this animal has a high probability of being the offshore ecotype. Observed effort was concentrated off ~~NJ~~New Jersey and scattered between ~~DE~~Delaware and ~~NC~~North Carolina from 1 to 50 miles off the beach. All bycatches were documented during January to April. Using the observed takes, the estimated annual mortality attributed to this fishery was 0 in 1995 through 1997, 4 (CV=0.7) in 1998, and 0 from 1999 through 2000 (Table 1). A bottlenose dolphin was captured in the region of overlap over the continental shelf in the mid-Atlantic gillnet fishery during May, 2001. Mortality estimates have not been developed for the offshore morphotype during 2001-2002 due to the uncertainties associated with the relative distribution of the two morphotypes.

Table 1. Summary of the incidental mortality of bottlenose dolphins (*Tursiops truncatus*) by commercial fishery including the years sampled (Years), the number of vessels active within the fishery (Vessels), the type of data used (Data Type), the annual observer coverage (Observer Coverage), the mortalities recorded by on-board observers (Observed Mortality), the estimated annual mortality (Estimated Mortality), the estimated CV of the annual mortality (Estimated CVs) and the mean annual mortality (CV in parentheses).

Fishery	Years	Vessels	Data Type ¹	Observer Coverage ²	Observed Serious Injury	Observed Mortality	Estimated Mortality	Estimated CVs	Mean Annual Mortality
New England Multisp. Sink Gillnet	96-00 <u>99-03</u>	301	Obs. Data Dealer Reports, Logbooks	.04, .06, .05, .06, .06, .04, .02, .03	0, 0, 0, 0, 0	0, 1, 0, 0, <u>0</u>	0, 132, 0, <u>0, 0</u>	0, 1.16, 0, <u>0,</u> 0, 0; 1320, 0, 0, 0 1.16	26 (1.16)
mid-Atlantic Coastal Gillnet	96-00 <u>99-03</u>	Unk ³	Obs. Data Dealer Reports	.04, .03, .05, .02, .02, .02, .01, .01	0, 0, 0, 0, 0	0, 0, 1, 0, 0, 0,	4, 0, 00, 0NA, <u>NA</u>	0.7, 0, 0, NA, <u>NA</u>	1 (0.7)
TOTAL									27 (1.12)

¹ Observer data (Obs. Data) are used to measure bycatch rates, and the data are collected within the Northeast Fisheries Science Center (NEFSC) Sea Sampling Program. Mandatory logbook (logbook) data collected by the Southeast Fisheries Science Center (SEFSC) are used to measure total effort for the pelagic drift gillnet fishery. The NEFSC collects landings data (Dealer Reports), and total landings are used as a measure of total effort for the gillnet fisheries. Mandatory vessel trip reports (Logbook) data are used to determine the spatial distribution of fishing effort in the Northeast multispecies sink gillnet fishery.

² Observer coverage of the Northeast multispecies sink gillnet fishery is measured as the percentage of trips observed. Observer coverage of the mid-Atlantic coastal gillnet fishery is measured as the percentage of tons of fish landed.

³ Number of vessels is not known.

Other Mortality

Bottlenose dolphins are one of the most frequently stranded small cetaceans along the Atlantic coast. Many of the animals show signs of human interaction (*i.e.*, net marks, mutilation, etc.). The estimated number of animals that represent the offshore stock is presently under evaluation.

STATUS OF STOCK

The status of this stock relative to OSP in the U.S. Atlantic EEZ is unknown. The western North Atlantic offshore bottlenose dolphin is not listed as threatened or endangered under the Endangered Species Act. There are insufficient data to determine the population trends for this species. This level is Average 1998-2002 annual fishery-related mortality and serious injury does not exceed the PBR therefore this is not a strategic stock. The total fishery-related mortality and serious injury for this stock is less than 10% of the calculated PBR and, therefore, can be considered to be insignificant and approaching zero mortality and serious injury rate. Average 1994-1998 annual fishery-related mortality and serious injury does not exceed the PBR; therefore, this is not a strategic stock.

REFERENCES

- Barlow, J., S. L. Swartz, T. C. Eagle and P. R. Wade. 1995. U.S. Marine Mammal Stock Assessments: Guidelines for Preparation, Background, and a Summary of the 1995 Assessments. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-6, 73 pp.
- Buckland, S. T., D. R. Anderson, K. P. Burnham and J. L. Laake, D. Distance Distance Borchers and L. Thomas. 2001. Introduction to distance sampling: Estimating abundance of biological populations. Chapman and Hall, New York Oxford University Press, 446-446432 pp.
- CETAP (Cetacean and Turtle Assessment Program). 1982. A Characterization of Marine Mammals and Turtles in the Mid- and North Atlantic Areas of the U.S. Outer Continental Shelf, Final Report, Contract AA551-CT8-48, U.S. NTIS PB83-215855, Bureau of Land Management, Washington, DC, 576 pp.
- Cramer J. 1994. Large pelagic logbook newsletter - 1993. NOAA Tech. Mem. NMFS-SEFSC-352, 19 pp. NMFS, Southeast Fisheries Science Center, Miami, FL.
- Curry, B. E. and J. Smith. 1997. Phylogeographic structure of the bottlenose dolphin (*Tursiops truncatus*): stock identification and implications for management. Pp. 327-247. in: A. E. Dizon, S. J. Chivers and W. F. Perrin (eds. editors), Molecular genetics of marine mammals. Spec. Publ. 3 Society for Marine Mammalogy.
- Duffield, D. A. 1986. Investigation of genetic variability in stocks of the bottlenose dolphin (*Tursiops truncatus*). Final report to the NMFS/SEFSC, Contract No. NA83-GA-00036, 53 pp.
- Duffield, D. A., S. H. Ridgway and L. H. Cornell. 1983. Hematology distinguishes coastal and offshore forms of dolphins (*Tursiops*). *Can. J. Zool.* 61: 930-933.
- Gerrior Garrison, P., A. S. Williams and D. J. Christensen. 1994. Observations of the 1992 U.S. pelagic pair trawl fishery in the Northwest Atlantic. U.S. Mar. Fish. Rev. 56(3): 24-27.
- Goudey, C.A. 1995. The 1994 experimental pair trawl fishery for tuna in the northwest Atlantic, Massachusetts Institute of Technology, Sea Grant, MITSG 95-6, Cambridge, MA. 10 pp.
- Goudey, C.A. 1996. The 1995 experimental pair trawl fishery for tuna in the northwest Atlantic, Massachusetts Institute of Technology, Sea Grant, MITSG 95-6, Cambridge, MA. 13 pp.
- L.P. 2003. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2001-2002. NOAA Technical Memorandum NMFS-SEFSC-515, 51 pp.
- Garrison, L. P. and P. M. Richards. 2004. Estimated Bycatch of Marine Mammals and Turtles in the U.S. Atlantic Pelagic Longline Fleet During 2003. NOAA Technical Memorandum NMFS-SEFSC-527, 57 p.
- Gowans S. and H. Whitehead. 1995. Distribution and habitat partitioning by small odontocetes in the Gully, a submarine canyon on the Scotian Shelf. *Can. J. Zool.* 73: 1599-1608.
- Hersh, S. L. and D. A. Duffield. 1990. Distinction between northwest Atlantic offshore and coastal bottlenose dolphins based on hemoglobin profile and morphometry. Pp. Pages 129-139. in: S. Leatherwood and R. Reeves (eds. editors), The bottlenose dolphin, Academic Press, San Diego, 653 pp.
- Johnson, D. R., C. A. Brown and C. Yeung. 1999. Estimates of marine mammal and marine turtle catch by the U.S. Atlantic pelagic longline fleet in 1992-1997. NOAA Tech. Mem. NOAA-SEFSC-418, 70 pp. NMFS, Southeast Fisheries Science Center, Miami, FL.
- Hiby, L. 1999. The objective identification of duplicate sightings in aerial survey for porpoise. Pages 179-189 in: Garner et al. (eds.). Marine Mammal Survey and Assessment Methods. Balkema, Rotterdam.
- Hoelzel, A. R., C. W. Potter and P. B. Best. 1998. Genetic differentiation between parapatric 'nearshore' and

- 'offshore' populations of the bottlenose dolphin. *Proc. R. Soc. Lond. B* 265:1177-1183.
- Kenney, R. D. 1990. Bottlenose dolphins off the northeastern United States. Pages 369-386, in In: S. Leatherwood and R. R. Reeves (~~eds~~editors), *The bottlenose dolphin*, Academic Press, San Diego, 653 pp.
- ~~Laake, J. L., S. T. Buckland, D. R. Anderson and K. P. Burnham. 1993. DISTANCE user's guide, V2.0. Colorado Cooperative Fish & Wildlife Research Unit, Colorado State University, Ft. Collins, Colorado, 72 pp.~~
- Mead, J. G. and C. W. Potter. 1995. Recognizing two populations for the bottlenose dolphin (*Tursiops truncatus*) off the Atlantic coast of North America: morphologic and ecologic considerations. *International Biological Research Institute Reports* 5:31-43.
- Mullin, K. D., and G. L. Fulling. (in review)2003. Abundance and distribution of cetaceans in the southern U.S. North Atlantic Ocean during summer 1998. *Fish. Bull., U.S.*
- ~~Northridge, S. 1996. Estimation of cetacean mortality in the U.S. Atlantic swordfish and tuna drift gillnet and pair trawl fisheries. Final report to the Northeast Fisheries Science Center, Contract No. 40ENNF500045. 101:603-613.~~
- Palka, D. 1995. Abundance estimate of the Gulf of Maine harbor porpoise. *Rep. int Whal. Commn.* Special Issue 16:27-50
- Palka, D. and Hammond, P.S. 2001. Accounting for responsive movement in line transect estimates of abundance. Can. J. Fish. Aquat. Sci. 58: 777-787
- Palka, D., G. Waring and D. Potter. (inIn review). Abundances of cetaceans and sea turtles in the northwest Atlantic during summer 1995 and 1998. *Fish. Bull., U.S.*
- ~~Scott, G.P. and C.A. Brown. 1997. Estimates of marine mammal and marine turtle catch by the U.S. Atlantic pelagic longline fleet in 1994-1995. Miami Laboratory Contribution MIA-96/97-28~~Thomas, L., J. L. Laake, J. F. Derry, S. T. Buckland, D. L. Borchers, D. R. Anderson, K. P. Burnham, S. Strindberg, S. L. Hedley, F. F. C. Marques, J. H. Pollard and R. M. Fewster. 1998. Distance 3.5. Research Unit for Wildlife Population Assessment, University of St. Andrews, St. Andrews, UK.
- Torres, L. G., P. E. Rosel, C. D'Agrosa and A. J. Read. 2003. Improving managment of overlapping bottlenose dolphin ecotypes through spatial analysis and genetics. Mar. Mamm. Sci. 19:502-514.
- Wade, P. R. and R. P. Angliss. 1997. Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-12, 93 pp.
- ~~Waring, G. T., P. Gerrior, P. M. Payne, B. L. Parry and J. R. Nicolas. 1990. Incidental take of marine mammals in foreign fishery activities off the northeast United States, 1977-1988. Fish. Bull., U.S. 88(2): 347-360.~~
- Wells, R. S., H. L. Rhinehart, P. Cunningham, J. Whaley, M. Baran, C. Koberna and D. P. Costa. 1999. Long distance offshore movements of bottlenose dolphins. *Mar. Mammal Sci.* 15(4):1098-1114.
- ~~Yeung, C. 1999a. Estimates of marine mammal and marine turtle bycatch by the U.S. Atlantic pelagic longline fleet in 1998. NOAA Tech. Memo. NMFS-SEFSC-430, 26 pp. NMFS, Southeast Fisheries Science Center, Miami, FL.~~
- ~~Yeung, C. 1999b. Revised Mortality Estimates of Marine Mammal Bycatch in 1992-1997 based on Serious Injury Guidelines. NOAA Tech. Memo. NMFS-SEFSC-429, 23 pp. NMFS, Southeast Fisheries Science Center, Miami, FL.~~
- ~~Yeung, C. Yeung, C. 2001. Estimates of marine mammal and marine turtle bycatch by the U.S. Atlantic pelagic longline fleet in 1999-2000. NOAA Tech. Memo. NMFS-SEFSC-467, 43 pp. NMFS, Southeast Fisheries Science Center, Miami, FL.~~

ATLANTIC SPOTTED DOLPHIN (*Stenella frontalis*): Western North Atlantic Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

There are two species of spotted dolphin in the Western Atlantic Ocean, the Atlantic spotted dolphin, *Stenella frontalis*, formerly *S. plagiodon* (Perrin *et al.* 1987), and the pantropical spotted dolphin, *S. attenuata*. These species are (Perrin *et al.* 1987). The Atlantic spotted dolphin occurs in two forms which may be distinct sub-species (Perrin *et al.* 1987, 1994; Rice 1998): the large, heavily spotted form which inhabits the continental shelf and is usually found inside or near the 200m isobath; and the smaller, less spotted island and offshore form which occurs in the Atlantic Ocean but is not known to occur in the Gulf of Mexico (Fulling *et al.* 2003; Mullin and Fulling 2003; Mullin and Fulling 2004). Where they co-occur, the offshore form of the Atlantic spotted dolphin and the pantropical spotted dolphin can be difficult to differentiate at sea.

Atlantic spotted dolphins are distributed in tropical and warm temperate waters of the western North Atlantic (Leatherwood *et al.* 1976). Their distribution is from southern New England, south through the Gulf of Mexico and the Caribbean to Venezuela (Leatherwood *et al.* 1976; Perrin *et al.* 1994). The large, heavily spotted form of the Atlantic spotted dolphin along the southeastern and Gulf coasts of the United States, which may warrant designation as a distinct sub-species (Rice 1998), (inhabits the continental shelf, usually being found inside or near the 200 m isobath (within 250-350 km 250-350km of the coast) but sometimes coming into very shallow water adjacent to the beach (Figure 1). Off the northeast USA coast, spotted dolphins are widely distributed on the continental shelf, along the continental shelf edge, and offshore over the deep ocean south of 40° N (CETAP 1982). Atlantic spotted dolphins regularly occur in the inshore waters south of Chesapeake Bay and near the continental shelf edge and continental slope waters north of this region (Payne *et al.* 1984; Mullin in review and Fulling 2003). Sightings have also been made along the north wall of the Gulf Stream and warm-core ring features (Waring *et al.* 1992). Stock structure in the western North Atlantic is unknown.

POPULATION SIZE

Total numbers of Atlantic spotted dolphins off the USA coast or Canadian Atlantic coast are unknown, although three estimates from selected regions of the habitat do exist for select time periods. Because *S. frontalis* and *S. attenuata* are difficult to differentiate at sea, the reported abundance estimates, prior to 1998, are for both species of spotted dolphins combined. Sightings were almost exclusively in the continental shelf edge and continental slope areas west of Georges Bank (Figure 1). concentrated in the slope waters north of Cape Hatteras, but in the shelf waters south of Cape Hatteras, with sightings extending into the deeper slope and offshore waters of the mid-Atlantic (Fig. 1).

An abundance of 6,107 undifferentiated spotted dolphins (CV=0.27) was estimated from an aerial survey program conducted from 1978 to 1982 on the continental, shelf and shelf edge waters between Cape Hatteras, North Carolina and Nova Scotia (CETAP 1982). As recommended in the GAMS Workshop Report (Wade and Angliss 1997), estimates older than eight years are deemed unreliable, therefore should not be used for PBR determinations. Further, due to changes in survey methodology these data should not be used to make comparisons to more current estimates.

An abundance of 4,772 (CV=1.27) undifferentiated spotted dolphins was estimated from a July to September 1995 sighting survey conducted by two ships and an airplane that covered waters from Virginia to the mouth of the Gulf of St. Lawrence (Table 1; Palka *et al.* in review Unpubl. Ms.). Total track line length was 32,600 km 600km. The ships covered waters between the 50 and 1000 fathom depth contour lines, the northern edge of




Figure 1. Distribution of Atlantic spotted dolphin sightings from NEFSC and SEFSC shipboard and aerial surveys during the summer in 1998 and 2004. Isobaths are at 100 m and 1,000 m.

the Gulf Stream, and the northern Gulf of Maine/Bay of Fundy region. The airplane covered waters in the mid-Atlantic from the coastline to the 50 fathom depth contour line, the southern Gulf of Maine, and shelf waters off Nova Scotia from the coastline to the 1000 fathom depth contour line. Data collection and analysis methods used were described in Palka (1996).

An abundance of 32,043 (CV=1.39) for offshore Atlantic spotted dolphins was estimated from a line transect sighting survey conducted during July 6 to September 6, 1998 by a ship and plane that surveyed 15,900 km of track line in waters north of Maryland (38° N) (Figure 1; Palka *et al.* in review Unpubl. Ms.). Shipboard data were analyzed using the modified direct duplicate method (Palka 1995) that accounts for school size bias and $g(0)$, the probability of detecting a group on the track line. Aerial data were not corrected for $g(0)$.

An abundance of 14,396 (CV=0.6263) for offshore, and 15,840 (CV=0.60) for coastal Atlantic spotted dolphins was estimated from a shipboard line transect sighting survey conducted between 8 July and 17 August 1998 that surveyed 5,457 km of track line in waters south of Maryland (38°N) (Figure 1; Mullin in review and Fulling 2003). Abundance estimates were made using the program DISTANCE (Buckland *et al.* 1993; Laake *et al.* 1993, 2001) where school size bias and ship attraction were accounted for.

The best available abundance estimate of 3,040 (CV=0.46) for the Atlantic spotted dolphins is the sum of the estimates from the two 1998 USA Atlantic surveys, 52,279 (CV=0.87), where the estimate from the northern USA Atlantic is 32,043 (CV=1.39) and estimates from the southern USA Atlantic are 4,396 (CV=0.62) and 15,840 (CV=0.60). was estimated from a line transect sighting survey conducted during June 12 to August 4, 2004 by a ship and plane that surveyed 10,761 km of track line in waters north of Maryland (about 38° N) to the Bay of Fundy (about 45° N) (Figure 1; Palka unpubl.). Shipboard data were collected using the two independent team line transect method and analyzed using the modified direct duplicate method (Palka 1995) accounting for biases due to school size and other potential covariates, reactive movements (Palka and Hammond 2001), and $g(0)$, the probability of detecting a group on the track line. Aerial data were collected using the Hiby circle-back line transect method (Hiby 1999) and analyzed accounting for $g(0)$ and biases due to school size and other potential covariates (Figure 1; Palka unpubl.).

A survey of the U.S. Atlantic outer continental shelf and continental slope (water depths >50m) between 27.5 – 38 °N latitude was conducted during June-August, 2004. The survey employed two independent visual teams searching with 50x bigeye binoculars. Survey effort was stratified to include increased effort along the continental shelf break and Gulf Stream front in the mid-Atlantic. The survey included 5,659 km of trackline, and there were a total of 473 cetacean sightings. Sightings were most frequent in waters North of Cape Hatteras, North Carolina along the shelf break. Data were analyzed to correct for visibility bias ($g(0)$) and group-size bias employing line transect distance analysis and the direct duplicate estimator (Palka 1995; Buckland *et al.*, 2001). The resulting abundance estimate for Atlantic spotted dolphins between Florida and Maryland was 47,400 (CV=0.45).

At their November 1999 meeting, the Atlantic SRG recommended that, without a genetic determination of stock structure, the abundance estimates for the coastal and offshore forms should be combined. There remains debate over how distinguishable both species are at sea, though in the waters south of Cape Hatteras identification to species is made with very high certainty. This does not, however, account for the potential for a mixed species herd, as has been recorded for several dolphin assemblages. Pending further genetic studies for clarification of this problem, a single species abundance estimate will be used as the best estimate of abundance, combining species specific data from the northern as well as southern portions of the species' ranges. This joint estimate is considered best because together these two surveys have the most complete coverage of the species' habitat. The best 2004 abundance estimate for Atlantic spotted dolphins is the sum of the estimates from the two 2004 western U.S. Atlantic surveys, 50,440 (CV=0.43), where the estimate from the northern U.S. Atlantic is 3,040 (CV=0.46), and from the southern U.S. Atlantic is 47,400 (CV=0.45).

Table 1. Summary of abundance estimates for both undifferentiated spotted dolphins (1995), and differentiated Atlantic spotted dolphins (1998 and 2004). Month, year, and area covered during each abundance survey, and resulting abundance estimate (N_{best}) and coefficient of variation (CV).

Month/Year	Area	N_{best}	CV
Jul-Sep 1998	Virginia to Gulf of St. Lawrence	32,043 ¹	1.39
Jul-Aug 1998	Florida to Maryland	14,396 ²	0.6263

Jul-Sep 1998	Florida to Gulf of St. Lawrence to Florida (COMBINED)	36,439¹ 1,224² 1	98
Jul-Sep 1998 <u>Jun-Aug 2004</u>	<u>Maryland to Bay of Fundy</u>	<u>3,040</u>	<u>0.46</u>
<u>Jun-Aug 2004</u>	Florida to Maryland	15,840 6,400	0. 60 <u>45</u>
<u>Jun-Aug 2004</u>	<u>Florida to Bay of Fundy (COMBINED)</u>	<u>50,440²</u>	<u>0.45</u>

¹ This represents the first estimate for the offshore Atlantic spotted dolphin.

² This is the combined estimate for the two survey regions

³ This estimate is a recalculation of the same data reported in previous SARs. For more details see Mullin and Fulling 2003.

⁴ This represents the first estimate for the coastal Atlantic spotted dolphin

Minimum Population Estimate

The minimum population estimate is the lower limit of the two-tailed 60% confidence interval of the log-normally distributed best abundance estimate. This is equivalent to the 20th percentile of the log-normal distribution as specified by Wade and Angliss (1997). ~~Until more definitive stock identification (i.e., genetic analysis) work is completed, the Atlantic Scientific Review Group recommends that the best estimate of abundance for Atlantic spotted dolphins is the combined estimates for the offshore 15,840 (CV=0.60) and coastal 36,439 (CV=1.22) forms of Atlantic spotted dolphins. This estimate is 52,279 (CV=0.87). The best abundance estimate is 50,440 (CV=0.43).~~ The minimum population estimates based on the combined offshore and coastal abundance estimates is ~~27,357 (CV=0.87)~~ 745.

Current Population Trend

There are insufficient data to determine the population trends for this species, given that surveys prior to 1998 did not differentiate between species of spotted dolphins.

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Current and maximum net productivity rates are unknown for this stock. For purposes of this assessment, the maximum net productivity rate was assumed to be 0.04. This value is based on theoretical modeling showing that cetacean populations may not grow at rates much greater than 4% given the constraints of their reproductive life history (Barlow *et al.*, 1995).

POTENTIAL BIOLOGICAL REMOVAL

Potential Biological Removal (PBR) is the product of minimum population size, one-half the maximum productivity rate, and a "recovery" factor (MMPA Sec. 3. 16 U.S.C. 1362; Wade and Angliss 1997). The minimum population size for the ~~combined offshore and coastal "forms" of~~ Atlantic spotted dolphins is ~~52,279 (CV=0.87)~~ 745. The maximum productivity rate is 0.04, the default value for cetaceans. The "recovery" factor, which accounts for endangered, depleted, threatened stocks, or stocks of unknown status relative to optimum sustainable population (OSP) is set to 0.5 because this stock is of unknown status. PBR for the combined offshore and coastal forms of Atlantic spotted dolphins is ~~278 (CV=0.87)~~ 357.

ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

Fishery Information

Detailed fishery information is reported in Appendix III. Total annual estimated average fishery-related mortality or serious injury to this stock during ~~1994-1998~~ 1999-2003 was ~~7.8 undifferentiated zero Atlantic~~ spotted dolphins (*Stenella* spp.) ~~CV=0.01 (Yeung 2001; Table 2).~~

Fishery Information

Garrison 2003; Garrison and Richards, 2004).

Earlier Interactions

No spotted dolphin mortalities were observed in 1977-1991 foreign fishing activities. ~~Data on current incidental takes in USA fisheries are available from several sources. In 1986, NMFS established a mandatory self-reported fisheries information system for large pelagic fisheries. Data files are maintained at the Southeast Fisheries Science Center (SEFSC). The Northeast Fisheries Science Center (NEFSC) Fisheries Observer Program was initiated in 1989 and since that year several fisheries have been covered by the program. In late 1992 and in 1993, the~~

SEFSC provided observer coverage of pelagic longline vessels fishing off the Grand Banks (Tail of the Banks) and provides observer coverage of vessels fishing south of Cape Hatteras. Total fishery-related mortality and serious injury cannot be estimated separately for the two species of spotted dolphins in the USA Atlantic Exclusive Economic Zone (EEZ) because of the uncertainty in species identification by fishery observers. The Atlantic Scientific Review Group advised adopting the risk-averse strategy of assuming that either species might have been subject to the observed fishery-related mortality and serious injury.

Bycatch ~~has had~~ been observed by NMFS Sea Samplers in the pelagic drift gillnet and pelagic longline fisheries, but no mortalities or serious injuries have been documented in the pelagic pair trawl, Northeast multispecies sink gillnet, mid-Atlantic coastal gillnet, and North Atlantic bottom trawl fisheries; and no takes have been documented in a review of Canadian gillnet and trap fisheries (Read 1994).

Pelagic Drift Gillnet

The estimated total number of hauls in the pelagic drift gillnet fishery increased from 714 in 1989 to 1,144 in 1990; thereafter, with the introduction of quotas, effort was severely reduced. The estimated number of hauls in 1991, 1992, 1993, 1994, 1995, 1996, and 1998 were 233, 243, 232, 197, 164, 149, and 113 respectively. In 1996 and 1997, NMFS issued management regulations which prohibited the operation of this fishery in 1997. Further, in January 1999 NMFS issued a Final Rule to prohibit the use of driftnets (*i.e.*, permanent closure) in the North Atlantic swordfish fishery (50 CFR Part 630). Fifty-nine different vessels participated in this fishery at one time or another between 1989 and 1993. Since 1994, between 10- and 13 vessels have participated in the fishery (Table 2). Observer coverage, expressed as percent of sets observed, was 8% in 1989, 6% in 1990, 20% in 1991, 40% in 1992, 42% in 1993, 87% in 1994, 99% in 1995, 64% in 1996, and 99% in 1998. Effort was concentrated along the southern edge of Georges Bank and off Cape Hatteras. Examination of the species composition of the catch and locations of the fishery throughout the year, suggested that the pelagic drift gillnet fishery be stratified into two strata, a southern or winter stratum, and a northern or summer stratum. Estimates of the total bycatch, from 1989 to 1993, were obtained using the aggregated (pooled 1989-1993) catch rates, by strata (Northridge 1996). Estimates of total annual bycatch for 1994 and 1995 were estimated from the sum of the observed caught and the product of the average bycatch per haul and the number of unobserved hauls as recorded in self-reported fisheries information. Variances were estimated using bootstrap re-sampling techniques.

Forty-nine undifferentiated spotted dolphins mortalities were observed in the drift gillnet fishery between 1989 and 1998 and occurred northeast of Cape Hatteras within the ~~183 m~~ **183m** isobath in February-April; and near Lydonia Canyon in October. Six whole animal carcasses that were sent to the Smithsonian were identified as Pantropical spotted dolphins (*S. attenuata*). The remaining animals were not identified to species. Estimated annual mortality and serious injury attributable to this fishery (CV in parentheses) was 25 in 1989 (.65), 51 in 1990 (.49), 11 in 1991 (.41), 20 in 1992 (0.18), 8.4 in 1993 (0.40), 29 in 1994 (0.01), 0 in 1995, 2 in 1996 (0.06), ~~NA in 1997, and 0 in 1998;~~ average annual mortality and serious injury during 1994-1998 was 7.8 (0.01) (Table 2).

Pelagic Longline fishery in 1997 and 0 in 1998.

The pelagic longline fishery operates in the ~~USA~~ **U.S.** Atlantic (including Caribbean) and Gulf of Mexico EEZ (~~SEFSC unpublished data~~). Interactions between the pelagic longline fishery and spotted dolphins have been reported; however, a vessel may fish in more than one statistical reporting area and it is not possible to separate estimates of fishing effort other than to subtract Gulf of Mexico effort from Atlantic fishing effort, which includes the Caribbean Sea. ~~This fishery has been monitored with about 5% observer coverage, in terms of trips observed, since 1992. Total effort from 1999-2003, excluding the Gulf of Mexico, for the pelagic longline fishery, based on mandatory self-reported fisheries information, was 11,279 sets in 1991, 9,869 sets in 1992, 9,862 sets in 1993, 9,481 sets in 1994, 10,129 sets in 1995, 9,885 sets in 1996, 8,023 sets in 1997, and 6,675 sets in 1998 (Cramer 1994; Scott and Brown 1997; Johnson *et al.* 1999; Yeung 1999b). Since 1992, this fishery has been monitored with about 5% observer coverage, in terms of trips observed, within every statistical reporting area within the EEZ and beyond. Off the USA Atlantic coast, the fishery has been observed from January to March off Cape Hatteras, in May and June in the entire mid-Atlantic, and in July through December in the mid-Atlantic Bight and off Nova Scotia. The 1994-1998, estimated take was based on a revised analysis of the observed incidental take and self-reported incidental take and effort data, and replace previous estimates for the 1992-1993 and 1994-1995 periods (Cramer 1994; Scott and Brown 1997; Johnson *et al.* 1999; Yeung 1999a). Further, Yeung (1999b), revised the 1992-1997 fishery mortality estimates in Johnson *et al.* (1999) to include seriously injured animals. The 1998 bycatch estimates were from Yeung (1999a). Most of the estimated marine mammal bycatch was from EEZ waters between South Carolina and Cape Cod (Johnson *et al.* 1999). Excluding the Gulf of Mexico where one animal was hooked and released alive (Appendix 1), no Atlantic spotted dolphin bycatches were observed for 1992-1998.~~

Table 2. Summary of the incidental mortality of undifferentiated spotted dolphins (*Stenella* sp.) by commercial fishery including the years sampled (Years), the number of vessels active within the fishery (Vessels), the type of data used (Data Type), the annual observer coverage (Observer Coverage), the mortalities recorded by on-board observers (Observed Mortality), the estimated annual mortality (Estimated Mortality), the estimated CV of the annual mortality (Estimated CVs) and the mean annual mortality (CV in parentheses).

Fishery Years Vessels

Data Type¹

Observed Serious Injury Observer Coverage² Observed Mortality Estimated Mortality⁵

Estimated CVs

Mean Annual Mortality Pelagic Drift Gillnet⁶ 94-98 1994=11³

1995=12 1996=10

1998=13 Obs. Data Logbook 0, 0, 0, 0, 0 .87, .99, .64, NA, .99 29, 0, 2, NA, 029, 0, 2⁴, NA, 0 .01, 0, 0, NA, 0 7.75
(0.01) TOTAL 7.8

(0.01)¹ Observer data (Obs. Data) are used to measure bycatch rates, and the data are collected within the Northeast Fisheries Science Center (NEFSC) Fisheries Observer Program. Mandatory logbook (Logbook) data are used to measure total effort for the pelagic drift gillnet fishery, and these data are collected at the Southeast Fisheries Science Center (SEFSC).

² The observer coverage for the pelagic drift gillnet and pair trawl fishery is measured in terms of sets, and the longline fishery is in trips.

³ 1994, 1995, 1996, and 1998 shown, other years not available on an annual basis.

⁴ Estimates were based on two seasons. The two observed takes were during the winter season when observer coverage was 100%.

⁵ Annual mortality estimates include animals seriously injured and released alive.

⁶ The fishery did not operate in 1997, the average annual mortality is based on the number of years (4; 1994-1998) that the fishery operated recorded.

Other Mortality

From ~~1995-1998~~ 1999-2003, ~~thirteen~~ 16 Atlantic spotted dolphins were stranded between ~~North Carolina~~ Massachusetts and Florida (NMFS unpublished data).

One animal stranded in North Carolina in 1999, 3 animals stranded in North Carolina and 1 stranded in Georgia in 2000, 2 animals stranded in North Carolina and 3 in Florida in 2001, 2 animals stranded in North Carolina and 2 in Florida in 2002, and 1 animal stranded in Massachusetts, 1 in North Carolina and 1 in Florida in 2003. None of these strandings had documented signs of human interactions.

Table 2. Atlantic spotted dolphin (*Stenella frontalis*) strandings along the U.S. Atlantic coast, 1999-2003

<u>STATE</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>TOTALS</u>
<u>Massachusetts</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>1</u>	<u>1</u>
<u>North Carolina</u>	<u>0</u>	<u>3</u>	<u>2</u>	<u>2</u>	<u>1</u>	<u>8</u>
<u>South Carolina</u>	<u>1</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>1</u>
<u>Georgia</u>	<u>0</u>	<u>1</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>1</u>
<u>Florida</u>	<u>0</u>	<u>0</u>	<u>3</u>	<u>2</u>	<u>1</u>	<u>6</u>
<u>TOTALS</u>	<u>1</u>	<u>4</u>	<u>5</u>	<u>4</u>	<u>3</u>	<u>17</u>

STATUS OF STOCK

The status of Atlantic spotted dolphins, relative to OSP in the ~~USA~~ U.S. Atlantic EEZ is unknown. The species is not listed as threatened or endangered under the Endangered Species Act. There are insufficient data to determine the population trends for this species. Total fishery-related mortality and serious injury for this stock is less than 10% of the calculated PBR and, therefore, can be considered to be insignificant and approaching zero mortality and serious injury rate. Average annual fishery-related mortality and serious injury does not exceed the PBR; therefore, this is not a strategic stock.

REFERENCES

- Barlow, J., S. L. Swartz, T. C. Eagle and P. R. Wade. 1995. U.S. Marine ~~M~~mammal ~~S~~stock ~~A~~assessments: Guidelines for ~~P~~reparation, ~~B~~ackground, and a ~~S~~ummary of the 1995 ~~A~~assessments. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-6, ~~73 pp~~ 73pp. Available from NMFS, Southwest Fisheries Science Center, 8604 La Jolla Shores Drive, La Jolla, CA, 92037-1508.
- Buckland, S. T., D. R. Anderson, K. P. Burnham and J. L. Laake. ~~1993~~ 2001. ~~D~~Borchers and L. Thomas. 2001. Introduction to distance ~~S~~ampling: Estimating abundance of biological populations. ~~Chapman and Hall~~ Oxford University Press, New York, ~~446~~ 32 pp.-
- CETAP. 1982. A characterization of marine mammals and turtles in the mid- and north Atlantic areas of the U.S. outer continental shelf. Cetacean and Turtle Assessment Program, University of Rhode Island. Final Report, Contract AA51-C78-48, Bureau of Land Management, Washington, DC, 538 pp.
- Cramer, J. 1994. Large pelagic logbook newsletter - 1993. NOAA Tech. Mem. NMFS-SEFSC-352, 19 pp.
- Garrison, L.P. 2003. Estimated bycatch of marine mammals and marine turtle catch bys in the U.S. Atlantic pelagic longline fleet in 1992-1997. SEFSC PRD-98/99-03. 68 pp.
- ~~Laake, J. L., S. T. Buckland, D. R. Anderson and K. P. Burnham. 1993. DISTANCE user's guide, V2.0. Colorado~~

- Cooperative Fish & Wildlife Research Unit, Colorado State University, Ft. Collins, Colorado, 72 during 2001-2002. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-SEFSC-515, 52 pp. Available from NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Road, Miami, FL 33149.
- Garrison, L.P. and P.M. Richards. 2004. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2003. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-SEFSC-527, 57 pp. Available from NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Road, Miami, FL 33149.
- Hiby, L. 1999. The objective identification of duplicate sightings in aerial survey for porpoise. pp. 179-189. In: G.W. Garner, S. C. Amstrup, J. L. Laake, B. F. J. Manly, L. L. McDonald, and D. G. Robertson (eds.) *Marine mammal survey and assessment methods*. Balkema, Rotterdam. 287 pp.
- Leatherwood, S., D. K. Caldwell and H. E. Winn. 1976. Whales, dolphins, and porpoises of the western North Atlantic. A guide to their identification. NOAA Tech. Rep. NMFS Circ. 396, U.S. Dept. Commer. Washington, DC 176 pp.
- Mullin, K. D. ~~(in review)~~ and G. L. Fulling. 2003. Abundance and ~~distribution of~~ cetaceans in the southern U.S. Atlantic Ocean during summer 1998. *Fish. Bull.*, U.S.
- ~~Northridge, S. 1996. Estimation of cetacean mortality in the U.S. Atlantic swordfish and tuna drift gillnet and pair trawl fisheries. Final report to the Northeast Fisheries Science Center, Contract No. 40ENNF500160.~~
- Palka, D. 1995. 101:603-613.
- Palka, D. 1995. Abundance estimate of the Gulf of Maine harbor porpoise. pp. 27-50 In: A. Bjørge and G.P. Donovan (eds.) *Biology of the Phocoenids*. Rep. Int. Whal. Commn. (Special Issue 16:27-50) I-x + 552 pp.
- Palka, D. 1996. Update on abundance of Gulf of Maine/Bay of Fundy harbor porpoises. NOAA/NMFS/NEFSC. Ref. Doc. 96-04; 37 pp. Available from: National Marine Fisheries Service, 166 Water Street, Woods Hole, MA 02543-1026.
- Palka, D., G. Waring, and D. Potter. (in review). Abundances of cetaceans and sea turtles in the northwest Atlantic during summer 1995 and 1998. *Fish. Bull., U.S. and P.S. Hammond. 2001. Accounting for responsive movement in line transect estimates of abundance. Can. J. Fish. Aquat. Sci. 58: 777-787.*
- Payne, P. M., L. A. Selzer and A. R. Knowlton. 1984. Distribution and density of cetaceans, marine turtles, and seabirds in the shelf waters of the northeastern United States, June 1980-December 1983, based on shipboard observations. NOAA/NMFS Contract No. NA-81-FA-C-00023. Available from: National Marine Fisheries Service, 166 Water Street, Woods Hole, MA 02543-1026.
- Perrin, W. F., E. D. Mitchell, J. G. Mead, D. K. Caldwell, M. C. Caldwell, P. J. H. van Bree, and W. H. Dawbin. 1987. Revision of the spotted dolphins, *Stenella* sp. *Mar. Mammal Sci.* 3(2): 99-170.
- Perrin, W. F., D. K. Caldwell, and M. C. Caldwell. 1994. Atlantic spotted dolphin. Pages pp. 173-190. In: S. H. Ridgway and R. Harrison (eds.). *Handbook of marine mammals, Volume 5: The first book of dolphins*. Academic Press, San Diego, 418 pp.
- Read, A. J. 1994. Interactions between cetaceans and gillnet and trap fisheries in the northwest Atlantic. pp. 133-147. In: W.F. Perrin, G.P. Donovan and J. Barlow (eds.) *Gillnets and Cetaceans*. Rep. Int. Whal. Commn. (Special Issue 15). I-ix + 629 pp.
- Rice, D. W. 1998. Marine mammals of the world, systematics and distribution. *Spec. Publ. No 4. Society for The Society for Marine Mammalogy*, Lawrence, KS. 231 pp.
- Scott, G. P. and C. A. Brown. 1997. Estimates of marine mammal and marine turtle catch by the U.S. Atlantic pelagic longline fleet in 1994-1995. Miami Laboratory Contribution MIA-96/97-28
- Wade, P.-R., and R.-P. Angliss. 1997. Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. U.S. Dep. Commer., NOAA Tech. Memo. -NMFS-OPR-12, 93 pp. Available from: NOAA, National Marine Fisheries Service, 166 Water St., Woods Hole, MA 02543.
- Waring, G. T., C. P. Fairfield, C. M. Ruhsam and M. Sano. 1992. Cetaceans associated with Gulf Stream features off the northeastern USA shelf. ICES Marine Mammals Comm. CM 1992/N:12, 29 pp.
- Yeung, C. ~~1999a~~ 2001. Estimates of marine mammal and marine turtle bycatch by the U.S. Atlantic pelagic longline fleet in 1998. NOAA Technical Memorandum NMFS-SEFSC-430, 26 pp.
- Yeung, C. 1999b. Revised Mortality Estimates of Marine Mammal Bycatch in 1992-1997 based on Serious Injury Guidelines. NOAA Technical Memorandum NMFS-SEFSC-429, 23 pp. 1999-2000. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-SEFSC-467, National Marine Fisheries Service, Miami, FL, 43 pp. Available from NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Road, Miami, FL 33149.

CLYMENE DOLPHIN (*Stenella clymene*): Western North Atlantic Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

The Clymene dolphin is endemic to tropical and sub-tropical waters of the Atlantic (Jefferson and Curry 2003). Clymene dolphins have been commonly sighted in the Gulf of Mexico since 1990 (Mullin *et al.* 1994; Fertl *et al.* 2003), and a Gulf of Mexico stock has been designated since 1995. Four Clymene dolphin groups were sighted during summer 1998 in the western North Atlantic (Mullin and Fulling 2003), and two groups were sighted in the same general area during a 1999 bottlenose dolphin survey (NMFS unpublished). These sightings and stranding records (Fertl *et al.* 2003) indicate that this species routinely occurs in the western North Atlantic. The western North Atlantic population is provisionally being considered a separate stock for management purposes, although there is currently no information to differentiate this stock from the northern Gulf of Mexico stock(s). Additional morphological, genetic and/or behavioral data are needed to provide further information on stock delineation.

POPULATION SIZE

The numbers of Clymene dolphins off the U.S. or Canadian Atlantic coast are unknown, and seasonal abundance estimates are not available for this species since it was rarely seen in any surveys.

Clymene dolphins were observed during earlier surveys along the U.S. Atlantic coast. Estimates of abundance were derived through the application of distance sampling analysis (Buckland *et al.* 2001) and the computer program DISTANCE (Thomas *et al.* 1998) to sighting data. Data were collected using standard line-transect techniques conducted from NOAA Ship *Relentless* during July and August 1998 between Maryland (38.00°N) and central Florida (28.00°N) from the 10 m isobath to the seaward boundary of the U.S. EEZ. Transect lines were placed perpendicular to bathymetry in a double saw-tooth pattern. Sightings of Clymene dolphins were primarily on the continental slope east of Cape Hatteras, North Carolina (Fig. 1). The best estimate of abundance for the Clymene dolphin was 6,086 (CV=0.93) (Mullin and Fulling 2003) and represents the first and only estimate to date for this species in the U.S. Atlantic EEZ.

Minimum Population Estimate

The minimum population estimate is the lower limit of the two-tailed 60% confidence interval of the log-normally distributed best abundance estimate. This is equivalent to the 20th percentile of the log-normal distributed abundance estimate as specified by Wade and Angliss (1997). The best estimate of abundance for the western North Atlantic Clymene dolphin stock, based on the 1998 surveys, is 6,086 (CV=0.93). The minimum population estimate for the western North Atlantic stock is 3,132 Clymene dolphins.




Figure 1. Distribution of Clymene dolphin sightings from NEFSC and SEFSC vessel and aerial summer surveys during 1998 and 2004. Isobaths are at 100 and 1,000 m.

Current Population Trend

There are insufficient data to determine the population trends for this ~~species~~stock.

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Current and maximum net productivity rates are unknown for this ~~species~~stock. For purposes of this assessment, the maximum net productivity rate was assumed to be 0.04. This value is based on theoretical modeling showing that cetacean populations may not grow at rates much greater than 4% given the constraints of their reproductive history (Barlow *et al.* 1995).

POTENTIAL BIOLOGICAL REMOVAL

Potential ~~B~~Biological ~~R~~Removal (PBR) is the product of minimum population size, one half the maximum net productivity rate, and a recovery factor (MMPA Sec. 3. 16 U.S.C. 1362; Wade and Angliss 1997). The minimum population size is 3,132. The maximum productivity rate is 0.04, the default value for cetaceans. The “recovery” factor, which accounts for endangered, depleted, threatened stocks, or stocks of unknown status relative to optimum sustainable population (OSP) is assumed to be 0.5 because this stock is of unknown status. PBR for the western North Atlantic Clymene dolphin stock is 31.

ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

~~Annual human-caused mortality and serious injury is unknown for this species.~~

Fisheries Information

~~Fishery-related~~Fishery Information

Detailed fishery information is reported in Appendix III. Total annual estimated fishery-related mortality and serious injury to this stock during 1999-2003 was zero Clymene dolphins, as there were no reports of mortalities or serious injury ~~is unknown for this species.~~

Other Mortality

~~to Clymene dolphins (Yeung 2001; Garrison 2003; Garrison and Richards, 2004).~~

Other Mortality

There have been 2 reported strandings of Clymene dolphins in the western North Atlantic between 1997-2002~~1999-2003~~. No signs of human interactions were noted in either stranding. There may be some uncertainty in the identification of this species due to similarities with other *Stenella* species.

STATUS OF STOCK

The status of ~~the~~ Clymene ~~dolphin stock in the western North Atlantic EEZ~~dolphins, relative to ~~OSPOPS~~in the EEZ is unknown. The species is not listed as threatened or endangered under the Endangered Species Act. There are insufficient data to determine the population trends for this ~~species~~stock. The total fishery-related mortality and serious injury for this stock is unknown, but assumed to be less than 10% of the calculated PBR and can be considered to be insignificant and approaching zero mortality and serious injury rate. This ~~is~~is not a strategic stock because the average annual fishery-related mortality and serious injury has not exceeded PBR for the last two years.

REFERENCES

- Barlow, J., S. L. Swartz, T. C. Eagle, and P. R. Wade. 1995. U.S. marine mammal stock assessment: guidelines for preparation, background, and a summary of the 1995 assessments. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-6, National Marine Fisheries Service, Seattle, WA, 73 pp. Available from: NOAA National Marine Fisheries Service, 7600 Sand Point Way NE, Seattle, WA 98115-0070.
- Buckland, S. T., D. R. Anderson, K. P. Burnham, J. L. Laake, D. L. Borchers and L. Thomas. 2001. *Introduction to distance sampling: estimating abundance of biological populations*. Oxford University Press, New York, 432 pp.
- Fertl, D., T. A. Jefferson, I. B. Moreno, A. N. Zerbini and K. D. Mullin. 2003. Distribution of the Clymene dolphin *Stenella clymene*. *Mammal Review* 33(3):253-271.

- Garrison, L.P. 2003. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2001-2002. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-SEFSC-515, 52 pp. Available from NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Road, Miami, FL 33149.
- Garrison, L.P. and P.M. Richards. 2004. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2003. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-SEFSC-527, 57 pp. Available from NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Road, Miami, FL 33149.
- Jefferson, T. A. and B. E. Curry. 2003. *Stenella clymene*. *Mammalian Species* 726:1-5.
- Mullin, K. D. and G. L. Fulling. 2003. Abundance of cetaceans in the southern U.S. North Atlantic Ocean during summer 1998. *Fish. Bull.* 101:603-613.
- Mullin, K. D., T. A. Jefferson, and L. J. Hansen. 1994. Sightings of the clymene dolphin (*Stenella clymene*) in the Gulf of Mexico. *Mar. Mamm. Sci.* 10:464-470.
- Palka, D. 1995. Abundance estimate of the Gulf of Maine harbor porpoise. pp. 27-50 In: A. Bjørge and G.P. Donovan (eds.) *Biology of the Phocoenids*, Rep. Int. Whal. Commn (Special Issue 16) i-x + 552 pp.
- Palka, D. and P.S. Hammond. 2001. Accounting for responsive movement in line transect estimates of abundance. *Can. J. Fish. Aquat. Sci.* 58: 777-787.
- Thomas, L., J. L. Laake, J. F. Derry, S. T. Buckland, D. L. Borchers, D. R. Anderson, K. P. Burnham, S. Strindberg, S. L. Hedley, F. F. C. Marques, J. H. Pollard and R. M. Fewster. 1998. Distance 3.5. Research Unit for Wildlife Population Assessment, University of St. Andrews, St. Andrews, UK.
- Wade, P. R. and R. P. Angliss. 1997. Guidelines for assessing marine mammal stocks: Report of the GAMMS workshop April 3-5, 1996, Seattle, Washington. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-12, National Marine Fisheries Service, Seattle, WA, 93pp. Available from: NOAA National Marine Fisheries Service, 7600 Sand Point Way NE, Seattle, WA 98115-0070.
- Yeung, C. 2001. Estimates of marine mammal and marine turtle bycatch by the U.S. Atlantic pelagic longline fleet in 1999-2000. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-SEFSC-467, National Marine Fisheries Service, Miami, FL, 43 pp. Available from NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Road, Miami, FL 33149.

DWARF SPERM WHALE (*Kogia sima*): Western North Atlantic Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

The dwarf sperm whale (*Kogia sima*) appears to be distributed worldwide in temperate to tropical waters (Caldwell and Caldwell 1989). There are no stranding records for the east Canadian coast (Willis and Baird 1998). Sightings of these animals in the western North Atlantic occur in oceanic waters (Mullin and Fulling 2003; NMFS unpublished data). Dwarf sperm whales and pygmy sperm whales (*K. breviceps*) are difficult to distinguish differentiate at sea (Caldwell and Caldwell 1989, Wursig *et al.* 2000), and sightings of either species are often categorized as *Kogia* spp. There is no information on stock differentiation for the Atlantic population. In a recent study using Duffield *et al.* (2003) propose using the molecular weights of myoglobin and hemoglobin, as determined by blood or muscle tissues of stranded animals, as a quick and robust way to provide species confirmation. Using hematological and as well as stable-isotope data, Barros *et al.* (1998) speculated that dwarf sperm whales may have a more pelagic distribution than pygmy sperm whales, and/or dive deeper during feeding bouts.

Diagnostic morphological characters have also been useful in distinguishing the two *Kogia* species (Barros and Duffield 2003), thus enabling researchers to use stranding data in distributional and ecological studies.. Specifically, the distance from the snout to the center of the blowhole in proportion to the animal's total length, as well as the height of the dorsal fin, in proportion to the animal's total length, can be used to differentiate between the two *Kogia* species when such measurements are obtainable (Barros and Duffield 2003).

POPULATION SIZE

Total numbers of dwarf sperm whales off the U.S. or Canadian Atlantic coast are unknown, although estimates from selected regions of the habitat do exist for select time periods. Because *Kogia sima* and *Kogia breviceps* are difficult to differentiate at sea, the reported abundance estimates are for both species of *Kogia*.

An abundance of 115 (CV=0.61) for *Kogia* spp. was estimated from a line-transect survey conducted from July 6 to September 6, 1998, by a ship and plane that surveyed 15,900 km of track line in waters north of Maryland (38° N) (Fig. 1; Palka *et al.*, in review Unpubl. Ms.). Shipboard data were analyzed using the modified direct duplicate method (Palka 1995) that accounts for school size bias and $g(0)$, the probability of detecting a group on the track line. Aerial data were not corrected for $g(0)$.

An abundance of 580 (CV=0.57) for *Kogia* spp. was estimated from a shipboard line-transect sighting survey conducted between 8 July and 17 August 1998 that surveyed 4,163 km of track line in waters south of Maryland (38°N) (Fig. 1; Mullin and Fulling 2003). Abundance estimates were made using the program DISTANCE (Buckland *et al.* 2001; Thomas *et al.* 1998).




Figure 1. Distribution of *Kogia* spp. sightings from NEFSC and SEFSC shipboard and aerial surveys during the summer in 1998 and 2004. Isobaths are at 100 m and 1,000 m.

The best available An abundance of 344 (CV= 0.32) for *Kogia* spp. was estimated from a line transect sighting survey conducted during June 12 to August 4, 2004 by a ship and plane that surveyed 10,761 km of track line in waters north of Maryland (about 38° N) to the Bay of Fundy (about 45° N) (Figure 1; Palka unpubl.). Shipboard data were collected using the two independent team line transect method and analyzed using the modified direct duplicate method (Palka 1995) accounting for biases due to school size and other potential covariates, reactive movements (Palka and Hammond 2001), and $g(0)$, the probability of detecting a group on the track line. Aerial data were collected using the Hiby circle-back line transect method (Hiby 1999) and analyzed accounting for $g(0)$ and biases due to school size and other potential covariates (Figure 1; Palka unpubl.).

A survey of the U.S. Atlantic outer continental shelf and continental slope (water depths > 50m) between 27.5 – 38 °N latitude was conducted during June-August, 2004. The survey employed two independent visual teams searching with 50x bigeye binoculars. Survey effort was stratified to include increased effort along the continental shelf break and Gulf Stream front in the mid-Atlantic. The survey included 5,659 km of trackline, and there were a total of 473 cetacean sightings. Sightings were most frequent in waters North of Cape Hatteras, North Carolina along the shelf break. Data were analyzed to correct for visibility bias ($g(0)$) and group-size bias employing line transect distance analysis and the direct duplicate estimator (Palka 1995; Buckland *et al.*, 2001). The resulting abundance estimate for *Kogia* spp. between Florida and Maryland was 37 (CV=0.75).

The best 2004 abundance estimate for *Kogia* spp. is the sum of the estimates from the two 1998-2004 U.S. Atlantic surveys, 695-381 (CV=0.4930), where the estimate from the northern U.S. Atlantic is 115-344 (CV=0.6132), and from the southern U.S. Atlantic is 580-37 (CV=0.5775). This joint estimate is considered the best because together these two surveys have the most complete coverage of the species' habitat.

A separate estimate of dwarf sperm whale abundance cannot be provided due to the uncertainty of species identification at sea.

Table 1. Summary of abundance estimates for the western North Atlantic *Kogia* spp. Month, year, and area covered during each abundance survey, and resulting abundance estimate (N_{best}) and coefficient of variation (CV).

<u>Month/Year</u>	<u>Area</u>	<u>N_{best}</u>	<u>CV</u>
<u>Jul-Sep 1998</u>	<u>Maryland to Gulf of St. Lawrence</u>	<u>115</u>	<u>0.61</u>
<u>Jul-Aug 1998</u>	<u>Florida to Maryland</u>	<u>580</u>	<u>0.57</u>
<u>Jul-Sep 1998</u>	<u>Florida to Gulf of St. Lawrence (COMBINED)</u>	<u>695¹</u>	<u>0.49</u>
<u>Jun-Aug 2004</u>	<u>Maryland to Bay of Fundy</u>	<u>344</u>	<u>0.32</u>
<u>Jun-Aug 2004</u>	<u>Florida to Maryland</u>	<u>37</u>	<u>0.75</u>
<u>Jun-Aug 2004</u>	<u>Bay of Fundy to Florida (COMBINED)</u>	<u>381¹</u>	<u>0.30</u>

¹This is the combined estimate for the two survey areas

Minimum Population Estimate

The minimum population estimate is the lower limit of the two-tailed 60% confidence interval of the log-normally distributed best abundance estimate. This is equivalent to the 20th percentile of the log-normal distribution as specified by Wade and Angliss (1997). The best estimate of abundance for *Kogia* spp. is 695-381 (CV=0.4930). The minimum population estimate for *Kogia* spp. is 470-298.

Current Population Trend

The available information is insufficient to evaluate trends in population size for this species in the western North Atlantic.-

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Current and maximum net productivity rates are unknown for this stock. For purposes of this assessment, the maximum net productivity rate was assumed to be 0.04. This value is based on theoretical modeling showing that cetacean populations may not grow at rates much greater than 4% given the constraints of their reproductive life history (Barlow *et al.*- 1995).

POTENTIAL BIOLOGICAL REMOVAL

Potential ~~b~~Biological ~~r~~Removal (PBR) is the product of minimum population size, one-half the maximum productivity rate, and a “recovery” factor (Wade and Angliss 1997). The minimum population size is ~~470~~298. The maximum productivity rate is 0.04, the default value for cetaceans. The “recovery” factor, which accounts for endangered, depleted, threatened stocks, or stocks of unknown status relative to optimum sustainable population (OSP) is assumed to be 0.5 because this stock is of unknown status. PBR for the western North Atlantic *Kogia* spp. is ~~43.7~~-.

ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

~~The level of past or current, direct, human-caused mortality of dwarf sperm whales in the U.S. Atlantic EEZ is unknown. Available information indicates there is likely little fisheries interaction with dwarf sperm whales in the U.S. Atlantic EEZ~~

Fishery Information

~~Detailed fishery information is reported in Appendix III. There has been no logbook report of fishery-related serious injury recorded off the east coast of Florida in the pelagic longline fishery in 2000 (Table 2) (Yeung 2001; Garrison 2003; Garrison and Richards, 2004). Total annual estimated average fishery-related mortality ~~or~~and serious injury to this stock during ~~1996-2000 was 0~~1999-2003 was zero for dwarf sperm whales, as there were no reports of mortality or serious injury to dwarf sperm whales (Table 1).~~Fishery Information
Yeung 2001; Garrison 2003; Garrison and Richards 2004).

Earlier Interactions

~~No dwarf sperm whale mortalities were observed in 1977-1991 foreign fishing activities.~~ Bycatch has been observed by NMFS Sea Samplers in the pelagic drift gillnet fishery, but no mortalities or serious injuries have been documented in other fisheries.

Pelagic Drift Gillnet

There was one report of mortality or serious injury to a dwarf sperm whale attributable to the pelagic drift gillnet fishery. Estimated annual fishery-related mortality and serious injury (CV in parentheses) was 0 dwarf sperm whales from 1991-1994, 1.0 in 1995 (CV=0), and 0 from ~~1996-1998.~~ Estimated average annual mortality and serious injury related to this fishery during 1994-1998 was 0.251996-2003.

Other Mortality

~~From 1999-2003, 37 dwarf sperm whales (CV=0). Other Mortality were reported stranded between North Carolina and Puerto Rico (Table 2). No dwarf sperm whales were reported to stranded in Nova Scotia from 1990-2004 (T. Wimmer, Nova Scotia Marine Animal Response Society, pers. comm.). The total includes 8 animals stranded in North Carolina and 1 in Georgia in 1999; 4 animals stranded in North Carolina, 1 in South Carolina, and 4 in Florida in 2000; 1 animal stranded in North Carolina, 1 in South Carolina, and 2 in Florida in 2001; 3 animals stranded in Florida and 2 in Puerto Rico in 2002; and 4 animals stranded in North Carolina, 2 in South Carolina, 2 in Georgia, and 2 in Florida in 2003. In addition to the above strandings of *Kogia sima*, there were 8 strandings reported as *Kogia* spp. as follows: 1 *Kogia* spp. stranded in Georgia in 2000, 1 stranded in North Carolina and 2 in Florida in 2002, and 1 stranded in Georgia and 3 in Florida in 2003.~~

Table 2. Dwarf sperm whale (*Kogia sima*) strandings along the Atlantic coast, 1999-2003

<u>STATE</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>TOTALS</u>
--------------	-------------	-------------	-------------	-------------	-------------	---------------

<u>North Carolina</u>	<u>8</u>	<u>4</u>	<u>1¹</u>	<u>0¹</u>	<u>4</u>	<u>17</u>
<u>South Carolina</u>	<u>0</u>	<u>1</u>	<u>1</u>	<u>0</u>	<u>2</u>	<u>4</u>
<u>Georgia</u>	<u>1</u>	<u>0¹</u>	<u>0</u>	<u>0¹</u>	<u>2¹</u>	<u>3</u>
<u>Florida</u>	<u>0</u>	<u>4</u>	<u>2</u>	<u>3²</u>	<u>2³</u>	<u>11</u>
<u>Puerto Rico</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>2</u>	<u>0</u>	<u>2</u>
<u>TOTALS</u>	<u>9</u>	<u>9</u>	<u>4</u>	<u>5</u>	<u>10</u>	<u>37</u>

~~¹1 additional *Kogia* spp. stranded~~

~~²2 additional *Kogia* spp. stranded~~

~~³3 additional *Kogia* spp. stranded~~

There were no documented strandings of dwarf sperm whales along the U.S. Atlantic coast during 1999-2003 which were classified as likely caused by fishery interactions.

Historical stranding records (1883-1988) of dwarf sperm whales in the southeastern U.S. (Credle 1988), and strandings recorded during 1988-1997 (Barros *et al.* 1998) indicate that this species accounts for about 17% of all *Kogia* strandings in this area. During the period 1990-October 1998, 3 dwarf sperm whale strandings occurred in the northeastern U.S. (Maryland, Massachusetts, and Rhode Island), whereas 43 strandings were documented along the U.S. Atlantic coast between North Carolina and the Florida Keys in the same period. A pair of latex examination gloves was retrieved from the stomach of a dwarf sperm whale stranded in Miami in 1987 (Barros *et al.*, 1990). In the period 1987-1994, 1 animal had possible propeller cuts on or near the flukes. ~~Three dwarf sperm whales and 3 undifferentiated *Kogia* spp. stranded in the U.S. Atlantic Ocean during 2002. In each case, signs of human interaction could not be assessed or were not identified~~

Rehabilitation challenges for *Kogia* spp. are numerous due to limited knowledge regarding even the basic biology of these species. Advances in recent rehabilitation success has potential implications for future release and tracking of animals at sea to potentially provide information on distribution, movements and habitat use of these species (Manire *et al.*, 2004).

STATUS OF STOCK

The status of ~~this stock~~ the pygmy sperm whale relative to OSP in the western U.S. Atlantic EEZ is unknown. This species is not listed as endangered or threatened under the Endangered Species Act. There is insufficient information with which to assess population trends. Total fishery-related mortality and serious injury for this stock is less than 10% of the calculated PBR and therefore, can be considered to be insignificant and approaching zero mortality and serious injury rate. This is not a strategic stock.

REFERENCES

- Barlow, J., S. L. Swartz, T. C. Eagle and P. R. Wade. 1995. U.S. Marine Mmammal Sstock Assessments: Guidelines for Preparation, Background, and a Summary of the 1995 Assessments. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-6, ~~73 pp.~~ 73pp. Available from NMFS, Southwest Fisheries Science Center, 8604 La Jolla Shores Drive, La Jolla, CA, 92037-1508.
- Barros, N.B. and D.A. Duffield. 2003. Unraveling the mysteries of Pygmy and Dwarf sperm whales. *Strandings Newsletter of the Southeast U.S. Marine Mammal Stranding Network*. December 2003. NOAA Tech. Memo NMFS-SEFSC-521, 11 pp. Available from NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Drive, Miami, FL 33149.
- Barros, N. B., D. A. Duffield, P. H. Ostrom, D. K. Odell and V. R. Cornish. -1998.- Nearshore vs. offshore ecotype differentiation of *Kogia breviceps* and *K. simus* based on hemoglobin, morphometric and dietary analyses. Abstracts. World Marine Mammal Science Conference Abstracts. Monaco. 20-24 January.
- Barros, N. B., D. K. Odell and G. W. Patton. 1990. Ingestion of plastic debris by stranded marine mammals from Florida. -Page 746. -In: R. S. Shomura, R. S. and M.-L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris. NOAA Tech. Memo. NOAA-TM-NMFS-SWFSC-154.

- Available from NMFS, Southwest Fisheries Science Center, 8604 La Jolla Shores Drive, La Jolla, CA, 92037-1508.
- Buckland, S. T., D. R. Anderson, K. P. Burnham, J. L. Laake, D. L. Borchers and L. Thomas. 2001. *Introduction to distance sampling: estimating abundance of biological populations*. Oxford University Press, New York, 432 pp.
- Caldwell, D. K. and M. C. Caldwell. 1989. Pygmy sperm whale *Kogia breviceps* (de Blainville; 1838): dwarf sperm whale *Kogia simus* Owen, 1866. Pages pp. 235-260. -In: S. H. Ridgway and R. Harrison (editors eds.), *Handbook of marine mammals, Vol. 4: River dolphins and the larger toothed whales*. -Academic Press, San Diego. pp.
- Credle, V. R. 1988. Magnetite and magnetoreception in dwarf and pygmy sperm whales, *Kogia simus* and *Kogia breviceps*. MSc. Thesis. -University of Miami. Coral Gables, FL.
- Duffield, D.A., N.B. Barros, E.O. Espinoza, S. Ploen, F.M.D. Gulland, and J.E. Heyning. 2003. Identifying Pygmy and Dwarf Sperm Whales (Genus *Kogia*) using electrospray ionization mass spectrometry of myoglobin and hemoglobin. *Mar. Mammal Sci.* 19(2):395-399.
- Garrison, L.P. 2003. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2001-2002. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-SEFSC-515, 52 pp. Available from NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Road, Miami, FL 33149.
- Garrison, L.P. and P.M. Richards. 2004. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2003. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-SEFSC-527, 57 pp. Available from NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Road, Miami, FL.
- Hiby, L. 1999. The objective identification of duplicate sightings in aerial survey for porpoise. pp. 179-189. In: G.W. Garner, S. C. Amstrup, J. L. Laake, B. F. J. Manly, L. L. McDonald, and D. G. Robertson (eds.) *Marine mammal survey and assessment methods*. Balkema, Rotterdam. 287 pp.
- Manire, C.A., H.L. Rhinehart, N.B. Barros, L. Byrd, and P. Cunningham-Smith. 2004. An approach to the rehabilitation of *Kogia* spp. *Aquatic Mamm.* 30(2):257-270.
- Mullin, K. D. and G. L. Fulling. 2003. Abundance and distribution of cetaceans in the southern U.S. North Atlantic Ocean during summer 1998. *Fish. Bull.*, U.S. 101:603-613.
- Palka, D. 1995. Abundance estimate of the Gulf of Maine harbor porpoise. pp. 27-50 In: A. Bjørge and G.P. Donovan (eds.) *Biology of the Phocoenids*. Rep. Int. Whal. Commn. (Special Issue 16: 27-50) i-x + 552 pp.
- Palka, D., G. Waring and D. Potter. In review. Abundances of cetaceans and sea turtles in the northwest Atlantic during summer 1995 and 1998. *Fish. Bull.*, U.S. and P.S. Hammond. 2001. Accounting for responsive movement in line-transect estimates of abundance. *Can. J. Fish. Aquat. Sci.* 58: 777-787.
- Thomas, L., J. L. Laake, J. F. Derry, S. T. Buckland, D. L. Borchers, D. R. Anderson, K. P. Burnham, S. Strindberg, S. L. Hedley, F. F. C. Marques, J. H. Pollard and R. M. Fewster. 1998. Distance 3.5. Research Unit for Wildlife Population Assessment, University of St. Andrews, St. Andrews, UK.
- Wade, P.-R., and R.-P. Angliss. -1997.- Guidelines for assessing marine mammal stocks:- Report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-12, 93 pp.
- Willis, P. M. and R. W. Baird. 1998. Status of the dwarf sperm whale, *Kogia simus*, with special reference to Canada. *Can. Field Nat.* 112:114-125. Available from: NOAA, National Marine Fisheries Service, 166 Water St., Woods Hole, MA 02543.
- Wursig, B., T.A. Jefferson, and D.J. Schmidly. 2000. *The marine mammals of the Gulf of Mexico*. Texas A&M University Press, College Station, TX, 256 pp.
- Yeung, C. 2001. Estimates of marine mammal and marine turtle bycatch by the U.S. Atlantic pelagic longline fleet in 1999-2000. NOAA Tech. Memo. NOAA-TM-SEFSC-467, 42 pp. Available from: NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Dr., Miami, FL, 33149.

FRASER'S DOLPHIN (*Lagenodelphis hosei*): Western North Atlantic Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

Fraser's dolphin is distributed worldwide in tropical waters (Perrin *et al.* 1994). A group of an estimated 250 Fraser's dolphins was sighted in waters 3300 m deep in the western North Atlantic off Cape Hatteras during a 1999 vessel survey (Figure 1; Anon. 1999). Fraser's dolphins are assumed to be part of the cetacean fauna of the tropical western North Atlantic. The paucity of sightings is probably due to naturally low abundance compared to other cetacean species. Sightings in the more extensively surveyed northern Gulf of Mexico are uncommon but occur on a regular basis. Fraser's dolphins have been observed in oceanic waters (>200 m) in the northern Gulf of Mexico during all seasons (Leatherwood *et al.* 1993; Hansen *et al.* 1996; Mullin and Hoggard 2000; Mullin and Fulling, 2004). The western North Atlantic population is provisionally being considered one stock for management purposes. Additional morphological, genetic and/or behavioral data are needed to provide further information on stock delineation.

POPULATION SIZE

The numbers of Fraser's dolphins off the U.S. or Canadian Atlantic coast are unknown, and seasonal abundance estimates are not available for this stock, since it was rarely seen in any surveys. A group of an estimated 250 Fraser's dolphins was sighted in waters 3300 m deep in the western North Atlantic off Cape Hatteras during a 1999 vessel survey (Figure 1; Anon. 1999). Abundances have not been estimated from the 1999 vessel survey in western North Atlantic (Anon. 1999); because the sighting was not made during line-transect sampling effort; therefore, the population size of Fraser's dolphins is unknown.

Minimum Population Estimate

The Present data are insufficient to calculate a minimum population estimate for Fraser's dolphins in the western North Atlantic is unknown this stock.

Current Population Trend

There are insufficient data to determine the population trends for this species stock.

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Current and maximum net productivity rates are unknown for this stock. For purposes of this assessment, the maximum net productivity rate was assumed to be 0.04. This value is based on theoretical modeling showing that cetacean populations may not grow at rates much greater than 4% given the constraints of their reproductive history (Barlow *et al.* 1995).

POTENTIAL BIOLOGICAL REMOVAL




Figure 1. Distribution of Fraser's dolphins from SEFSC vessel surveys during 1998-2002. All sightings are shown. Solid lines indicate the 200 and 2000 m isobaths.

Potential ~~b~~Biological ~~r~~Removal level (PBR) is the product of the minimum population size, one-half the maximum ~~net~~ productivity rate, and a recovery factor (MMPA Sec. 3.16 U.S.C. 1362; Wade and Angliss 1997). The minimum population size is unknown. The maximum productivity rate is 0.04, the default value for cetaceans. The “recovery” factor, which accounts for endangered, depleted, threatened stocks, or stocks of unknown status relative to optimum sustainable population (OSP), is assumed to be 0.5 because this stock is of unknown status. PBR for the western North Atlantic Fraser’s dolphin stock is unknown because the minimum population size is unknown.

ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

~~There has been no reported fishing-related~~
Fishery Information

Detailed fishery information is reported in Appendix III. Total annual estimated average fishery-related mortality of a Fraser’s dolphin (Yeung 1999, Yeung 2001).

Fisheries Information

~~The level of past or current, direct, human-caused mortality of~~ and serious injury to this stock during 1999-2003 was zero Fraser’s dolphins in the western North Atlantic is unknown. Pelagic swordfish, tunas, and billfish are the targets of the longline fishery operating in the western North Atlantic. There, as there were no reports of mortality or serious injury to Fraser’s dolphins ~~by this fishery~~ (Yeung 2001; Garrison 2003; Garrison and Richards, 2004).

Other Mortality

~~There were no reported strandings of~~ From 1999-2003, 12 Fraser’s dolphins ~~in the western North Atlantic during 1997-2002.~~

~~Stranding data probably underestimate the extent of fishery-related mortality and serious injury because not all of the marine mammals which die or are seriously injured in fishery interactions wash ashore, not all that wash ashore are discovered, reported or investigated, nor will all of those that do wash ashore necessarily show signs of entanglement or other fishery-interaction. Finally, the level of technical expertise among stranding network personnel varies widely as does the ability to recognize signs of fishery interactions.~~

were reported stranded between Maine and Puerto Rico (Table 1). The total includes 1 animal stranded in Puerto in 1999 and 1 in 2002, and 10 mass stranded live animals in April 2003 in Lee, Florida. There were no indications of human interactions for these stranded animals.

Table 1. Fraser’s dolphin (*Lagenodelphis hosei*) strandings along the U.S. Atlantic coast, 1999-2003

<u>STATE</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>TOTALS</u>
<u>North Carolina</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>
<u>South Carolina</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>
<u>Georgia</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>
<u>Florida</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>10</u>	<u>10</u>
<u>Puerto Rico</u>	<u>1</u>	<u>0</u>	<u>0</u>	<u>1</u>	<u>0</u>	<u>2</u>
<u>TOTALS</u>	<u>1</u>	<u>0</u>	<u>0</u>	<u>1</u>	<u>10</u>	<u>12</u>

~~Florida live mass stranding of 10 animals in Lee, Florida on April 4, 2003~~

STATUS OF STOCK

The status of ~~Fraser’s dolphins~~ melon-headed whales, relative to OPS, in the ~~northern~~ U.S. western North Atlantic, ~~relative to OSP~~, EEZ is unknown. The species is not listed as threatened or endangered under the

Endangered Species Act. There are insufficient data to determine the population ~~trends for this species. The size or trends and PBR cannot be calculated for this stock. No fishery-related mortality and serious injury has been observed since 1999; therefore,~~ total fishery-related mortality and serious injury ~~for this stock is unknown, but assumed to be less than 10% of the calculated PBR and rate~~ can be considered ~~to be~~ insignificant and approaching zero mortality and serious injury ~~rate~~. This is not a strategic stock.

REFERENCES

- Anon. 1999. Cruise results. Summer Atlantic Ocean marine mammal survey. NOAA Ship *Oregon II* cruise 236 (99-05), 4 August - 30 September 1999. Available from SEFSC, 3209 Frederic Street, Pascagoula, MS 39567.
- Barlow, J., S. L. Swartz, T. C. Eagle and P. R. Wade. 1995. U.S. Marine mammal stock assessments: Guidelines for preparation, background, and a summary of the 1995 assessments. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-6, 73pp.
- ~~Hansen, L. J., K. D. Mullin and C. L. Roden. 1995. Estimates of cetacean abundance in the northern Gulf of Mexico from vessel surveys. Southeast Available from NMFS, Southwest Fisheries Science Center, Miami Laboratory, Contribution No. MIA-94/95-25, 98604 La Jolla Shores Drive, La Jolla, CA, 92037-1508.~~
- ~~Garrison, L.P. 2003. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2001-2002. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-SEFSC-515, 52 pp. Available from: NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Dr., Miami, FL, 33149.~~
- ~~Yeung, Garrison, L.P. and P.M. Richards. 2001. Estimates of marine mammals and marine turtle bycatch by in the U.S. Atlantic pelagic longline fleet in 1999-2000 during 2003. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-SEFSC-465, 43 pp. Available from: NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Dr., Miami, FL, 33149.~~
- Hansen, L. J., K. D. Mullin, T. A. Jefferson and G. P. Scott. 1996. Visual surveys aboard ships and aircraft. ~~Pages pp.~~ 55-132. In: R. W. Davis and G. S. Fargion (editors), *Distribution and abundance of marine mammals in the north-central and western Gulf of Mexico: Final report. Volume II: Technical report.* OCS Study MMS 96- 0027. Minerals Management Service, Gulf of Mexico OCS Region, New Orleans.
- Leatherwood, S., T. A. Jefferson, J. C. Norris, W. E. Stevens, L. J. Hansen and K. D. Mullin. 1993. Occurrence and sounds of Fraser's dolphin in the Gulf of Mexico. *Texas J. Sci.* 45(4):349-354.
- ~~Mullin, K. D. and G. L. Fulling. In review. Abundance of cetaceans in the oceanic northern Gulf of Mexico. Mar. Mamm. Sci.~~
- Mullin, K. D. and W. Hoggard. 2000. Visual surveys of cetaceans and sea turtles from aircraft and ships. Pages 111-172. In R. W. Davis, W. E. Evans, and B. Würsig (editors), *Cetaceans, sea turtles and seabirds in the northern Gulf of Mexico: Distribution, abundance and habitat associations. Volume II: Technical report.* OCS Study MMS 96-0027. Minerals Management Service, Gulf of Mexico OCS Region, New Orleans.
- ~~Mullin, K. D. and G. L. Fulling. 2004. Abundance of cetaceans in the oceanic northern Gulf of Mexico, 1996-2001. Mar. Mamm. Sci. 20(4):787-807.~~
- Perrin, W. F., S. Leatherwood and A. Collet. 1994. Fraser's dolphin *Lagenodelphis hosei* (Fraser 1956). ~~Pages pp.~~ 225-240. In: S. H. Ridgway and R. Harrison (editors), *Handbook of marine mammals, Vol. 5: The first book of dolphins.* Academic Press, London, 416 pp.
- Wade, P.-R., and R.-P. Angliss. -1997.- Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, ~~WA Washington.~~ U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-12, ~~93pp.~~
- ~~Yeung, C. 1999. 93 pp. Available from: NOAA, National Marine Fisheries Service, 166 Water St., Woods Hole, MA 02543.~~
- ~~Yeung, C. 2001. Estimates of marine mammal and marine turtle bycatch by the U.S. Atlantic pelagic longline fleet in 1998-1999-2000. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-SEFSC-430, 26 NMFS-SEFSC-467, 43 pp. Available from: NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Dr., Miami, FL, 33149.~~

MELON-HEADED WHALE (*Peponocephala electra*): Western North Atlantic Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

The melon-headed whale is distributed worldwide in tropical to sub-tropical waters (Jefferson *et al.*, 1994). A group of melon-headed whales was sighted during both a 1999 (20 whales) and 2002 (80 whales) vessel survey of the western North Atlantic off of Cape Hatteras, North Carolina in waters >2500 m deep (Figure 1; Anon. 1999; Anon. 2002). Melon-headed whales are assumed to be part of the cetacean fauna of the tropical western North Atlantic. The paucity of sightings is probably due to a naturally low number of groups compared to other cetacean species.

Sightings in the more extensively surveyed northern Gulf of Mexico occur in oceanic waters (Mullin *et al.* 1994; Mullin and Fulling *in review*, 2004). Sightings of melon-headed whales in the northern Gulf of Mexico were documented in all seasons during aerial surveys of the northern Gulf of Mexico between 1992 and 1998 (Hansen *et al.* 1996; Mullin and Hoggard 2000). The western North Atlantic population is provisionally being considered one stock for management purposes. Additional morphological, genetic and/or behavioral data are needed to provide further information on stock delineation.

POPULATION SIZE

The numbers of melon-headed whales off the U.S. or Canadian Atlantic coast are unknown, and seasonal abundance estimates are not available for this stock, since it was rarely seen in any surveys. A group of melon-headed whales was sighted during both a 1999 (20 whales) and 2002 (80 whales) vessel survey of the western North Atlantic off of Cape Hatteras, North Carolina in waters >2500 m deep (Figure 1; Anon. 1999; Anon. 2002). Abundances have not been estimated from the 1999 and 2002 vessel surveys in western North Atlantic (Anon. 1999; Anon. 2002); because the sighting was not made during line-transect sampling effort; therefore the population size of melon-headed whales is unknown.

Minimum Population Estimate

The Present data are insufficient to calculate a minimum population estimate for the western North Atlantic this stock of melon-headed whales unknown.

Current Population Trend

There are insufficient data to determine the population trends for this species stock.

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Current and maximum net productivity rates are unknown for this stock. For purposes of this assessment, the maximum net productivity rate was assumed to be 0.04. This value is based on theoretical modeling showing that cetacean populations may not grow at rates much greater than 4% given the constraints of their reproductive history (Barlow *et al.* 1995).




Figure 1. Distribution of melon-headed whales from SEFSC vessel surveys during 1998-2002. All sightings are shown. Solid lines indicate the 200 and 2000 m isobaths.

POTENTIAL BIOLOGICAL REMOVAL

Potential ~~b~~Biological ~~r~~Removal level (PBR) is the product of the minimum population size, one-half the maximum ~~net~~ productivity rate, and a recovery factor (MMPA Sec. 3.16 U.S.C. 1362; Wade and Angliss 1997). The minimum population size is unknown. The maximum productivity rate is 0.04, the default value for cetaceans. The “recovery” factor, which accounts for endangered, depleted, threatened stocks, or stocks of unknown status relative to optimum sustainable population (OPS), is assumed to be 0.5 because this stock is of unknown status. PBR for the western North Atlantic stock of ~~melon-headed whales~~ is unknown because the minimum population size is unknown.

ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

~~There has been no reported fishing-related mortality of a melon-headed whale between 1997 and 2001 (Yeung 1999; Yeung 2001).~~

~~Fisheries~~Fishery Information

~~The level of past or current, direct, human-caused mortality of~~ Detailed fishery information is reported in Appendix III. Total annual estimated average fishery-related mortality and serious injury to this stock during 1999-2003 was zero ~~melon-headed whales in the western North Atlantic is unknown. There has historically been some take of this species in small cetacean fisheries in the Caribbean (Caldwell *et al.* 1976). Pelagic swordfish, tunas, and billfish are the targets of the longline fishery operating in the western North Atlantic. There, as there~~ were no reports of mortality or serious injury to melon-headed whales ~~by this fishery (Yeung 2001; Garrison 2003; Garrison and Richards, 2004).~~

Other Mortality

From 1999-2003, 1 melon-headed whale was reported stranded in Puerto Rico. There was one additional reported stranding of a melon-headed whale in the western North Atlantic between 1997 and 2002, ~~though there was n~~. No evidence of human interaction was apparent for this ~~either~~ stranded animal. Stranding data probably underestimate the extent of fishery-related mortality and serious injury because not all of the marine mammals which die or are seriously injured in fishery interactions wash ashore, not all that wash ashore are discovered, reported or investigated, nor will all of those that do wash ashore necessarily show signs of entanglement or other fishery-interaction. Finally, the level of technical expertise among stranding network personnel varies widely as does the ability to recognize signs of fishery interactions.

STATUS OF STOCK

The status of melon-headed whales, relative to OPS, in the western North Atlantic, ~~relative to OSP, EEZ~~ is unknown. The species is not listed as threatened or endangered under the Endangered Species Act. There are insufficient data to determine the population trends for this species. The size or trends and PBR cannot be calculated for this stock. No fishery-related mortality and serious injury has been observed since 1999; therefore, total fishery-related mortality and serious injury for this stock is unknown, but assumed to be less than 10% of the calculated PBR and rate can be considered ~~to be~~ insignificant and approaching zero mortality and serious injury ~~rate~~. This is not a strategic stock.

REFERENCES

- Anon. 1999. Cruise results. Summer Atlantic Ocean marine mammal survey. NOAA Ship *Oregon II* cruise 236 (99-05), 4 August - 30 September 1999. Available from SEFSC, 3209 Frederic Street, Pascagoula, MS 39567.
- Anon. 2002. Cruise results. Mid-Atlantic cetacean survey. NOAA Ship *Gordon Gunter* cruise GU-02-01, 6 February - 8 April 2002. Available from SEFSC, 3209 Frederic Street, Pascagoula, MS 39567.
- Barlow, J., S. L. Swartz, T. C. Eagle and P. R. Wade. 1995. U.S. Marine mammal stock assessments: Guidelines for preparation, background, and a summary of the 1995 assessments. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-6, 73pp.
- ~~Caldwell, D. K., M. C. Caldwell and R. V. Walker. 1976. First records for Fraser's dolphin (*Lagenodelphis hosei*) in the Atlantic and the melon-headed whale (*Peponocephala electra*) in the western Atlantic. *Cetology* 25:1-4.~~
- ~~Hansen, L. J., K. D. Mullin and C. L. Roden. 1995. Estimates of cetacean abundance in the northern Gulf of~~

- ~~Mexico from vessel surveys. Southeast Available from NMFS, Southwest Fisheries Science Center, Miami Laboratory, Contribution No. MIA-94/95-25, 9 pp. 8604 La Jolla Shores Drive, La Jolla, CA, 92037-1508.~~
- Garrison, L.P. 2003. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2001-2002. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-SEFSC-515, 52 pp. Available from: NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Dr., Miami, FL, 33149.
- Road, Miami, FL 33149.
- ~~Yeung~~Garrison, L.P. and P.M. Richards. 2001. Estimates~~Estimated bycatch~~ of marine mammals and marine turtle bycatch by~~in~~ the U.S. Atlantic pelagic longline fleet in 1999-2000~~during 2003~~. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-SEFSC-465, 43 pp. Available from: NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Dr., Miami, FL, 33149.
- Hansen, L. J., K. D. Mullin, T. A. Jefferson and G. P. Scott. 1996. Visual surveys aboard ships and aircraft. Pages 55-132. In: R. W. Davis and G. S. Fargion (editors), *Distribution and abundance of marine mammals in the north-central and western Gulf of Mexico: Final report. Volume II: Technical report*. OCS Study MMS 96-0027. Minerals Management Service, Gulf of Mexico OCS Region, New Orleans.
- Jefferson, T. A., S. Leatherwood, and M. A. Weber. 1994. Marine mammals of the world. FAO, Rome, 320 pp.
- Mullin, K. D. ~~and G. L. Fulling. In review. Abundance of cetaceans in the oceanic northern Gulf of Mexico. Mar. Mamm. Sci.~~
- Mullin, K. D., T. A. Jefferson, L. J. Hansen and W. Hoggard. 1994. First sightings of melon-headed whales (*Peponocephala electra*) in the Gulf of Mexico. Mar. Mam. Sci. 10(3): 342-348.
- Mullin, K. D. and W. Hoggard. 2000. Visual surveys of cetaceans and sea turtles from aircraft and ships. Pages 111-172. In: R. W. Davis, W. E. Evans, and B. Würsig (editors), *Cetaceans, sea turtles and seabirds in the northern Gulf of Mexico: Distribution, abundance and habitat associations. Volume II: Technical report*. OCS Study MMS 96-0027. Minerals Management Service, Gulf of Mexico OCS Region, New Orleans.
- Mullin, K. D. and G. L. Fulling. 2004. Abundance of cetaceans in the oceanic northern Gulf of Mexico, 1996-2001. Mar. Mamm. Sci. 20(4):787-807.
- Wade, P.-R., and R.-P. Angliss. -1997.- Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, ~~WA~~Washington. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-12, ~~93~~pp.
- Yeung, C. 1999. 93 pp. Available from: NOAA, National Marine Fisheries Service, 166 Water St., Woods Hole, MA 02543.
- Yeung, C. 2001. Estimates of marine mammal and marine turtle bycatch by the U.S. Atlantic pelagic longline fleet in 1998-1999-2000. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-SEFSC-430, 26 NMFS-SEFSC-467, 43 pp. Available from: NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Dr., Miami, FL, 33149.

PANTROPICAL SPOTTED DOLPHIN (*Stenella attenuata*): Western North Atlantic Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

There are two species of spotted dolphin in the Western Atlantic —Ocean, the Atlantic spotted dolphin, *Stenella frontalis*, formerly *S. plagiodon* (Perrin *et al.* 1987), and the pantropical spotted dolphin, *S. attenuata*. These species are (Perrin *et al.* 1987). The Atlantic spotted dolphin occurs in two forms which may be distinct sub-species (Perrin *et al.* 1987, 1994; Rice 1998): the large, heavily spotted form which inhabits the continental shelf and is usually found inside or near the 200m isobath; and the smaller, less spotted island and offshore form which occurs in the Atlantic Ocean but is not known to occur in the Gulf of Mexico (Fulling *et al.* 2003; Mullin and Fulling 2003; Mullin and Fulling 2004). Where they co-occur, the offshore form of the Atlantic spotted dolphin and the pantropical spotted dolphin can be difficult to differentiate at sea.

The pantropical spotted dolphin is distributed worldwide in tropical and some sub-tropical oceans (Perrin *et al.* 1987; Perrin and Hohn 1994). Sightings of this species in the northern Gulf of Mexico occur over the deeper waters, and rarely over the continental shelf or continental shelf edge (Mullin *et al.* 1991; SEFSC, unpublished data). Pantropical spotted dolphins were seen in all seasons during recent seasonal aerial surveys of the northern Gulf of Mexico, and during recent winter aerial surveys offshore of the southeastern USA. Atlantic coast (SEFSC unpublished data). Some of the Pacific populations have been divided into different geographic stocks based on morphological characteristics (Perrin *et al.* 1987; Perrin and Hohn 1994); however, there is no information on stock differentiation in the Atlantic population.

POPULATION SIZE

Total numbers of pantropical spotted dolphins off the USA, or Canadian Atlantic coast are unknown, although three estimates from selected regions of the habitat do exist for select time periods. Because *S. frontalis* and *S. attenuata* are difficult to differentiate at sea, the reported abundance estimates, prior to 1998, are for both species of spotted dolphins combined. Sightings were almost exclusively in the continental shelf edge and continental slope areas west of Georges Bank (Figure 1). concentrated in the southeastern edge of Georges Bank, along the Florida shelf and to a more limited degree the Florida slope waters, and offshore in Gulf Stream waters southeast of Cape Hatteras (Fig. 1).

An abundance of 6,107 undifferentiated spotted dolphins (CV=0.27) was estimated from an aerial survey program conducted from 1978 to 1982 on the continental, shelf and shelf edge waters between Cape Hatteras, North Carolina and Nova Scotia (CETAP 1982). As recommended in the GAMS Workshop Report (Wade and Angliss 1997), estimates older than eight years are deemed unreliable, therefore should not be used for PBR determinations. Further, due to changes in survey methodology these data should not be used to make comparisons to more current estimates.

An abundance of 4,772 (CV=1.27) undifferentiated spotted dolphins was estimated from a July to September 1995 sighting survey conducted by two ships and an airplane that covered waters from Virginia to the mouth of the Gulf of St. Lawrence (Table 1; Palka *et al.* in review Unpubl. Ms.). Total track line length was 32,600 km. The ships covered waters between the 50 and 1000 fathom depth contour lines, the northern edge of the Gulf Stream, and the northern Gulf of Maine/Bay of Fundy region. The airplane covered waters in the mid-Atlantic from the coastline to the 50 fathom depth contour line, the southern Gulf of Maine, and shelf waters off Nova Scotia from the coastline to the 1000 fathom depth contour line. Data collection and analysis methods used were described in Palka (1996).

An abundance of 343 (CV=1.03) for pantropical spotted dolphins was estimated from a line transect sighting survey conducted during July 6 to September 6, 1998 by a ship and plane that surveyed 15,900 km of track line




Figure 1. Distribution of pantropical spotted dolphin sightings from NEFSC and SEFSC shipboard and aerial surveys during the summer in 1998 and 2004. Isobaths are at 100m and 1,000m.

in waters north of Maryland (38° N) (Figure 1; Palka *et al.* ~~in review~~ Unpubl. Ms.). Shipboard data were analyzed using the modified direct duplicate method (Palka 1995) that accounts for school size bias and $g(0)$, the probability of detecting a group on the track line. Aerial data were not corrected for $g(0)$.

An abundance of ~~12,774~~ 747 (CV=0.~~57~~56) for pantropical spotted dolphins was estimated from a shipboard line transect sighting survey conducted between 8 July and 17 August 1998 that surveyed ~~5,457 km~~ 163 km of track line in waters south of Maryland (38°N) (Figure 1; Mullin ~~in review~~); and Fulling 2003). This estimate is a recalculation of the same data reported in previous SARs. For more details see Mullin and Fulling (2003). Abundance estimates were made using the program DISTANCE (Buckland *et al.* 1993; Laake *et al.* 1993~~2003~~) where school size bias and ship attraction were accounted for.

~~The best available~~ An abundance of zero for Atlantic spotted dolphins was estimated from a line transect sighting survey conducted during June 12 to August 4, 2004 by a ship and plane that surveyed 10,761 km of track line in waters north of Maryland (about 38° N) to the Bay of Fundy (about 45° N) (Figure 1; Palka unpubl.), as no dolphins of this species were observed). Shipboard data were collected using the two independent team line transect method and analyzed using the modified direct duplicate method (Palka 1995) accounting for biases due to school size and other potential covariates, reactive movements (Palka and Hammond 2001), and $g(0)$, the probability of detecting a group on the track line. Aerial data were collected using the Hiby circle-back line transect method (Hiby 1999) and analyzed accounting for $g(0)$ and biases due to school size and other potential covariates (Figure 1; Palka unpubl.).

A survey of the U.S. Atlantic outer continental shelf and continental slope (water depths = 50m) between 27.5 – 38 °N latitude was conducted during June-August, 2004. The survey employed two independent visual teams searching with 50x bigeye binoculars. Survey effort was stratified to include increased effort along the continental shelf break and Gulf Stream front in the mid-Atlantic. The survey included 5,659 km of trackline, and there were a total of 473 cetacean sightings. Sightings were most frequent in waters North of Cape Hatteras, North Carolina along the shelf break. Data were analyzed to correct for visibility bias ($g(0)$) and group-size bias employing line transect distance analysis and the direct duplicate estimator (Palka 1995; Buckland *et al.*, 2001). The resulting abundance estimate for pantropical Atlantic spotted dolphins is the sum of the estimates from the two 1998 USA Atlantic surveys, 13,117 (CV=0.56), where the estimate between Florida and Maryland was 4,439 (CV=0.49).

At their November 1999 meeting, the Atlantic SRG recommended that, without a genetic determination of stock structure, the abundance estimates for the coastal and offshore forms should be combined. There remains debate over how distinguishable both species are at sea, though in the waters south of Cape Hatteras identification to species is made with very high certainty. This does not, however, account for the potential for a mixed species herd, as has been recorded for several dolphin assemblages. Pending further genetic studies for clarification of this problem, a single species abundance estimate will be used as the best estimate of abundance, combining species specific data from the northern USA Atlantic is 343 (CV=1.03) and from the southern USA Atlantic is 12,774 (CV=0.57) as well as southern portions of the species' ranges. This joint estimate is considered best because together these two surveys have the most complete coverage of the species' habitat. The best 2004 abundance estimate for pantropical spotted dolphins is the sum of the estimates from the two 2004 western U.S. Atlantic surveys, 4,439 (CV=0.49), where the estimate from the northern U.S. Atlantic is 0, and from the southern U.S. Atlantic is 4,439 (CV=0.49).

Table 1. Summary of abundance estimates for both undifferentiated spotted dolphins (1995) and differentiated pantropical spotted dolphins (1998, 2004). Month, year, and area covered during each abundance survey, and resulting abundance estimate (N_{best}) and coefficient of variation (CV).

Month/Year	Area	N_{best}	CV
Jul-Sep 1995 Virginia to Gulf of St. Lawrence 4,772¹ 1.27 Jul-Sep 1998	Maryland to Gulf of St. Lawrence	343 ^{2,1}	1.03
Jul-Aug 1998	Florida to Maryland	12, 774 ^{2,7} 47¹	0. 57 <u>56</u>
Jul-Aug 1998	<u>Florida to</u> Gulf of St. Lawrence <u>(COMBINED)</u>	<u>13,090</u>	<u>0.55</u>
<u>Jun-Aug 2004</u>	<u>Maryland to Bay of Fundy</u>	<u>0</u>	<u>0</u>
<u>Jun-Aug 2004</u>	Florida <u>to Maryland</u>	<u>4,439</u>	<u>0.48</u>

<u>Jun-Aug 2004</u>	<u>Florida to Bay of Fundy (COMBINED)</u>	<u>13,117³ 439</u>	<u>0.5648</u>
---------------------	---	-----------------------------------	---------------

¹ Because of uncertain species identification in the 1995 survey, all spotted dolphins were lumped together.

²

¹ This represents the first estimates for pantropical spotted dolphin.

³ This represents the combined estimates for both regions.

Minimum Population Estimate

The minimum population estimate is the lower limit of the two-tailed 60% confidence interval of the log-normally distributed best abundance estimate. This is equivalent to the 20th percentile of the log-normal distribution as specified by Wade and Angliss (1997). The best estimate of abundance for pantropical spotted dolphins is 13,117³439 (CV=0.5649). The minimum population estimate for pantropical spotted dolphins is 83,450 (CV=0.1056).

Current Population Trend

There are insufficient data to determine the population trends for this species, because prior to 1998 spotted dolphins (*Stenella* spp) were not differentiated during surveys.

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Current and maximum net productivity rates are unknown for this stock. For purposes of this assessment, the maximum net productivity rate was assumed to be 0.04. This value is based on theoretical modeling showing that cetacean populations may not grow at rates much greater than 4% given the constraints of their reproductive life history (Barlow *et al.* 1995).

POTENTIAL BIOLOGICAL REMOVAL

Potential Biological Removal (PBR) is the product of minimum population size, one-half the maximum productivity rate, and a "recovery" factor (MMPA Sec. 3, 16 U.S.C. 1362; Wade and Angliss 1997). The minimum population size for pantropical spotted dolphins is 83,450 (CV=0.5649). The maximum productivity rate is 0.04, the default value for cetaceans. The "recovery" factor, which accounts for endangered, depleted, threatened stocks, or stocks of unknown status relative to optimum sustainable population (OSP) is assumed to be 0.5 because this stock is of unknown status. PBR for pantropical dolphins is 8430.

ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

Total annual estimated average fishery-related mortality or serious injury to this stock during 1996-2000 was 0 undifferentiated spotted dolphin (*Stenella* sp.).

Fisheries Fishery Information

No spotted dolphin mortalities were observed in 1977-1991 foreign fishing activities. Data on current incidental takes in USA fisheries are available from several sources. In 1986, NMFS established a mandatory self-reported fisheries information system for large pelagic fisheries. Data files are maintained at the Southeast Fisheries Science Center (SEFSC). The Northeast Fisheries Science Center (NEFSC) Fisheries Observer Program was initiated in 1989 and since that year several fisheries have been covered by the program. In late 1992 and in 1993, the SEFSC provided observer coverage of pelagic longline vessels fishing off the Grand Banks (Tail of the Banks) and provides observer coverage of vessels fishing south of Cape Hatteras. Total

Detailed fishery information is reported in Appendix III. Total annual estimated average fishery-related mortality and/or serious injury cannot be estimated separately for the two species of to this stock during 1999-2003 was zero pantropical spotted dolphins, as there were no reports of mortality or serious injury to pantropical spotted dolphins in the USA Atlantic Exclusive Economic Zone (EEZ) because of the uncertainty in species identification by fishery observers. The Atlantic Scientific Review Group advised adopting the risk-averse strategy of assuming that either species might have been subject to the observed fishery-related mortality and serious injury.

(Yeung 2001; Garrison 2003; Garrison and Richards, 2004).

Earlier Interactions

No spotted dolphin mortalities were observed in 1977-1991 foreign fishing activities. Bycatch has been observed by NMFS Sea Samplers in the pelagic drift gillnet and pelagic longline fisheries, but no mortalities or serious injuries have been documented in the pelagic pair trawl, Northeast multispecies sink gillnet, mid-Atlantic coastal gillnet, and North Atlantic bottom trawl fisheries; and no takes have been documented in a review of Canadian gillnet and trap fisheries (Read 1994).

Pelagic Drift Gillnet

The estimated total number of hauls in the pelagic drift gillnet fishery increased from 714 in 1989 to 1,144 in 1990; thereafter, with the introduction of quotas, effort was severely reduced. The estimated number of hauls in 1991, 1992, 1993, 1994, 1995, 1996, and 1998 were 233, 243, 232, 197, 164, 149, 113 respectively. In 1996 and 1997, NMFS issued management regulations which prohibited the operation of this fishery in 1997. Further, in January 1999 NMFS issued a Final Rule to prohibit the use of driftnets (*i.e.*, permanent closure) in the North Atlantic

swordfish fishery (50 CFR Part 630). Fifty-nine different vessels participated in this fishery at one time or another between 1989 and 1993. Since 1994, between 10 and 13 vessels have participated in the fishery. Observer coverage, expressed as percent of sets observed, was 8% in 1989, 6% in 1990, 20% in 1991, 40% in 1992, 42% in 1993, 87% in 1994, 99% in 1995, 64% in 1996, and 99% in 1998. Effort was concentrated along the southern edge of Georges Bank and off Cape Hatteras. Examination of the species composition of the catch and locations of the fishery throughout the year, suggested that the pelagic drift gillnet fishery be stratified into two strata, a southern or winter stratum, and a northern or summer stratum. Estimates of the total bycatch from 1989 to 1993 were obtained using the aggregated (pooled 1989-1993) catch rates, by strata (Northridge 1996). Estimates of total annual bycatch for 1994 and 1995 were estimated from the sum of the observed caught and the product of the average bycatch per haul and the number of unobserved hauls as recorded in self-reported fisheries information. Variances were estimated using bootstrap re-sampling techniques. Forty-nine undifferentiated spotted dolphin mortalities were observed in the drift gillnet fishery between 1989 and 1998 and occurred northeast of Cape Hatteras within the ~~183 m~~ 183m isobath in February-April, and near Lydonia Canyon in October. Six whole animal carcasses that were sent to the Smithsonian were identified as Pantropical spotted dolphins (*S. attenuata*). The remaining animals were not identified to species. Estimated annual mortality and serious injury attributable to this fishery (CV in parentheses) was 25 in 1989 (.65), 51 in 1990 (.49), 11 in 1991 (.41), 20 in 1992 (0.18), 8.4 in 1993 (0.40), 29 in 1994 (0.01), 0 in 1995, 2 in 1996 (0.06), no fishery in 1997, and 0 in 1998.

Pelagic Longline

The pelagic longline fishery operates in the USA U.S. Atlantic (-including Caribbean) and Gulf of Mexico EEZ (SEFSC unpublished data). Interactions between the pelagic longline fishery and spotted dolphins have been reported; however, a vessel may fish in more than one statistical reporting area and it is not possible to separate estimates of fishing effort other than to subtract Gulf of Mexico effort from Atlantic fishing effort, which includes the Caribbean Sea. ~~This fishery has been monitored with about 5% observer coverage, in terms of trips observed, since 1992. Total effort, excluding the Gulf of Mexico, for the pelagic longline fishery, based on mandatory self-reported fisheries information, was 11,279 sets in 1991, 9,869 sets in 1992, 9,862 sets in 1993, 9,481 sets in 1994, 10,129 sets in 1995, 9,885 sets in 1996, 8,023 sets in 1997, and 6,675 sets in 1998 (Cramer 1994; Scott and Brown 1997; Johnson *et al.* 1999; Yeung, 1999a). Since 1992, this fishery has been monitored with about 5% observer coverage, in terms of trips observed, within every statistical reporting area within the US Atlantic EEZ and beyond. Off the US Atlantic coast, the fishery has been observed from January to March off Cape Hatteras, in May and June in the entire mid-Atlantic, and in July through December in the mid-Atlantic Bight and off Nova Scotia. The 1994-1998, estimated take was based on a revised analysis of the observed incidental take and self-reported incidental take and effort data, and replaces previous estimates for the 1992-1993 and 1994-1995 periods (Cramer 1994; Scott and Brown 1997; Johnson *et al.* 1999; Yeung 1999b). Further, Yeung (1999b) revised the 1992-1997 fishery mortality estimates in Johnson *et al.* (1999) to include seriously injured animals. The 1998 bycatch estimates were from Yeung (1999a). Most of the estimated marine mammal bycatch was from US Atlantic EEZ waters between South Carolina and Cape Cod (Johnson *et al.* 1999). Excluding the Gulf of Mexico where ~~one~~ 1 animal was hooked and released alive, no pantropical spotted dolphin bycatches were observed ~~for 1992-1998~~ during 1999-2003.~~

Other Mortality

From ~~1995-1998~~ 1999-2003, ~~158~~ pantropical spotted dolphins were stranded between North Carolina and Florida Puerto Rico (NMFS unpublished data). The ~~158~~ mortalities includes the ~~1996 mass stranding of 11 animals in Florida (NMFS unpublished data)~~ 4 animals stranded in Florida in 1999, 1 animal stranded in North Carolina and 1 in Florida in both 2002 and 2003. There were no documented signs of human interactions in any of these strandings.

Table 2. Pantropical spotted dolphin (*Stenella attenuata*) strandings along the U.S. Atlantic coast, 1999-2003

<u>STATE</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>TOTALS</u>
<u>North Carolina</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>
<u>South Carolina</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>
<u>Georgia</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>
<u>Florida</u>	<u>4</u>	<u>0</u>	<u>1</u>	<u>1</u>	<u>0</u>	<u>6</u>
<u>Puerto Rico</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>
<u>TOTALS</u>	<u>4</u>	<u>0</u>	<u>1</u>	<u>1</u>	<u>0</u>	<u>6</u>

STATUS OF STOCK

The status of pantropical spotted dolphins, relative to OSP in the US western U.S. Atlantic EEZ is unknown.

The species is not listed as threatened or endangered under the Endangered Species Act. There are insufficient data to determine the population trends for this species. Total fishery-related mortality and serious injury for this stock is less than 10% of the calculated PBR and, therefore, can be considered to be insignificant and approaching zero mortality and serious injury rate. Average annual fishery-related mortality and serious injury does not exceed the PBR; therefore, this is not a strategic stock

REFERENCES

- Barlow, J., S. L. Swartz, T. C. Eagle, and P. R. Wade. 1995. U.S. Marine Mammal Stock Assessments: Guidelines for Preparation, Background, and a Summary of the 1995 Assessments. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-6, 73 pp. Available from NMFS, Southwest Fisheries Science Center, 8604 La Jolla Shores Drive, La Jolla, CA, 92037-1508.
- Buckland, S. T., D. R. Anderson, K. P. Burnham, and J. L. Laake. 1993. *Distance 3.52*. Borchers and L. Thomas. 2001. *Introduction to distance sampling: estimating abundance of biological populations*. Chapman and Hall/Oxford University Press, New York, NY. 443 pp.
- CETAP. 1982. A characterization of marine mammals and turtles in the mid- and north Atlantic areas of the U.S. outer continental shelf. Cetacean and Turtle Assessment Program, University of Rhode Island. Final Report, Contract AA51-C78-48, Bureau of Land Management, Washington, DC. 538 pp.
- Cramer, J. 1994. Large pelagic logbook newsletter - 1993. NOAA Tech. Memo. NMFS-SEFSC-352, 19 pp. Garrison, L.P. 2003. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2001-2002. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-SEFSC-515, 52 pp. Available from NMFS, Southeast Fisheries Science Center, Miami, FL.
- Johnson, D. R., C. A. Brown, and C. Yeung. 1999. Estimates of marine mammal and marine turtle catch by the U.S. Atlantic pelagic longline fleet in 1992-1997. SEFSC PRD-98/99-03. 68 pp.
- Laake, J. L., S. T. Buckland, D. R. Anderson and K. P. Burnham. 1993. DISTANCE user's guide, V2.0. Colorado Cooperative Fish & Wildlife Research Unit, Colorado State University, Ft. Collins, Colorado, 72 pp. during 2003. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-SEFSC-527, 57 pp. Available from NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Road, Miami, FL 33149.
- Hiby, L. 1999. The objective identification of duplicate sightings in aerial survey for porpoise. pp. 179-189. In: G.W. Garner, S. C. Amstrup, J. L. Laake, B. F. J. Manly, L. L. McDonald, and D. G. Robertson (eds.) *Marine mammal survey and assessment methods*. Balkema, Rotterdam. 287 pp.
- Mullin, K. D. (in review) and G. L. Fulling. 2003. Abundance and distribution of cetaceans in the southern U.S. Atlantic Ocean during summer 1998. *Fish. Bull., U.S.* 101:603-613.
- Mullin, K., W. Hoggard, C. Roden, R. Lohoefer, C. Rogers and B. Taggart. 1991. Cetaceans on the upper continental slope in the north-central Gulf of Mexico. OCS Study/MMS 91-0027. U.S. Dept. of the Interior, Minerals Management Service, Gulf of Mexico OCS Regional Office, New Orleans, Louisiana, 108 pp.
- Northridge, S. 1996. Estimation of cetacean mortality in the U.S. Atlantic swordfish and tuna drift gillnet and pair trawl fisheries. Final report to the Northeast Fisheries Science Center, Contract No. 40ENNF500160.
- Palka, D. 1995. Abundance estimate of the Gulf of Maine harbor porpoise. pp. 27-50 In: A. Bjørge and G.P. Donovan (eds.) *Biology of the Phocoenids*. Rep. Int. Whal. Commn. (Special Issue 16:27-50) I-x + 552 pp.
- Palka, D. 1996. Update on abundance of Gulf of Maine/Bay of Fundy harbor porpoises. NOAA-NMFS-NEFSC. Ref. Doc. 96-04; 37p. Available from: National Marine Fisheries Service, 166 Water Street, Woods Hole, MA 02543-1026.
- Palka, D., G. Waring, and D. Potter. (in review). Abundances of cetaceans and sea turtles in the northwest Atlantic during summer 1995 and 1998. *Fish. Bull., U.S.* and P.S. Hammond. 2001. Accounting for responsive movement in line transect estimates of abundance. *Can. J. Fish. Aquat. Sci.* 58: 777-787.
- Perrin, W. F., E. D. Mitchell, J. G. Mead, D. K. Caldwell, M. C. Caldwell, P. J. H. van Bree, and W. H. Dawbin. 1987. Revision of the spotted dolphins, *Stenella* sp. *Mar. Mammal Sci.* 3(2): 99-170.
- Perrin, W. F. and A. A. Hohn. 1994. Pantropical spotted dolphin *Stenella attenuata*. Pp. 71-98 in: S. H. Ridgway and R. Harrison (eds.). Handbook of marine mammals, Vol. 5: The first book of dolphins. Academic Press, San Diego, 418 pp.
- Read, A. J. 1994. Interactions between cetaceans and gillnet and trap fisheries in the northwest Atlantic. pp. 133-147. In: W.F. Perrin, G.P. Donovan and J. Barlow (eds.) *Gillnets and Cetaceans*. Rep. Int. Whal. Commn. (Special Issue 15). I-x + 629 pp.
- Scott, G. P. and C. A. Brown. 1997. Estimates of marine mammal and marine turtle catch by the U.S. Atlantic pelagic longline fleet in 1994-1995. Miami Laboratory Contribution MIA-96/97-28
- Wade, P.-R., and R.-P. Angliss. 1997. Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-012, 93 pp.
- Waring, G. T., D. L. Palka, P. J. Clapham, S. Swartz, M. C. Rossman, T. V. N. Cole, L. J. Hansen, K. D. Bisack, K. D. Mullin, R. S. Wells, D. K. Odell, and N. B. Barros. 1999. U.S. Atlantic and Gulf of Mexico marine mammal stock assessments - 1999. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-NE-153, 196pp.
- Yeung, C. 1999a. Available from: NOAA, National Marine Fisheries Service, 166 Water St., Woods Hole, MA

02543.
Yeung, C. 2001. Estimates of marine mammal and marine turtle bycatch by the U.S. Atlantic pelagic longline fleet in ~~1998~~ 1999-2000. U.S. Dep. Commer., NOAA Tech. Memo. ~~NMFS-SEFSC-430, 26~~ NMFS-SEFSC-467, 43 pp. Available from: NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Dr., Miami, FL, 33149.

~~Yeung, C. 1999b. Revised Mortality Estimates of Marine Mammal Bycatch in 1992-1997 based on Serious Injury Guidelines. NOAA Tech. Memo. NMFS-SEFSC-429, 23 pp. NMFS, Southeast Fisheries Science Center, Miami, FL.~~

PYGMY KILLER WHALE (*Feresa attenuata*): Western North Atlantic Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

The pygmy killer whale is distributed worldwide in tropical to sub-tropical waters (Jefferson *et al.* 1994). A group of six pygmy killer whales was sighted during a 1992 vessel survey of the western North Atlantic off of Cape Hatteras, North Carolina, in waters >1500 m deep (Hansen *et al.* 1994), but this species was not sighted during subsequent surveys in 1998, 1999 or 2002 (Anon. 1999; Anon. 2002; Mullin and Fulling 2003). Pygmy killer whales are assumed to be part of the cetacean fauna of the tropical western North Atlantic. The paucity of sightings is probably due to a naturally low number of groups compared to other cetacean species. Sightings in the more extensively surveyed northern Gulf of Mexico occur in oceanic waters (Mullin *et al.* 1994; Mullin and Fulling, in review 2004). Sightings of pygmy killer whales were documented in all seasons during aerial surveys of the northern Gulf of Mexico between 1992 and 1998 (Hansen *et al.* 1996; Mullin and Hoggard 2000). The western North Atlantic population is provisionally being considered one stock for management purposes. Additional morphological, genetic and/or behavioral data are needed to provide further information on stock delineation.

POPULATION SIZE

The numbers of pygmy killer whales off the U.S. or Canadian Atlantic coast are unknown, and seasonal abundance estimates are not available for this stock, since it was rarely seen in any surveys. A group of 6 pygmy killer whales was sighted during a 1992 vessel survey of the western North Atlantic off of Cape Hatteras, North Carolina, in waters >1500 m deep (Hansen *et al.* 1994), but this species was not sighted during subsequent surveys (Anon. 1999; Anon. 2002; Mullin and Fulling 2003). Abundance was not estimated for pygmy killer whales from the 1992 vessel survey in the western North Atlantic (Hansen *et al.* 1994) because the sighting was not made during line-transect sampling effort; therefore, the population size of pygmy killer whales is unknown.

Minimum Population Estimate

The minimum population estimate is the lower limit of the two-tailed 60% confidence interval of the log-normal distributed abundance estimate. This is equivalent to the 20th percentile of the log-normal distributed abundance estimate as specified by Wade and Angliss (1997). The best estimate of abundance for pygmy killer whales is unknown. The present data are insufficient to calculate a minimum population estimate for the western North Atlantic this stock of pygmy killer whales is unknown.

Current Population Trend

There are insufficient data to determine the population trends for this speciesstock.

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Current and maximum net productivity rates are unknown for this stock. For purposes of this assessment, the maximum net productivity rate was assumed to be 0.04. This value is based on theoretical modeling showing that cetacean populations may not grow at rates much greater than 4% given the constraints of their reproductive history (Barlow *et al.* 1995).

POTENTIAL BIOLOGICAL REMOVAL

Potential Biological Removal level (PBR) is the product of the minimum population size, one-half the maximum net productivity rate, and a recovery factor (MMPA Sec. 3.16 U.S.C. 1362; Wade and Angliss 1997). The minimum population size is unknown. The maximum productivity rate is 0.04, the default value for cetaceans. The “recovery” factor, which accounts for endangered, depleted, threatened stocks, or stocks of unknown status relative to optimum sustainable population (OPS), is assumed to be 0.5 because this stock is of unknown status. PBR for the western North Atlantic stock of pygmy killer whales is unknown.

ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

There has been no reported fishing-related mortality of a pygmy killer whale between 1997 and 2001 (Yeung 1999; Yeung 2001).

Fisheries

The level of past or current, direct, human-caused mortality of Detailed fishery information is reported in Appendix III. Total annual estimated average fishery-related mortality and serious injury to this stock during 1999-2003 was zero pygmy killer whales, as there were no reports of mortality or serious injury to pygmy killer whales in the western North Atlantic is unknown (Yeung 2001; Garrison 2003; Garrison and Richards, 2004).

There has historically been some take of this species in small cetacean fisheries in the Caribbean (Caldwell and Caldwell 1971). Pelagic swordfish, tunas, and billfish are the targets of the longline fishery operating in the western North Atlantic. There were no reports of mortality or serious injury to pygmy killer whales by this fishery.

Other Mortality

There were three reported strandings of From 1999-2003, 2 pygmy killer whales in the western North Atlantic between 1997 and 2002. There was no evidence were reported stranded between Maine and Puerto Rico (Table 1). The total includes 1 animal stranded in South Carolina and 1 in Georgia in 2003, though there were no indications of human interactions for these stranded animals. Stranding data probably underestimate the extent of fishery-related mortality and serious injury because not all of the marine mammals which die or are seriously injured in fishery interactions wash ashore, not all that wash ashore are discovered, reported or investigated, nor will all of those that do wash ashore necessarily show signs of entanglement or other fishery interaction. Finally, the level of technical expertise among stranding network personnel varies widely as does the ability to recognize signs of fishery interactions.

Table 1. Pygmy killer whale (*Feresa attenuata*) strandings along the U.S. Atlantic coast, 1999-2003

<u>STATE</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>TOTALS</u>
<u>North Carolina</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>
<u>South Carolina</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>1</u>	<u>1</u>
<u>Georgia</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>1</u>	<u>1</u>
<u>Florida</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>
<u>Puerto Rico</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>
<u>TOTALS</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>2</u>	<u>2</u>

STATUS OF STOCK

The status of pygmy killer whales, relative to OPS, in the U.S. western North Atlantic, relative to OSP, EEZ is unknown. The species is not listed as threatened or endangered under the Endangered Species Act. There are insufficient data to determine the population trends for this species. The size or trends and PBR cannot be calculated for this stock. No fishery-related mortality and serious injury has been observed since 1999; therefore, total fishery-related mortality and serious injury for this stock is unknown, but assumed to be less than 10% of the PBR and rate can be considered to be insignificant and approaching zero mortality and serious injury rate. This is not a strategic stock.

REFERENCES

- Anon. 1999. Cruise results. Summer Atlantic Ocean marine mammal survey. NOAA Ship *Oregon II* cruise 236 (99-05), 4 August - 30 September 1999. Available from: SEFSC, 3209 Frederic Street, Pascagoula, MS, 39567.
- Anon. 2002. Cruise results. Mid-Atlantic cetacean survey. NOAA Ship *Gordon Gunter* cruise GU-02-01, 6 February - 8 April 2002. Available from: SEFSC, 3209 Frederic Street, Pascagoula, MS 39567.
- Barlow, J., S. L. Swartz, T. C. Eagle and P. R. Wade. 1995. U.S. Marine mammal stock assessments: Guidelines for

preparation, background, and a summary of the 1995 assessments. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-6, 73pp.

Available from: NMFS, Southwest Fisheries Science Center, 8604 La Jolla Shores Drive, La Jolla, CA, 92037-1508.

Caldwell, D. K. and M. C. Caldwell. 1971. The pygmy killer whale, *Feresa attenuata*, in the western Atlantic, with a summary of world records. *J. Mamm.* 52:206-209.

~~Yeung~~Garrison, L.P. 2001~~3~~. ~~Estimates~~Estimated bycatch of marine mammals and ~~marine~~ turtle ~~bycatch~~ by in the U.S. Atlantic pelagic longline fleet ~~in 1999-2000~~during 2001-2002. U.S. Dep. Commer., NOAA Tech. Memo. ~~NMFS-SEFSC-467~~NMFS-SEFSC-515, ~~43~~52 pp. Available from: NMFS, Southeast Fisheries Science Center, 75 Virginia Beach ~~Dr.~~Road, Miami, ~~FL~~FL 33149.

Garrison, L.P. and P.M. Richards. 2004. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2003. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-SEFSC-527, 57 pp. Available from NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Road, Miami, FL 33149.

Hansen, L. J., K. D. Mullin and C. L. Roden. 1994. Preliminary estimates of cetacean abundance in the northern Gulf of Mexico from vessel surveys, and of selected cetacean species in the U.S. Atlantic Exclusive Economic Zone from vessel surveys. Southeast Fisheries Science Center, Miami Laboratory, Contribution No. MIA-93/94-58. Available from: NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Dr., Miami, FL, 33149.

Hansen, L. J., K. D. Mullin, T. A. Jefferson and G. P. Scott. 1996. Visual surveys aboard ships and aircraft. Pages 55-132. In: R. W. Davis and G. S. Fargion (editors), *Distribution and abundance of marine mammals in the north-central and western Gulf of Mexico: Final report. Volume II: Technical report*. OCS Study MMS 96-0027. Minerals Management Service, Gulf of Mexico OCS Region, New Orleans.

Jefferson, T. A., S. Leatherwood, and M. A. Weber. 1994. Marine mammals of the world. FAO, Rome, 320 pp.

Mullin, K. D. and G. L. Fulling. 2003. Abundance of cetaceans in the southern U.S. North Atlantic Ocean during summer 1998. *Fish. Bull.* 101:603-613.

Mullin, K. D. and G. L. Fulling. ~~In review~~2004. Abundance of cetaceans in the oceanic northern Gulf of Mexico, 1996-2001. *Mar. Mamm. Sci.* 20(4):787-807.

Mullin, K. D. and W. Hoggard. 2000. Visual surveys of cetaceans and sea turtles from aircraft and ships. Pages 111-172. In: R. W. Davis, W. E. Evans and B. Würsig (editors), *Cetaceans, sea turtles and seabirds in the northern Gulf of Mexico: Distribution, abundance and habitat associations. Volume II: Technical report*. OCS Study MMS 96-0027. Minerals Management Service, Gulf of Mexico OCS Region, New Orleans.

Wade, P.-R., and R.-P. Angliss. -1997.- Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, ~~WA~~Washington. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-12, ~~93~~93pp.

Yeung, C. 199993 pp. Available from: NOAA, National Marine Fisheries Service, 166 Water St., Woods Hole, MA 02543.

Yeung, C. 2001. Estimates of marine mammal and marine turtle bycatch by the U.S. Atlantic pelagic longline fleet in ~~1998~~1999-2000. U.S. Dep. Commer., NOAA Tech. Memo. ~~NMFS-SEFSC-430, 26~~NMFS-SEFSC-467, 43 pp. Available from: NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Dr., Miami, FL, 33149.

PYGMY SPERM WHALE (*Kogia breviceps*): Western North Atlantic Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

The pygmy sperm whale (*Kogia breviceps*) appears to be distributed worldwide in temperate to tropical waters (Caldwell and Caldwell 1989). Sightings of these animals in the ~~W~~western North Atlantic occur in oceanic waters (Mullin and Fulling 2003; SEFSC unpublished data). Pygmy sperm whales and dwarf sperm whales (*K. sima*) are difficult to ~~distinguish~~ differentiate at sea (Caldwell and Caldwell 1989, Wursig *et al.* 2000), and sightings of either species are often categorized as *Kogia spp.*. There is no information on stock differentiation for the Atlantic population. ~~In a recent study using~~ Duffield *et al.* (2003) propose using the molecular weights of myoglobin and hemoglobin, as determined by blood or muscle tissues of stranded animals, as a quick and robust way to provide species confirmation. Using hematological ~~and~~ as well as stable-isotope data, Barros *et al.* (1998) speculated that dwarf sperm whales may have a more pelagic distribution than pygmy sperm whales, and/or dive deeper during feeding bouts.

Diagnostic morphological characters have also been useful in distinguishing the two *Kogia* species (Barros and Duffield 2003), thus enabling researchers to use stranding data in distributional and ecological studies.. Specifically, the distance from the snout to the center of the blowhole in proportion to the animal's total length, as well as the height of the dorsal fin, in proportion to the animal's total length, can be used to differentiate between the two *Kogia* species when such measurements are obtainable (Barros and Duffield 2003).

POPULATION SIZE

Total numbers of pygmy sperm whales off the U.S. or Canadian Atlantic coast are unknown, although estimates from selected regions of the habitat do exist for select time periods. Because *Kogia breviceps* and *Kogia sima* are difficult to differentiate at sea, the reported abundance estimates are for both species of *Kogia*.

An abundance of 115 (CV=0.61) for *Kogia spp.* was estimated from a ~~line-transect~~ line-transect survey conducted from July 6 to September 6, 1998, by a ship and plane that surveyed 15,900 km of track line in waters north of Maryland (38° N) (Fig. 1; Palka *et al.*; in review Unpubl. Ms.). Shipboard data were analyzed using the modified direct duplicate method (Palka 1995) that accounts for school size bias and $g(0)$, the probability of detecting a group on the track line. Aerial data were not corrected for $g(0)$.

An abundance of 580 (CV=0.57) for *Kogia spp.* was estimated from a shipboard ~~line-transect~~ line-transect sighting survey conducted between 8 July and 17 August 1998 that surveyed 4,163 km of track line in waters south of Maryland (38°N) (Fig. 1; Mullin and Fulling 2003). Abundance estimates were made using the program DISTANCE (Buckland *et al.* 2001; Thomas *et al.* 1998).

The best available An abundance of 344 (CV= 0.32) for *Kogia spp.* was estimated from a line-transect sighting survey conducted during June 12 to August 4, 2004 by a ship and plane that surveyed 10,761 km of track line in waters north of Maryland (about 38° N) to the Bay of Fundy (about 45° N) (Figure 1; Palka unpubl.). Shipboard data were collected using the two independent team line-transect method and analyzed using the modified direct duplicate method (Palka 1995) accounting for biases due to school size and other potential covariates, reactive movements (Palka and Hammond 2001), and $g(0)$, the probability of detecting a group on the track line. Aerial data




Figure 1. Distribution of *Kogia spp.* sightings from NEFSC and SEFSC shipboard and aerial surveys during the summer in 1998 and 2004. Isobaths are at 100 m and 1,000 m.

were collected using the Hiby circle-back line-transect method (Hiby 1999) and analyzed accounting for g(0) and biases due to school size and other potential covariates (Figure 1; Palka unpubl.).

A survey of the U.S. Atlantic outer continental shelf and continental slope (water depths > 50m) between 27.5 – 38 °N latitude was conducted during June-August, 2004. The survey employed two independent visual teams searching with 50x bigeye binoculars. Survey effort was stratified to include increased effort along the continental shelf break and Gulf Stream front in the mid-Atlantic. The survey included 5,659 km of trackline, and there was a total of 473 cetacean sightings. Sightings were most frequent in waters north of Cape Hatteras, North Carolina along the shelf break. Data were analyzed to correct for visibility bias (g(0)) and group-size bias employing line-transect distance analysis and the direct duplicate estimator (Palka 1995; Buckland *et al.*, 2001). The resulting abundance estimate for *Kogia* spp. between Florida and Maryland was 37 (CV=0.75).

The best 2004 abundance estimate for *Kogia* spp. is the sum of the estimates from the two 1998-2004 U.S. Atlantic surveys, 695-381 (CV=0.4930), where the estimate from the northern U.S. Atlantic is 115-344 (CV=0.6132), and from the southern U.S. Atlantic is 580-37 (CV=0.5775). This joint estimate is considered the best because together these two surveys have the most complete coverage of the species' habitat.

A separate estimate of pygmy sperm whale abundance cannot be provided due to the uncertainty of species identification at sea.

Table 1. Summary of abundance estimates for the western North Atlantic *Kogia* pp. Month, year, and area covered during each abundance survey, and resulting abundance estimate (N_{best}) and coefficient of variation (CV).

Month/Year	Area	N_{best}	CV
Jul-Sep 1998	Maryland to Gulf of St. Lawrence	115	0.61
Jul-Aug 1998	Florida to Maryland	580	0.57
Jul-Sep 1998	Florida to Gulf of St. Lawrence (COMBINED)	695 ¹	0.49
Jun-Aug 2004	Maryland to Bay of Fundy	344	0.32
Jun-Aug 2004	Florida to Maryland	37	0.75
Jun-Aug 2004	Florida to Bay of Fundy (COMBINED)	381 ¹	0.30

¹ This is the combined estimate for the two survey areas

Minimum Population Estimate

The minimum population estimate is the lower limit of the two-tailed 60% confidence interval of the log-normally distributed best abundance estimate. This is equivalent to the 20th percentile of the log-normal distribution as specified by Wade and Angliss (1997). The best estimate of abundance for *Kogia* spp. is 695-381 (CV=0.4930). The minimum population estimate for *Kogia* spp. is 470-298.

Current Population Trend

The available information is insufficient to evaluate trends in population size for this species in the western North Atlantic.

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Current and maximum net productivity rates are unknown for this stock. For purposes of this assessment, the maximum net productivity rate was assumed to be 0.04. This value is based on theoretical modeling showing that cetacean populations may not grow at rates much greater than 4% given the constraints of their reproductive life history (Barlow *et al.*, 1995).

POTENTIAL BIOLOGICAL REMOVAL

Potential Biological Removal (PBR) is the product of minimum population size, one-half the maximum productivity rate, and a "recovery" factor (Wade and Angliss 1997). The minimum population size is 470-298. The maximum productivity rate is 0.04, the default value for cetaceans. The "recovery" factor, which accounts for endangered, depleted, threatened stocks, or stocks of unknown status relative to optimum sustainable population (OSP) is assumed to be 0.5 because this stock is of unknown status. PBR for the western North Atlantic *Kogia* spp. is 43.7.

ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

Fishery Information

Detailed fishery information is reported in Appendix III. There has been one logbook report of fishery-related serious injury recorded off the east coast of Florida in 2000 (Table 1) (Yeung 2001). The resultant estimated mean annual mortality is 6 pygmy sperm whales (CV=1.00) for the years 1996-2000. Fisheries Information — One observed serious injury was reported in 2000 in the pelagic longline fishery (Table 1). Table 1 in 2000 (Table 2) (Yeung 2001; Garrison 2003; Garrison and Richards, 2004). Total annual estimated average fishery-related mortality and serious injury to this stock during 1999-2003 was 6 (CV=1.0) *Kogia* spp.

Table 2. Summary of the incidental mortality and serious injury of pygmy sperm whales (*Kogia breviceps*) by commercial fishery including the years sampled (Years), the number of vessels active within the fishery (Vessels), the type of data used (Data Type), the annual observer coverage (Observer Coverage), the observed mortalities and serious injuries recorded by on-board observers, the estimated annual mortality and serious injury, the combined annual estimates of mortality and serious injury (Estimated Combined Mortality), the estimated CV of the combined estimates (Estimated CVs) and the mean of the combined estimates (CV in parentheses).

Fishery	Years	Vessels ³	Data Type	Observer Coverage	Observed Serious Injury	Observed Mortality	Estimated Serious Injury	Estimated Mortality	Estimated Combined Mortality	Estimated CVs	Mean Annual Mortality
Pelagic Longline ²	96-00 <u>99-03</u>	253, 245, <u>205,</u> 193, <u>186, 180,</u> 161, 149, <u>127</u>	Obs. Data Logbook	.03 <u>.04,</u> .04, <u>.05,</u> .03 <u>.02,</u> .04, <u>.04,</u> .04, <u>.04,</u> .01 <u>.02</u>	0, 0, 0, 1, 0, <u>0</u>	0, 0, 0, 0, <u>280</u>	- 0, <u>0, 28,</u> <u>0, 0</u>	0, 0, 0, 0, 0	0, <u>0,</u> 0, 0, <u>28,</u> <u>0, 0,</u>	0, 0, 1, <u>0, 0</u>	6 <u>(1.0)</u>
TOTAL											6 <u>(1.0)</u>

¹ Observer data (Obs. Data) are used to measure bycatch rates, and the data are collected within the Southeast Fisheries Science Center (SEFSC) Observer Program. NEFSC collects landings data (Weighout), and total landings are used as a measure of total effort for the coastal gillnet fishery. Observed bycatch rates are raised to total fishing effort reported to the SEFSC Atlantic Large Pelagic Logbook.

² The 2000 mortality estimates were taken from Table 10 in Yeung 2001 (NMFS Miami Laboratory PRD 00/01-17), and exclude the Gulf of Mexico.

³ Number of vessels in the fishery are based on vessels reporting effort to the pelagic longline logbook.

Other Mortality

From 1999-2003, 125 pygmy sperm whales were reported stranded between Maine and Puerto Rico (Table 3). The total includes 7 animals stranded in Florida in 1999; 3 animals stranded in North Carolina, 1 in South Carolina, 7 in Florida and 1 in Puerto Rico in 2000; 1 animal stranded in North Carolina, 4 in South Carolina, 3 in Georgia, and 24 in Florida in 2001; 7 animals stranded in North Carolina, 5 in South Carolina, 4 in Georgia, and 15 in Florida in 2002; and 1 animal stranded in Nova Scotia, 4 animals in North Carolina, 7 in Georgia, and 31 in Florida in 2003. In addition to the above strandings of *Kogia breviceps*, there were 8 strandings reported as *Kogia* spp. as follows: 1 *Kogia* spp. stranded in Georgia in 2000, 1 stranded in North Carolina and 2 in Florida in 2002, 1 stranded in Georgia and 3 in Florida in 2003.

Table 3. Pygmy sperm whale (*Kogia breviceps*) strandings along the Atlantic coast, 1999-2003

<u>STATE</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>TOTALS</u>
<u>Nova Scotia</u> ¹					<u>1</u>	<u>1</u>
<u>North Carolina</u>	<u>0</u>	<u>3</u>	<u>1</u> ^{2,3}	<u>7</u> ³	<u>4</u>	<u>15</u>
<u>South Carolina</u>	<u>0</u>	<u>1</u>	<u>4</u>	<u>5</u>	<u>0</u>	<u>10</u>
<u>Georgia</u>	<u>0</u>	<u>0</u> ³	<u>3</u>	<u>4</u> ³	<u>7</u> ³	<u>14</u>
<u>Florida</u>	<u>7</u> ²	<u>7</u>	<u>24</u>	<u>15</u> ⁴	<u>31</u> ⁵	<u>84</u>
<u>Puerto Rico</u>	<u>0</u>	<u>1</u> ²	<u>0</u>	<u>0</u>	<u>0</u>	<u>1</u>
<u>TOTALS</u>	<u>7</u>	<u>12</u>	<u>32</u>	<u>31</u>	<u>43</u>	<u>125</u>

¹ Data supplied by Tonya Wimmer, Nova Scotia Marine Animal Response Society (pers. comm.)

² Signs of human interaction reported

³ 1 additional *Kogia* spp. stranded

⁴ 2 additional *Kogia* spp. stranded

There were ~~no~~3 documented strandings of pygmy sperm whales along the U.S. Atlantic coast during ~~1987-2002~~1999-2003 which were classified as likely caused by fishery interactions. ~~Stranding data probably underestimate the extent of fishery-related mortality and serious injury because not all of the marine mammals which die or are seriously injured may wash ashore, nor will all of those that do wash ashore necessarily show signs of entanglement or other fishery interaction. Finally, the level of technical expertise among stranding network personnel varies widely as does the ability to recognize signs of fishery interaction.~~

Other Mortality: 1 in Florida in 1999, 1 in Puerto Rico in 2000 and 1 in North Carolina in 2001. In one of the strandings in 2002 of a pygmy sperm whale, red plastic debris was found in the stomach along with squid beaks.

Historical stranding records (1883-1988) of pygmy sperm whales in the southeastern U.S. (Credle 1988), and strandings recorded during 1988-1997 (Barros *et al.* 1998) indicate that this species accounts for about 83% of all *Kogia* spp. strandings in this area. During the period 1990-October 1998, 21 pygmy sperm whale strandings occurred in the northeastern U.S. (Delaware, New Jersey, New York and Virginia), whereas 194 strandings were documented along the U.S. Atlantic coast between North Carolina and the Florida Keys in the same period. Remains of plastic bags and other marine debris have been retrieved from the stomachs of 13 stranded pygmy sperm whales in the southeastern U.S. (Barros *et al.* 1990, 1998), and at least on one occasion the ingestion of plastic debris is believed to have been the cause of death. During the period 1987-1994, 1 animal had possible propeller cuts on its flukes. ~~In 2002, 28 pygmy sperm whales and 3 undifferentiated *Kogia* spp. stranded on the U.S. Atlantic. In one of the strandings of a pygmy sperm whale, red plastic debris was found in the stomach along with squid beaks. In the other 2002 strandings, signs of human interactions were not present or could not be assessed~~

Rehabilitation challenges for *Kogia* spp. are numerous due to limited knowledge regarding even the basic biology of these species. Advances in recent rehabilitation success has potential implications for future release and tracking of animals at sea to potentially provide information on distribution, movements and habitat use of these species (Manire *et al.*, 2004).

STATUS OF STOCK

The status of the pygmy sperm whale ~~relative~~ to OSP in the ~~western~~ U.S. Atlantic EEZ is unknown. This species is not listed as endangered or threatened under the Endangered Species Act. There is insufficient information with which to assess population trends. Total fishery-related mortality and serious injury for this stock is not less than 10% of the calculated PBR and therefore, cannot be considered to be insignificant and approaching zero mortality and serious injury rate. This is a strategic stock because the ~~1996-2000~~1999-2003 estimated average annual fishery-related mortality to pygmy sperm whales exceeds PBR.

REFERENCES

- Barlow, J., S. L. Swartz, T. C. Eagle and P. R. Wade. 1995. U.S. Marine ~~M~~mammal ~~S~~stock ~~A~~assessments: Guidelines for ~~P~~reparation, ~~B~~ackground, and a ~~S~~summary of the 1995 ~~A~~assessments. U.S. Dep. Commer., NOAA Tech. Memo. ~~NMFS-OPR-6, 73 pp.~~ NMFS-OPR-6, 73pp. Available from NMFS, Southwest Fisheries Science Center, 8604 La Jolla Shores Drive, La Jolla, CA, 92037-1508.
- Barros, N.B. and D.A. Duffield. 2003. Unraveling the mysteries of Pygmy and Dwarf sperm whales. *Strandings Newsletter of the Southeast U.S. Marine Mammal Stranding Network*. December 2003. NOAA Tech. Memo NMFS-SEFSC-521, 11 pp. Available from NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Drive, Miami, FL 33149.
- Barros, N. B., D. A. Duffield, P. H. Ostrom, D. K. Odell and V. R. Cornish. 1998. Nearshore vs. offshore ecotype differentiation of *Kogia breviceps* and *K. simus* based on hemoglobin, morphometric and dietary analyses. Abstracts. World Marine Mammal Science Conference Abstracts. Monaco. 20-24 January.
- Barros, N. B., D. K. Odell and G. W. Patton. 1990. Ingestion of plastic debris by stranded marine mammals from Florida. Page 746. -In: R. S. Shomura and M.L. Godfrey (editors eds.), Proceedings of the Second International Conference on Marine Debris. NOAA Tech. Memo. NOAA-TM-NMFS-SWFSC-154. Available from NMFS, Southwest Fisheries Science Center, 8604 La Jolla Shores Drive, La Jolla, CA, 92037-1508.
- Buckland, S. T., D. R. Anderson, K. P. Burnham, J. L. Laake, D. L. Borchers and L. Thomas. 2001. Introduction to distance sampling: estimating abundance of biological populations. Oxford University Press, New York, 432 pp.
- Caldwell, D. K. and M. C. Caldwell. 1989. Pygmy sperm whale *Kogia breviceps* (de Blainville 1838): dwarf sperm whale *Kogia simus* Owen, 1866. Pages pp. 235-260. -In: S. H. Ridgway and R. Harrison (editors eds.), Handbook of marine mammals, Vol. 4: ~~R~~iver dolphins and the larger toothed whales. -Academic Press, San Diego. pp.
- Credle, V. R. 1988. Magnetite and magnetoreception in dwarf and pygmy sperm whales, *Kogia simus* and *Kogia breviceps*. MSc. Thesis. University of Miami. Coral Gables, FL.
- Mullin, K. D. and G. L. Fulling. 2003. Abundance and distribution of cetaceans in the southern U.S. North Atlantic Ocean during summer 1998. *Fish. Bull.*, U.S. 101:603-613.
- Palka, D. 1995. Abundance estimate of the Gulf of Maine harbor porpoise. *Rep. int Whal. Commn. Special Issue*

- Palka, D., G. Waring and D. Potter. In review. Abundances of cetaceans and sea turtles in the northwest Atlantic during summer 1995 and 1998. *Fish. Bull., U.S.*
- Thomas, L., J. L. Laake, J. F. Derry, S. T. Buckland, D. L. Borchers, D. R. Anderson, K. P. Burnham, S. Strindberg, S. L. Hedley, F. F. C. Marques, J. H. Pollard and R. M. Fewster. 1998. Distance 3.5. Research Unit for Wildlife Population Assessment, University of St. Andrews, St. Andrews, UK.
- Wade P. R. and R. P. Angliss. 1997. Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-12, 93 pp.
- Duffield, D.A., N.B. Barros, E.O. Espinoza, S. Ploen, F.M.D. Gulland, and J.E. Heyning. 2003. Identifying Pygmy and Dwarf Sperm Whales (Genus *Kogia*) using electrospray ionization mass spectrometry of myoglobin and hemoglobin. *Mar. Mammal Sci.* 19(2):395-399.
- Garrison, L.P. 2003. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2001-2002. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-SEFSC-515, 52 pp. Available from NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Road, Miami, FL 33149.
- Garrison, L.P. and P.M. Richards. 2004. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2003. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-SEFSC-527, 57 pp. Available from NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Road, Miami, FL.
- Wursig, B., T.A. Jefferson, and D.J. Schmidly. 2000. *The marine mammals of the Gulf of Mexico*. Texas A&M University Press, College Station, TX, 256 pp.
- Mullin, K. D. and G. L. Fulling. 2003. Abundance and distribution of cetaceans in the southern U.S. North Atlantic Ocean during summer 1998. *Fish. Bull., U.S.* 101:603-613.
- Hiby, L. 1999. The objective identification of duplicate sightings in aerial survey for porpoise. pp. 179-189. In: G.W. Garner, S. C. Amstrup, J. L. Laake, B. F. J. Manly, L. L. McDonald, and D. G. Robertson (eds.) *Marine mammal survey and assessment methods*. Balkema, Rotterdam. 287 pp.
- Manire, C.A., H.L. Rhinehart, N.B. Barros, L. Byrd, and P. Cunningham-Smith. 2004. An approach to the rehabilitation of *Kogia* spp. *Aquatic Mamm.* 30(2):257-270.
- Palka, D. 1995. Abundance estimate of the Gulf of Maine harbor porpoise. pp. 27-50 In: A. Bjørge and G.P. Donovan (eds.) *Biology of the Phocoenids*. Rep. Int. Whal. Commn (Special Issue 16) I-x + 552 pp.
- Palka, D. and P.S. Hammond. 2001. Accounting for responsive movement in line-transect estimates of abundance. *Can. J. Fish. Aquat. Sci.* 58: 777-787.
- Thomas, L., J. L. Laake, J. F. Derry, S. T. Buckland, D. L. Borchers, D. R. Anderson, K. P. Burnham, S. Strindberg, S. L. Hedley, F. F. C. Marques, J. H. Pollard and R. M. Fewster. 1998. Distance 3.5. Research Unit for Wildlife Population Assessment, University of St. Andrews, St. Andrews, UK.
- Wade, P.R., and R.P. Angliss. 1997. Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-12, 93 pp. Available from: NOAA, National Marine Fisheries Service, 166 Water St., Woods Hole, MA 02543.
- Yeung, C. 2001. Estimates of marine mammal and marine turtle bycatch by the U.S. Atlantic pelagic longline fleet in 1999-2000. NOAA Tech. Memo. NOAA-TM-SEFSC-467, 42 pp. Available from: NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Dr., Miami, FL, 33149.

SHORT-FINNED PILOT WHALE (*Globicephala macrorhynchus*): Western North Atlantic Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

There are two species of pilot whales in the ~~W~~western ~~North~~ Atlantic: the Atlantic or long-finned pilot whale, (*Globicephala melas*;) and the short-finned pilot whale, (*G. macrorhynchus*). These species are difficult to ~~identify~~~~differentiate~~ to the species level at sea; therefore, some of the descriptive material below refers to *Globicephala* spp. and is identified as such. The species boundary is considered to be in the New Jersey to Cape Hatteras area. Sightings north of this area are likely *G. melas*. The short-finned pilot whale is distributed worldwide in tropical to warm temperate waters (Leatherwood and Reeves 1983). The northern extent of the range of this species within the ~~USA~~~~U.S.~~ Atlantic Exclusive Economic Zone (EEZ) is generally thought to be Cape Hatteras, North Carolina (Leatherwood and Reeves 1983). Sightings of these animals in ~~US~~~~U.S.~~ Atlantic EEZ occur ~~primarily within the Gulf Stream [Southeast Fisheries Science Center (SEFSC) unpublished data]~~~~in oceanic waters (Mullin and Fulling 2003)~~ and along the continental shelf and continental slope in the northern Gulf of Mexico (~~Mullin Hansen et al. 1991~~~~6; SEFSC unpublished data~~~~Mullin and Hoggard 2000; Mullin and Fulling 2003~~). There is no information on stock differentiation for the Atlantic population.

POPULATION SIZE

The total number of short-finned pilot whales off the eastern ~~USA~~~~U.S.~~ and Canadian Atlantic coast is unknown, although ~~ten~~ estimates from selected regions of the habitat do exist for select time periods. Sightings were almost exclusively in the continental shelf edge and continental slope areas (Figure Fig. 1). Two estimates were derived from catch data and population models that estimated the abundance of the entire stock. ~~Seven~~ ~~seasonal~~ ~~Seasonal~~ estimates are available from selected regions in ~~USA~~~~U.S.~~ waters during spring, summer and autumn 1978-82, August 1990, June-July 1991, August-September 1991, June-July 1993, -July-September 1995, and July-August 1998, ~~and June-August 2004~~. Because long-finned and short-finned pilot whales are difficult to identify at sea, seasonal abundance estimates were reported for *Globicephala* spp., both long-finned and short-finned pilot whales. One estimate is available from the Gulf of St. Lawrence.

Mitchell (1974) used cumulative catch data from the 1951-61 drive fishery off Newfoundland to estimate the initial population size (ca. 50,000 animals).

Mercer (1975) used population models to estimate a population in the same region of between 43,000-96,000 long-finned pilot whales, with a range of 50,000-60,000 being considered the best estimate.

An abundance of 11,120 (CV=0.29)

Globicephala spp. was estimated from an aerial survey program conducted from 1978 to 1982 on the continental shelf and shelf edge waters between Cape Hatteras, North Carolina, and Nova Scotia (CETAP 1982). An abundance of 3,636 (CV=0.36) *Globicephala* spp. was estimated from a June and July 1991 shipboard line-transect sighting survey conducted primarily between the 200 and 2,000 m isobaths from Cape Hatteras to Georges Bank (Waring et al. 1992; Waring 1998). An abundance of 3,368 (CV=0.28) and 5,377 (CV=0.53) *Globicephala* spp. was estimated from line-transect aerial surveys conducted from August to September 1991 using the Twin Otter and AT-11, respectively (Anon. 1991). As recommended in the GAMMS Workshop Report (Wade and Angliss 1997), estimates older than eight years are deemed unreliable, and therefore should not be used for PBR determinations. Further, due to changes in survey methodology, these data should not be used to make comparisons to more current estimates.

An abundance of 668 (CV=0.55) *Globicephala* spp. was estimated from a June and July 1993 shipboard




Figure 1. Distribution of pilot whale sightings from NEFSC and SEFSC vessel and aerial summer surveys during 1998 and 2004. Isobaths are at 100 and 1,000 m.

line-transect sighting survey conducted principally between the 200m and 2,000 m isobaths from the southern edge of Georges Bank, across the Northeast Channel to the southeastern edge of the Scotian Shelf (Table 1; Anon. 1993b). Data were collected by two alternating teams that searched with 25x150 binoculars and were analyzed using DISTANCE (Buckland *et al.* 1993, 2001; Laake Thomas *et al.* 1993, 8). Estimates include school-size bias, if applicable, but do not include corrections for $g(0)$ or dive-time. Variability was estimated using bootstrap resampling techniques.

An abundance of 8,176 (CV=0.65) *Globicephala* spp. was estimated from a July to September 1995 sighting survey conducted by two ships and an airplane that covered waters from Virginia to the mouth of the Gulf of St. Lawrence (Table 1; Palka *et al.*, in review Unpubl. Ms.). Total track line length was 32,600 km. The ships covered waters between the 50 and 1000 fathom depth contour lines, the northern edge of the Gulf Stream, and the northern Gulf of Maine/Bay of Fundy region. The airplane covered waters in the mid-Atlantic from the coastline to the 50 fathom depth contour line, the southern Gulf of Maine, and shelf waters off Nova Scotia from the coastline to the 1000 fathom depth contour line. Data collection and analysis methods used were described in Palka (1996).

Kingsley and Reeves (1998) obtained an abundance estimate of 1,600 long-finned pilot whales (CV=0.65) from a late August and early September aerial survey of cetaceans in the Gulf of St. Lawrence in 1995 and 1998 (Table 1). Based on an examination of long-finned pilot whale summer distribution patterns and information on stock structure, it was deemed appropriate to combine these estimates with NMFS 1995 summer survey data. The best 1995 abundance estimate for *Globicephala* spp., 9,776 (CV=0.55), is was the sum of the estimates from the USA U.S. and Canadian surveys, where the estimate from the USA U.S. survey is was 8,176 (CV=0.65) and from the Canadian, 1,600 (CV=0.65).

An abundance of 9,800 (CV=0.34) *Globicephala* spp. was estimated from a line-transect sighting survey conducted during July 6 to September 6, 1998, by a ship and plane that surveyed 15,900 km of track line in waters north of Maryland (38° N) (Figure Fig. 1; Palka *et al.*, in review Unpubl. Ms.). Shipboard data were analyzed using the modified direct duplicate method (Palka 1995) that accounts for school size bias and $g(0)$, the probability of detecting a group on the track line. Aerial data were not corrected for $g(0)$.

An abundance of 4,724 (CV=0.61) *Globicephala* spp. was estimated from a shipboard line-transect sighting survey conducted between 8 July and 17 August 1998 that surveyed 54,570 km of track line in waters south of Maryland (38° N) (Figure Fig. 1; Mullin in review and Fulling 2003). Abundance estimates were made using the program DISTANCE. This estimate is a recalculation of the same data reported in previous SARs. For more details see Mullin and Fulling (2003).

An abundance of 15,436 (CV= 0.33) for *Globicephala* spp. was estimated from a line transect sighting survey conducted during June 12 to August 4, 2004 by a ship and plane that surveyed 10,761 km of track line in waters north of Maryland (about 38° N) to the Bay of Fundy (about 45° N) (Figure 1; Palka unpubl.). Shipboard data were collected using the two independent team line transect method and analyzed using the modified direct duplicate method (Palka 1995) accounting for biases due to school size and other potential covariates, reactive movements (Palka and Hammond 2001), and $g(0)$, the probability of detecting a group on the track line. Aerial data were collected using the Hiby circle-back line transect method (Hiby 1999) and analyzed accounting for $g(0)$ and biases due to school size and other potential covariates (Figure 1; Palka unpubl.).

A shipboard survey of the U.S. Atlantic outer continental shelf and continental slope (water depths >50m) between Florida and Maryland (27.5 and 38° N latitude) was conducted during June-August, 2004. The survey employed two independent visual teams searching with 50x bigeye binoculars. Survey effort was stratified to include increased effort along the continental shelf break and Gulf Stream front in the mid-Atlantic. The survey included 5,659 km of trackline, and there were a total of 473 cetacean sightings. Sightings were most frequent in waters north of Cape Hatteras, North Carolina along the shelf break. Data were analyzed to correct for visibility bias ($g(0)$) and group-size bias employing line transect distance analysis and the direct duplicate estimator (Palka 1995; Buckland *et al.* 1993; Laake *et al.* 1993) where school size bias and ship attraction were accounted for.

The best available, 2001). The resulting abundance estimate for *Globicephala* spp., 14,524 (CV=0.30), spp. between Florida and Maryland was 15,411 (CV=0.43).

The best 2004 abundance estimate for *Globicephala* spp. is the sum of the estimates from the two 1998 USA 2004 U.S. Atlantic surveys, 30,847 (CV=0.27), where the estimate from the northern USA U.S. Atlantic is 915,800 (CV=0.34), and from the southern USA U.S. Atlantic is 4,724 (CV=0.61). This joint estimate is considered the best because together these two surveys have the most complete coverage of the species' habitat.

Table 1. Summary of abundance estimates for the western North Atlantic *Globicephala* spp. Month, year, and area covered during each abundance survey, and resulting abundance estimate (N_{best}) and coefficient of variation (CV).

Month/Year	Area	N_{best}	CV
Jun-Jul 1993 Jul-Sep 1995	Virginia Maryland to Gulf of St. Lawrence	9,776 559,800	0.34

Jul-Sep 1998 <u>Jul-Aug 1998</u>	Maryland to Gulf of St. Lawrence <u>Florida to Maryland</u>	9,800.34 <u>14,524¹</u>	0.61 <u>0.26</u>
Jul-Sep 1998	<u>Florida to Gulf of St. Lawrence to Florida (COMBINED)</u>	<u>14,524¹</u>	<u>0.26</u>
<u>Jun-Aug 2004</u>	<u>Maryland to Bay of Fundy</u>	<u>15,436</u>	<u>0.30</u>
<u>Jun-Aug 2004</u>	<u>Florida to Maryland</u>	<u>15,411</u>	<u>0.43</u>
<u>Jun-Aug 2004</u>	<u>Florida to Bay of Fundy (COMBINED)</u>	<u>30,847¹</u>	<u>0.27</u>

¹ This is the combined estimate for the two survey areas

Minimum Population Estimate

The minimum population estimate is the lower limit of the two-tailed 60% confidence interval of the log-normally distributed best abundance estimate. This is equivalent to the 20th percentile of the log-normal distribution as specified by Wade and Angliss (1997). The best estimate of abundance for *Globicephala* sp. is 14,524 30,847 (CV=0.30²⁷). The minimum population estimate for *Globicephala* spp. is 14,524 30,847.

Current Population Trend

There are insufficient data to determine the population trends for this species *Globicephala* spp.

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Current and maximum net productivity rates are unknown for this stock. For purposes of this assessment, the maximum net productivity rate was assumed to be 0.04. This value is based on theoretical modeling showing that cetacean populations may not grow at rates much greater than 4% given the constraints of their reproductive life history (Barlow *et al.*, 1995).

POTENTIAL BIOLOGICAL REMOVAL

Potential Biological Removal (PBR) is the product of minimum population size, one-half the maximum productivity rate, and a "recovery" factor (MMPA Sec. 3, 16 U.S.C. 1362; Wade and Angliss 1997). The minimum population size for *Globicephala* spp. is 14,524 30,847 (CV=0.30²⁷). The maximum productivity rate is 0.04, the default value for cetaceans. The "recovery" factor, which accounts for endangered, depleted, threatened stocks, or stocks of unknown status relative to optimum sustainable population (OSP) is assumed to be 0.48 because the CV of the average mortality estimate is between 0.3-0.6 (Wade and Angliss 1997), and because this stock is of unknown status. PBR for the western North Atlantic *Globicephala* spp. is 108 237.

ANNUAL HUMAN-CAUSED MORTALITY

Fishery Information

Detailed fishery information are reported in Appendix III. Total fishery-related mortality and serious injury cannot be estimated separately for the two species of pilot whales in the US U.S. Atlantic EEZ because of the uncertainty in species identification by fishery observers. The Atlantic Scientific Review Group advised adopting the risk-averse strategy of assuming that either species might have been subject to the observed fishery-related mortality and serious injury. Total annual estimated average fishery-related mortality or serious injury of this stock during 1996-2000 1999-2003 in the USA U.S. fisheries listed below was 193 201 pilot whales (CV=0.43⁰) (Table 2). The Canadian average annual mortality estimate for 1996 from the Nova Scotia trawl fisheries is 6

The distribution of long-finned pilot whales. It is not possible to estimate variance of the Canadian estimate. The total average annual mortality estimate for 1996-2000 from the USA and Nova Scotia trawl fisheries is 199 (CV = 0.43) (Table 2).

Fishery Information

USA

The level of past or current, direct, human-caused mortality of whale, a northern species, overlaps with that of the short-finned pilot whales in the US Atlantic EEZ is unknown. The short-finned whale, a predominantly southern species, between 35°30'N to 38°00'N (Leatherwood *et al.* 1976). Although long-finned pilot whale has been taken in the pelagic longline fishery in Atlantic waters off the southeastern USA (Lee *et al.* 1994; SEFSC unpublished data), whales are most likely taken in the waters north of Delaware Bay, many of the pilot whale takes are not identified to species and bycatch does occur in the overlap area. In this summary, therefore, fishing interactions are considered for undifferentiated pilot whales (*Globicephala* spp.).

Earlier Interactions

Prior to 1977, there was no documentation of marine mammal bycatch in distant-water fleet (DWF) activities off the northeast coast of the USA. A fishery observer program, which has collected fishery data and information on incidental bycatch of marine mammals, was established in 1977 with the implementation of the Magnuson Fisheries Conservation and Management Act (MFCMA). DWF effort in the US Atlantic EEZ under MFCMA has been directed primarily towards Atlantic mackerel and squid. An average of 120 different foreign vessels per year (range 102-161) operated within the US Atlantic EEZ during 1977 through 1982. In 1982, there were 112 different foreign vessels; 18 (16%) were Japanese tuna longline vessels operating along the USA Atlantic coast. This was the first year that the Northeast Regional Observer Program assumed responsibility for observer coverage of the longline vessels. The number of foreign vessels operating within the US Atlantic EEZ each year between 1983 and 1991 averaged 33 and ranged from 9 to 67. The number of Japanese longline vessels included among the DWF vessels averaged 6 and ranged from 3 to 8 between 1983 and 1988. MFCMA observer coverage on DWF vessels was 25-35% during 1977-82, increased to 58%, 86%, 95%, and 98%, respectively, during 1983-86, and 100% observer coverage was maintained from 1987-91. Foreign fishing operations for squid ceased at the end of the 1986 fishing season, and for mackerel, at the end of the 1991 fishing season.

During 1977-1991, observers in this program recorded 436 pilot whale mortalities in foreign-fishing activities (Waring *et al.* 1990; Waring 1995). A total of 391 (90%) were taken in the mackerel fishery, and 41 (9%) occurred during *Loligo* and *Illex* squid-fishing operations. This total includes 48 documented takes by USA vessels involved in joint venture fishing operations in which USA captains transfer their catches to foreign processing vessels. Due to temporal fishing restrictions, the bycatch occurred during winter/spring (December to May) in continental shelf and continental shelf edge waters (Fairfield *et al.* 1993; Waring 1995); however, the majority of the takes occurred in late spring along the 100 m isobath. Two animals were also caught in both the hake fishery and tuna longline fisheries (Waring *et al.* 1990).

The distribution of long-finned pilot whale, a northern species, overlaps with that of the short-finned pilot whale, a predominantly southern species, between 35°30'N to 38°00'N (Leatherwood et al. 1976). Although long-finned pilot whales are most likely taken in the waters north of Delaware Bay, many of the pilot whale takes are not identified to species and bycatch does occur in the overlap area. In this summary, therefore, long-finned pilot whales (Globicephala melas) and unidentified pilot whales (Globicephala sp.) are considered together.

Data on current incidental takes in USA fisheries are available from several sources. In 1986, NMFS established a mandatory self-reported fisheries information system for large pelagic fisheries. Data files are maintained at the Southeast Fisheries Science Center (SEFSC). The Northeast Fisheries Science Center (NEFSC) Fisheries Observer Program was initiated in 1989, and since that year several fisheries have been covered by the program. In late 1992 and in 1993, the SEFSC provided observer coverage of pelagic longline vessels fishing off the Grand Banks (Tail of the Banks) and provides observer coverage of vessels fishing south of Cape Hatteras.

1990). Bycatch has been observed by NMFS Sea Samplers in the pelagic drift gillnet, pelagic longline, and pelagic pair trawl, bluefin tuna purse seine, North Atlantic bottom trawl, Atlantic squid, mackerel, butterfish trawl, and Mid-Atlantic coastal gillnet fisheries, but no mortalities or serious injuries have been documented in the Northeast multispecies sink gillnet fishery.

Pelagic Drift Gillnet

The estimated total number of hauls in the pelagic drift gillnet fishery increased from 714 in 1989 to 1,144 in 1990; thereafter, with the introduction of quotas, effort was severely reduced. The estimated number of hauls in 1991, 1992, 1993, 1994, 1995, 1996, and 1998 were 233, 243, 232, 197, 164, 149, and 113 respectively. In 1996 and 1997, NMFS issued management regulations which prohibited the operation of this fishery in 1997. Further, in January 1999, NMFS issued a Final Rule to prohibit the use of driftnets (*i.e.*, permanent closure) in the North Atlantic swordfish fishery (50 CFR Part 630). Fifty-nine different vessels participated in this fishery at one time or another between 1989 and 1993. From 1994 to 1998, between 10 and 13 vessels participated in the fishery. Observer coverage, expressed as percent of sets observed, was 8% in 1989, 6% in 1990, 20% in 1991, 40% in 1992, 42% in 1993, 87% in 1994, 99% in 1995, 64% in 1996, no fishery in 1997, and 99% in 1998. Effort was concentrated along the southern edge of Georges Bank and off Cape Hatteras. Examination of the species composition of the catch and locations of the fishery throughout the year, suggested that the pelagic drift gillnet fishery be stratified into two strata, a southern or winter stratum, and a northern or summer stratum. Estimates of the total bycatch, from 1989 to 1993, were obtained using the aggregated (pooled 1989-1993) catch rates, by stratum (Northridge 1996). Estimates of total annual bycatch for 1994 and 1995 were estimated from the sum of the observed caught and the product of the average bycatch per haul and the number of unobserved hauls as recorded in self-reported fisheries information. Variances were estimated using bootstrap re-sampling techniques. Between 1989 and 1998, 87 mortalities were observed in the large pelagic drift gillnet fishery. The annual fishery-related mortality (CV in parentheses) was 77 in 1989 (0.24), 132 in 1990 (0.24), 30 in 1991 (0.26), 33 in 1992 (0.16), 31 in 1993 (0.19), 20 in 1994 (0.06), 9.1 in 1995 (0), 11 in 1996 (0.17), no fishery in 1997, and 12 in 1998 (0). Since this fishery no longer exists it has been excluded from Table 2. Pilot whales were taken along the continental shelf edge, northeast of Cape Hatteras in January and February. Takes were recorded at the continental shelf edge east of Cape Charles, Virginia, in June. Pilot whales were taken from Hydrographer Canyon along the Great South Channel to Georges Bank from July to November. Takes occurred at the Oceanographer Canyon continental shelf break and along the continental shelf northeast of Cape Hatteras in October-November.

Pelagic Pair Trawl

Effort in the pelagic pair trawl fishery has increased during the period 1989 to 1993, from zero hauls in 1989 and 1990, to an estimated 171 hauls in 1991, and then to an estimated 536 hauls in 1992, 586 in 1993, 407 in 1994, and 440 in operated as an experimental fishery from 1991 to 1995. This fishery ceased operations in 1996; when NMFS rejected a petition to consider pair trawl gear as an authorized gear type in the Atlantic tunas fishery. The fishery operated from August-November in 1991, from June-November in 1992, from June-October in 1993, and from mid-summer to November in 1994 and 1995. Fisheries Observer began in October 1992 (Gerrior *et al.* 1994), and 48 sets (9% of the total) were sampled in that season, 102 hauls (17% of the total) were sampled in 1993. In 1994 and 1995, 52% and 54%, respectively, of the sets were observed. Twelve vessels have operated in this fishery. The fishery extends from 35°N to 41°N, and from 69°W to 72°W. Approximately 50% of the total effort was within a one degree square at 39°N, 72°W, around Hudson Canyon. Examination of the 1991-1993 locations and species composition of the bycatch, showed little seasonal change for the six months of operation and did not warrant any seasonal or areal stratification of this fishery (Northridge 1996). Five pilot whale (*Globicephala* sp.) mortalities were reported in the self-reported fisheries information in 1993. In 1994 and 1995 observers reported 1 and 12 mortalities, respectively. The estimated fishery-related mortality to pilot whales in the western U.S. Atlantic attributable to this fishery in 1994 was 2.0 (CV=0.49) and 22 (CV=0.33) in 1995. Since this fishery- no longer exists, it has been excluded from Table 2.

During the 1994 and 1995 experimental fishing seasons, fishing gear experiments were conducted to collect data on environmental parameters, gear behavior, and gear handling practices to evaluate factors affecting catch and bycatch (Goudcy 1995, 1996). Results of these studies were inconclusive in identifying factors responsible for marine mammal bycatch.

Pelagic Longline

Total effort, excluding the Gulf of Mexico, for the pelagic longline fishery, based on mandatory self-reported fisheries information, was 11,279 sets in 1991, 10,311 sets in 1992, 10,444 sets in 1993, 11,082 sets in 1994, 11,493 sets in 1995, 9,864 sets in 1996, 9,499 sets in 1997, 7,589 sets in 1998, 6,770 sets in 1999, and 6,582 sets in 2000 from 1991 to 2000. (Cramer 1994; Scott and Brown 1997; Johnson *et al.* 1999; Yeung 1999a; Yeung *et al.* 2000; Yeung,). In the 2001 Stock Assessment Report, the annual effort has been recalculated to include those sets targeting other species in conjunction with tuna/swordfish, instead of just effort that exclusively targeted tuna/swordfish as in previous reports (Johnson *et al.* 1999; Yeung 1999a). The result is an average increase in self-reported effort of roughly 10% (Yeung *et al.* 2000). The fishery has been observed from January to March off Cape Hatteras, in May and June in the entire mid-Atlantic, and in July through December in the mid-Atlantic Bight and off Nova Scotia. This fishery has been monitored with 3-6% observer coverage, in terms of sets observed, since 1992. The 1993-1997 estimated take was based on a revised analysis of the observed incidental take and self-reported incidental take and effort data, and replaces previous estimates for the 1990-1993 and 1994-1995 periods (Cramer 1994; Scott and Brown 1997; Johnson *et al.* 1999). Further, Yeung (1999b) revised the 1992-1997 fishery mortality estimates in Johnson *et al.* (1999) to include seriously injured animals. The 1998, 1999 and 2000 bycatch estimates were from Yeung (1999a), Yeung *et al.* (2000), and Yeung (2001), respectively.

Most of the estimated marine mammal bycatch was from USU.S. Atlantic EEZ waters between South Carolina and Cape Cod (Johnson *et al.* 1999). Pilot whales are frequently observed feeding to feed on hooked fish, particularly big-eye tuna (NMFS unpublished data). Between 1992-2000, 70 pilot whales (including 2 identified as a short-finned pilot whales) were released alive, including 36 that were considered seriously injured (of which 1 was identified as a short-finned pilot whale), and 45 mortalities were observed. January-March bycatch was concentrated on the continental shelf edge northeast of Cape Hatteras. Bycatch was recorded in this area during April-June, and takes also occurred north of Hydrographer Canyon off the continental shelf in water over 1,000 fathoms during April-June. During the July-September period, takes occurred on the continental shelf edge east of Cape Charles, Virginia, and on Block Canyon slope in over 1,000 fathoms of water. October-December bycatch occurred along the 20 to 50 fathom contour lines between Barnegatt Bay and Cape Hatteras.

The estimated fishery-related mortality to pilot whales in the USU.S. Atlantic (excluding the Gulf of Mexico) attributable to this fishery was: 127 in 1992 (CV=1.00) and 93 in 1999 (CV=1.00) 24 in 2000 (CV=1.0), 20 in 2001 (CV=0.7), 2 (CV=0.54) in 2002, and 0 (zero) in 2003. The estimated serious injuries were 40 (CV=0.71) in 1992, 19 (CV=1.00) in 1993, 232 (CV=0.53) in 1994, 345 (CV=0.51) in 1995, 0 from 1996 to 1998, 288 (CV=0.79) in 1999, and 109 (CV=0.88) in 2000. This (includes 37 estimated short-finned pilot whales in 1995 (CV=1.00); 50 (CV=0.48) in 2001, and 52 (CV=0.49) in 2002, and 21 (CV=NA) in 2003. The average 'combined' annual mortality between 1996 and 2000 was 103 pilot whales (CV=0.63) (Table 2). Seriously injured in 1999-2003 was 132 pilot whales (CV=0.49) (Table 2). Animals released alive but judged to have been seriously injured and released alive animals are combined with mortalities in the category 'combined mortality'. No pilot whale takes have been observed in the Gulf of Mexico fishery. The majority of pilot whale bycatch was concentrated along the continental shelf break between New Jersey and Cape Hatteras, occurring primarily in the late summer to early fall (Yeung 2001; Garrison, 2003; Garrison and Richards, 2004).

Bluefin Tuna Purse Seine

The tuna purse seine fishery between Cape Hatteras and Cape Cod is directed at small and medium bluefin and skip jack/skipjack for the canning industry, while north of Cape Cod, purse seine vessels are directed at large

medium and giant bluefin tuna (NMFS 1995). The latter fishery is entirely separate from any other Atlantic tuna purse seine fishery. Spotter aircraft are used to locate fish schools. The official start date is August 15, set by regulation. Individual vessel quotas (IVQs) and a limited access system prevent a derby fishery situation. Catch rates are high with this gear and consequently, the season usually only lasts a few weeks for large mediums and giants. The 1996 regulations allocated 250 MT (5 IVQs) with a minimum of 90% giants and 10% large mediums. Limited observer data are available for the bluefin tuna purse seine fishery. Out of 45 total trips made in 1996, 43 trips (95.6%) were observed. Forty-four sets were made on the 43 observed trips and all sets were observed. A total of 136 days were covered. Two interactions with pilot whales were observed in 1996. In one interaction, the net was actually pursued around one pilot whale, the rings were released and the animal escaped alive, condition unknown. This set occurred east of the Great South Channel and just north of the Cultivator Shoals region on Georges Bank. In a second interaction, 5 pilot whales were encircled in a set. The net was opened prior to pursuing to let the whales swim free, apparently uninjured. This set occurred on the Cultivator Shoals region on Georges Bank. Since 1996, this fishery has not been observed.

North Atlantic Bottom Trawl

Vessels in the North Atlantic bottom trawl fishery;

Southern New England/Mid-Atlantic Squid, Mackerel, Butterfish Trawl Fisheries

Because of spatial and temporal differences in the harvesting of *Illex* and *Loligo* squid, and Atlantic mackerel, each one of these sub-fisheries are described separately. The *Illex* and *Loligo* squid fisheries are managed by moratorium permits, gear and area restrictions, quotas, and trip limits. The Atlantic mackerel and butterfish fisheries are managed by an annual quota system.

Historically, the mid-Atlantic mackerel and squid trawl fisheries were combined into the Atlantic mid-water trawl fishery in the revised proposed list of fisheries in 1995. The mackerel trawl fishery was classified as a Category II fishery since 1990 and the squid fishery was originally classified as a Category II fishery in 1990, but was reclassified as a Category III fishery under the MMPA, were observed in order to meet fishery management needs, rather than marine mammal management needs. An average of 970 (CV=0.04) vessels (full and part time) participated annually in the fishery during 1989-1993. The in 1992. The combined fishery was then reclassified as a Category II fishery in 1995.

Illex Squid

The U.S. domestic fishery, ranging from Southern New England to Cape Hatteras North Carolina, reflects patterns in the seasonal distribution of *Illex* squid (*Illex illecebrosus*). *Illex* are harvested offshore mainly by small mesh otter trawlers when they are distributed in continental shelf and slope waters during the summer months (June-September) (Clark ed. 1998). Since 1996, 45% of all pilot whale takes observed were caught incidental to *Illex* squid fishing operations; 1 in 1996, 1 in 1998 and 2 in 2000. Annual observer coverage of this fishery has varied widely and reflects only the months when the fishery is active in New England in all seasons. One mortality was documented in 1990, and one animal was released alive and uninjured in 1993. The estimated fishery-related mortality to of pilot whales in the USA Atlantic attributable to this fishery was: 0 in 1994-1998, 228 in 1999, and 0 in 2000 45 in 1996 (CV=1.27), 0 in 1997, 85 in 1998 (CV=0.65), 0 in 1999, 34 in 2000 (CV=0.65) and 0 in 2001. The average annual mortality between 1996-2000 1998 and 2002 was 4630 pilot whales (CV=1.03 CV=0.50) (Table 2).

Loligo Squid

The U.S. domestic fishery for *Loligo* squid (*Loligo pealeii*) occurs mainly in Southern New England and mid-Atlantic waters. Fishery patterns reflect *Loligo* seasonal distribution where most effort is directed offshore near the edge of the continental shelf during the fall and winter months (October-March), and inshore during the spring and summer months (April-September) (Clark ed. 1998). This fishery is dominated by small-mesh otter trawlers, but substantial landings are also taken by inshore pound nets and fish traps during the spring and summer months (Clark ed. 1998). Only one pilot whale incidental take has been observed in *Loligo* squid fishing operations since 1996. The one take was observed in 1999 in the offshore fishery. No pilot whale takes have been observed in the inshore fishery. The estimated fishery-related mortality of pilot whales attributable to the fall/winter offshore fishery was 0 between 1996 and 1998, 49 in 1999 (CV=0.97) and 0 between 2000 and 2003. The average annual mortality between 1999 and 2003 was 10 pilot whales (CV=0.97) (Table 2). However, these estimates should be viewed with caution due to the extremely low (<1%) observer coverage.

Atlantic Squid, Mackerel, Butterfish Trawl

The mid-Atlantic

The U.S. domestic fishery for Atlantic mackerel and squid trawl fisheries were combined into the Atlantic mid-water trawl fishery in the revised proposed list of fisheries in 1995. The fishery occurs along the USA mid-Atlantic continental shelf region between New Brunswick, Canada, and Cape Hatteras year around. The (*Scomber scombrus*) occurs primarily in the Southern New England and mid-Atlantic waters between the months of January and May (Clark ed. 1998). This fishery is dominated by mid-water (pelagic) trawls. No incidental takes of pilot whales have been observed in the domestic mackerel fishery.

A U.S. joint venture (JV) fishery was conducted in the mid-Atlantic region from February to May 1998. NMFS maintained 100% observer coverage of the foreign joint venture vessels where 152 transfers from the U.S. vessels were observed. No incidental takes of pilot whales have been observed in the mackerel fishery. The former

distant water fleet fishery has been non-existent since 1977. There is also a mackerel trawl fishery was classified as a Category II fishery since 1990 and the squid fishery was originally classified as a Category II fishery in 1990, but was reclassified as a Category III fishery in 1992. The combined fishery was then reclassified as a Category II fishery in 1995. In 1996, mackerel, squid, and butterfish trawl fisheries were combined into the Atlantic squid, mackerel, butterfish trawl fishery, and maintained a Category II classification. Three fishery-related mortalities in the Gulf of Maine that generally occurs during the summer and fall months (May-December) (Clark ed. 1998). There have been no observed incidental takes of pilot whales were reported in self-reported fisheries information from the mackerel trawl fishery between 1990-1992. Six mortalities were observed in 1996, and one in years 1998 and 1999. The 1996 and 1998 bycatch occurred in the *Illex* squid fishery, and the 1999 in the *Loligo* fishery reported for the Gulf of Maine fishery.

Southern New England/Mid-Atlantic Mixed Groundfish Trawl Fisheries

This fishery occurs year round, ranging from Cape Cod Massachusetts to Cape Hatteras North Carolina. It represents a variety of individual sub-fisheries that include but are not limited to: monkfish, summer flounder (fluke), winter flounder, silver hake (whiting), spiny and smooth dogfish, scup, and black sea bass. There was one observed take in this fishery reported in 1999. The estimated fishery-related mortality to for pilot whales in the USA Atlantic attributable to this fishery was: 45 in 1996 (CV=1.27), 0 in 1997, 85 in 1998 (CV=0.65), 49 in 1999 (CV=0.97) and 34 in 2000 (CV=0.65); 0 in 1996-1998, 228 in 1999 and 0 in 2000-2003. The average annual mortality between 1996-1999 and 2000-2003 was 436 pilot whales (CV=0.45 CV=1.03) (Table 2). However, these estimates should be viewed with caution due to the extremely low (<1%) observer coverage.

Northeast Atlantic (Gulf of Maine/Georges Bank) Herring Fishery

There were no marine mammal takes observed from the domestic mid-water trawl fishing trips during 1999-2003. A U.S. joint venture (JV) mid-water (pelagic) trawl fishery was conducted on Georges Bank from August - December 2001. A Total Allowable Level of Foreign Fishing (TALFF) was also granted during the same time period. Eight pilot whales were incidentally captured in a single mid-water trawl during JV fishing operations (TALFF) (Table 2). The total mortality attributed to the Atlantic herring mid-water trawl fishery from 1999-2003 was 2 animals (Table 2).

Mobile Gear Restricted Areas

Mobile gear restricted areas (GRA's) were put in place for fishery management purposes in November 2000. The intent of the GRA is to reduce bycatch of scup. The GRA's are spread out in time and space along the edge of the Southern New England and mid-Atlantic continental shelf region (between 100-1000 meters). These seasonal closures are targeted at trawl gear with small mesh sizes (<4.5 inches). The Atlantic herring and Atlantic mackerel trawl fisheries are exempt from the GRA's. A temporary exemption was also granted for the *Loligo* squid fishery. For detailed information regarding GRA's refer to FR/Vol. 66, No. 41.

Mid-Atlantic Coastal Gillnet

Observer coverage of the USA Atlantic coastal gillnet fishery was initiated by the NEFSC Fisheries Observer program in July 1993; and from July to December 1993, 20 trips were observed. During 1994 and 1995, 221 and 382 trips were observed, respectively. This fishery, which extends from North Carolina to New York, is actually a combination of small vessel fisheries that target a variety of fish species, some of which operate right off the beach. The number of vessels in this fishery is unknown, because records which are held by both state and federal agencies have not been centralized and standardized. Observer coverage, expressed as percent of tons of fish landed, was 5%-4%, 3%, 5%, 2%, and 2% for 1995, 1996, 1997, 1998, 1999, and 2000, respectively (Table 2).

No pilot whales were taken in observed trips during 1993-1997. One pilot whale was observed taken in 1998, 0 in 1999 and 2000 during 1999-2003 (Table 2). Observed effort was concentrated off NJ and scattered between DE New York and NC North Carolina from 1 to 50 miles off the beach. All bycatches were documented during January to April. Using the observed takes, the estimated annual mortality (CV in parentheses) attributed to this fishery was 7 in 1998 (1.1). Average annual estimated fishery-related mortality attributable to this fishery during 1996-2000 between 1999 and 2003 was ±0 (zero) pilot whale (CV=1.1) whales.

CANADA

An unknown number of pilot whales have also been taken in Newfoundland; and Labrador, and in Bay of Fundy groundfish gillnets, Atlantic Canada and Greenland salmon gillnets, and Atlantic Canada cod traps (Read 1994). The Atlantic Canadian and Greenland salmon gillnet fishery is seasonal, with the peak from June to September, depending on location. During 1989 in southern and eastern Newfoundland and in Labrador, 2,196 nets 91 m long were used. There are no effort data available for the Greenland fishery; however, the fishery was terminated in 1993 under an agreement between Canada and North Atlantic Salmon Fund (Read 1994).

There were 3,121 cod traps operating in Newfoundland and Labrador during 1979, and about 7,500 in 1980 (Read 1994). This fishery was closed at the end of 1993 due to collapse of Canadian groundfish resources.

Between January 1993 and December 1994, 36 Spanish deep-water trawlers, covering 74 fishing trips (4,726 fishing days and 14,211 sets), were observed in NAFO Fishing Area 3 (off the Grand Banks) (Lens 1997). A total of 47 incidental catches were recorded, which included 1 long-finned pilot whale. The incidental mortality rate for pilot

whales was 0.007/set.

In Canada, the fisheries observer program places observers on all foreign fishing vessels, on between ~~25-40%~~ 25% and 40% of large Canadian vessels (greater than ~~100 ft~~ 100ft), and on approximately 5% of small vessels (Hooker *et al.* 1997).- Fishery observer effort off the coast of Nova Scotia during 1991-1996 varied on a seasonal and annual basis, reflecting changes in fishing effort (see Figure 3, Hooker *et al.* 1997). During the 1991-~~1996~~ period, long-finned pilot whales were bycaught (number of animals in parentheses) in bottom trawl (65); midwater trawl (6); and longline (1) gear. Recorded bycatches by year were: 16 in 1991, 21 in 1992, 14 in 1993, 3 in 1994, 9 in 1995; and 6 in 1996. Pilot whale bycatches occurred in all months except January-March and September (Hooker *et al.* 1997).

Table 2. Summary of the incidental mortality and serious injury of pilot whales (*Globicephala sp.*) by commercial fishery including the years sampled (Years), the number of vessels active within the fishery (Vessels), the type of data used (Data Type), the annual observer coverage (Observer Coverage), the observed mortalities and serious injuries recorded by on-board observers, the estimated annual mortality and serious injury, the combined annual estimates of mortality and serious injury (Estimated Combined Mortality), the estimated CV of the combined estimates (Estimated CVs) and the mean of the combined estimates (CV in parentheses).

Fishery	Years	Vessels ¹	Data Type ¹	Observer Coverage ²	Observed Serious Injury	Observed Mortality	Estimated Serious Injury	Estimated Mortality	Estimated Combined Mortality	Estimated CVs	Mean Annual Mortality
<u>*SNE/mid-Atlantic</u> <u>Illex Squid Trawl</u>	<u>99-03</u>	<u>73⁵</u>	Obs. Data <u>Logbook Dealer</u>	<u>.0328, .111, .003, .03, .04, .04, .04</u>	<u>0, 0, 4, 40, 0, 0, 1, 10, 0, 0, 288, 1090, 0, 0, 930, 24</u>	<u>0, 0, 2, NA, NA, 0, 381, 133</u>	<u>0, 0, 0, 0.79, 0.88, 0, 0, NA, NA, 0</u>	<u>.0 34, NA, NA, 0</u>	<u>.0 34, NA, NA, 0</u>	<u>0, 0.65, NA, NA, 0</u>	<u>1.03 (0.65)</u>
<u>*SNE/mid-Atlantic</u> <u>Coastal Gillnet-Loligo</u> <u>Squid Trawl</u> <u>(offshore)</u>	<u>1996-2000</u> <u>99-03</u>	<u>NA</u> <u>384⁵</u>	Obs. Data <u>Logbook Dealer</u>	<u>.0309, .0311, .03, .02012, .02005, NA</u>	<u>0, 0, 0, 0, 0</u>	<u>0, 0, 0, 0</u>	<u>0, 0, 0, 0, 0, 0</u>	<u>0, 49, 0, 0, 0, 0</u>	<u>.49, 0, 0, 0, 0</u>	<u>0, 0.797, 0, 0.6, 1.1, 0, 0</u>	<u>10 (0.797)</u>
<u>Nova Scotia</u> <u>*SNE/ mid-Atlantic</u> <u>Bottom Trawl</u> <u>fisheries</u>	<u>1996</u> <u>99-03</u>	NA	Obs. Data <u>Dealer</u>	<u>.003, .003, .004, .005, NA</u>	<u>0, 0, 0, 0, 0</u>	<u>1, 0, 0, 0, 0</u>	<u>0, 0, 0, 0, 0</u>	<u>.0 228, 0, 0, 0, 0</u>	<u>.0 228, 0, 0, 0, 0</u>	<u>1.03, 0, 0, 0, 0</u>	<u>46 (1.03)</u>
<u>GOM/GB Herring</u> <u>Mid-Water Trawl</u> <u>IV and TALFF²</u>	<u>99-03</u>	<u>1999-2000=0 2001=10⁸ 2002-2003=0</u>	Obs. Data	<u>NA, NA 1.00, NA, NA</u>	<u>0, 0, 0, 0, 0</u>	<u>0, 0, 0, 11, 0, 0</u>	<u>0, 0, 0, 0, 0</u>	<u>0, 0, 0, 11, 0, 0</u>	<u>.0 0, 0, 0, 11, 0, 0</u>	NA	<u>6 (NA) 2 (NA)</u>
<u>Pelagic³</u> <u>Longline (excluding</u> <u>NED-E)¹⁰</u>	<u>99-03</u>	<u>. 198, 180, 152, 135, 116</u>	Obs. Data <u>Logbook</u>	<u>.04 .04, .02 .04, .02</u>	<u>.4, 4, 4, 4, 2</u>	<u>.1, 1, 1, 0, 0</u>	<u>.288, 109, 50, 52, 21</u>	<u>.93, 24, 20, 2, 0</u>	<u>.381, 133, 70, 54, 21</u>	<u>.79, .88, .50, .46, .77</u>	<u>132 (0.49)</u>
<u>Pelagic Longline -</u> <u>NED-E area only^{3, 10}</u>	<u>2001-2003</u>	<u>180 sets, 482, 535 NA, NA, 9, 14, 11</u>	Obs. Data <u>Logbook</u>	<u>1, 1, 1</u>	<u>0, 0, 0</u>	<u>0, 0, 0</u>	<u>0, 0, 0</u>	<u>0, 0, 0</u>	<u>0, 0, 0</u>	<u>0</u>	<u>0</u>
<u>Mid-Atlantic</u> <u>Coastal Gillnet</u>	<u>99-03</u>	<u>NA</u>	Obs. Data <u>Dealer</u>	<u>.02, .02, .02, .01, .01</u>	<u>0, 0, 0, 0, 0</u>	<u>0, 0, 0, 0, 0</u>	<u>0, 0, 0, 0, 0</u>	<u>.0 0, 0, 0, NA¹⁰, 0</u>	<u>.0 0, 0, 0, NA¹⁰, 0</u>	<u>0, 0, 0, NA¹⁰, 0</u>	<u>0 (0)</u>
TOTAL		<u>199</u>									<u>261 (0.49)</u>

¹ Observer data (Obs. Data) are used to measure bycatch rates, and the data are collected within the Northeast Fisheries Science Center (NEFSC) Fisheries Observer Sea Sampling Program. Mandatory logbook data were used to measure total effort for the longline fishery. These data are collected at the Southeast Fisheries Science Center (SEFSC).

² Observer coverage of the mid-Atlantic coastal gillnet fishery is measured in tons of fish landed. Observer coverage for the longline fishery

are in terms of sets. The trawl fisheries are measured in trips.

³ In 1997, 1998, and 2000, the observed pilot whales were taken from the *Illex* squid otter trawl subfishery. The 1999 observed pilot whales were taken from the *Loligo* squid and N. Atlantic otter trawl subfisheries.

⁴ 1996-2000/1997-1998 mortality estimates were taken from Table 9a in Yeung *et al.* (NMFS Miami Laboratory PRD 99/00-13), and excludes the Gulf of Mexico. 2000/1999-2000 mortality estimates were taken from Table 10 in Yeung 2000 (NMFS, Miami Lab. Ref. Doc. 00/01-17) and excludes the Gulf of Mexico.

⁵ NOAA Technical Memorandum NMFS-SEFSC-467). 2001-2002 and 2003 mortality estimates were taken from Tables 5,7, and 10 in Garrison (2003) and Garrison and Richards (2004).

⁶ Number of vessels in the fishery are based on vessels reporting effort to the pelagic longline logbook.

⁷ These are numbers of potential fishing vessels based on permit holders in the 2002 fishery. Many of these vessels participate in the other fisheries and therefore the reported number of vessels are not additive across the squid, mackerel and butterfish fisheries. (67FR 65937).

⁸ The incidental take was observed on a trip than landed either hake as the primary species.

⁹ During joint venture fishing operations, nets that are transferred from the domestic vessel to the foreign vessels for processing are observed on board the foreign vessel. There may be nets fished by domestic vessels that do not get transferred to a foreign vessel for processing and therefore would not be observed. During IALFF fishing operations all nets fished by the foreign vessel are observed.

¹⁰ These foreign vessels and seven American vessels.

¹¹ NA - No joint venture fishing effort for Atlantic mackerel.

¹² An experimental program to test effects of gear characteristics, environmental factors, and fishing practices on marine turtle bycatch rates in the Northeast Distant (NED-E) water component of the fishery was conducted from June 1, 2001-December 31, 2003. Observer coverage was 100% during this experimental fishery. Summaries are provided for the pelagic longline EXCLUDING the NED-E area in one row and for ONLY the NED in the second row. No mortalities nor serious injuries were observed for pilot whales in the NED-E, though 1 pilot whale was caught alive and released without injury (Garrison, 2003; Garrison and Richards, 2004).

¹³ Sixty-five percent of sampling in the mid-Atlantic coastal gillnet by the NEFSC fisheries observer program was concentrated in one area off the coast of Virginia. Because of the low level of sampling that was not distributed proportionately throughout the mid-Atlantic region, observed mortality is considered unknown in 2002. The previous four year average (1999-2002, and 2003) estimated mortality was applied as the best representative estimate.

Other Mortality

Pilot whales have a propensity to mass strand throughout their range, but the role of human activity in these events is unknown. Between 2 and 120 pilot whales have stranded annually, either individually or in groups, in NMFS Northeast Region (Anon. 1993b) since 1980. From 1992-2000, 98 pilot whales (*Globicephala sp.*) have been reported stranded between Nova Scotia and Florida (Table 3), including several mass strandings as follows: 14 live mass stranded whales in 2000 and 3 in 2001 in Judique, Inverness County; 4 live mass stranded whales at Point Tupper, Inverness County in 2002; 11 mass stranded whales in 2000 and 57 in 2002 in Massachusetts; and 28 whales that stranded in Content Passage, Monroe County, FL (ocean side) on April 18, 2003. The fate of the animals is specified in Table 3.

Table 3. Pilot whales (*Globicephala macrohynchus*) strandings along the Atlantic coast, 1999-2003

<u>STATE</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>TOTALS</u>
<u>Nova Scotia¹</u>	<u>1</u>	<u>16²</u>	<u>3³</u>	<u>7⁴</u>	<u>2</u>	<u>29</u>
<u>Maine</u>	<u>0</u>	<u>0</u>	<u>5</u>	<u>2</u>	<u>1</u>	<u>8</u>
<u>New Hampshire</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>
<u>Massachusetts</u>	<u>6</u>	<u>13⁵</u>	<u>3</u>	<u>67⁶</u>	<u>5</u>	<u>94</u>
<u>Rhode Island</u>	<u>0</u>	<u>0</u>	<u>1</u>	<u>1</u>	<u>0</u>	<u>2</u>
<u>Connecticut</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>
<u>New York</u>	<u>1</u>	<u>1</u>	<u>1</u>	<u>0</u>	<u>0</u>	<u>3</u>
<u>New Jersey</u>	<u>1</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>6</u>	<u>7</u>
<u>Delaware</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>
<u>Maryland</u>	<u>1</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>1</u>
<u>Virginia</u>	<u>2</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>2</u>
<u>North Carolina</u>	<u>2⁷</u>	<u>0</u>	<u>2⁸</u>	<u>0</u>	<u>1⁹</u>	<u>5</u>
<u>South Carolina</u>	<u>0</u>	<u>0</u>	<u>1</u>	<u>0</u>	<u>1¹⁰</u>	<u>2</u>
<u>Georgia</u>	<u>0</u>	<u>1</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>1</u>
<u>Florida</u>	<u>2¹¹</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>29^{12, 13}</u>	<u>31</u>

<u>TOTALS</u>	<u>16</u>	<u>31</u>	<u>16</u>	<u>77</u>	<u>45</u>	<u>185</u>
---------------	-----------	-----------	-----------	-----------	-----------	------------

¹ Data supplied by Tonya Wimmer, Nova Scotia Marine Animal Response Society (pers. comm.). NOTE: These strandings were NOT included in the long-finned pilot whale section, thus the table totals will be different, and will be updated in the next revision of the long-finned pilot whale stranded between South Carolina and Maine, including 22 and 11 animals that mass stranded in 1992 and 2000, respectively, along the Massachusetts coast (NMFS unpublished data). Four of 6 animals from stock assessment.

² Includes 14 mass live strandings at Judique, Inverness County on August 6, 2000 - 11 returned to sea
³ Three mass live stranded animals at Judique, Inverness County on July 19, 2001 - all returned to sea

⁴ Includes 4 mass live strandings at Point Tupper, Inverness County on January 11, 2002 - fate unreported.

⁵ Includes mass stranding of 11 animals in July 2000

⁶ Includes mass stranding of 57 animals in July 2002

⁷ Two long-finned pilot whales stranded in NC in 1999, reported to species

⁸ Two pilot whales stranded in NC in 2001 not identified to species

⁹ One pilot whale stranded in NC in 2003 not identified to species. NOTE: This stranding was NOT included in the long-finned pilot whale section, thus the table totals will be different, and will be updated in the next revision of the long-finned pilot whale stock assessment.

¹⁰ Only moderate confidence on species identification as long-finned pilot whale

¹¹ Two long-finned pilot whales reported in Florida identified to species

¹² Includes mass live stranding event in Massachusetts in 2000 were rehabilitated and released. In addition 11 pilot whales that live stranded on Nantucket of 28 short-finned pilot whales in Content Passage, Monroe County, FL (Ocean side) on April 19, 2003 - 12 animals died or were euthanized at the scene, 9 were returned to sea, 7 were taken into rehabilitation of which 2 subsequently died and 5 were released to sea on August 10, 2003. NOTE: These strandings were NOT included in the long-finned pilot whale section, thus the table totals will be different, and will be updated in the next revision of the long-finned pilot whale stock assessment.

¹³ Signs of human interaction reported on 1 stranded animal (not part of the live mass stranding). NOTE: These strandings were NOT included in the long-finned pilot whale section, thus the table totals will be different, and will be updated in the next revision of the long-finned pilot whale stock assessment.

Short-finned pilot whales (*Globicephala macrorhynchus*) have been reported stranded as far north as Nova Scotia (1990) Block Island, Rhode Island (2001) and long-finned pilot whales (173 individuals) were reported on Sable Island, Nova Scotia from 1970-1998 (Lucas and Hooker 1997; Lucas and Hooker 2000). This included 130 animals that mass stranded in December 1976, and 2 smaller groups (<10 each) in autumn 1979 and summer 1992. Fourteen strandings were also recorded along Nova Scotia from 1991-1996 (Hooker *et al.* 1997). *Globicephala melas* as far south as South Carolina.

A potential human-caused source of mortality is from polychlorinated biphenyls (PCBs) and chlorinated pesticides (DDT, DDE, dieldrin, etc.), moderate levels of which have been found in pilot whale blubber (Taruski 1975; Muir *et al.* 1988; Weisbrod *et al.* 2000). Weisbrod *et al.* (2000) reported that bioaccumulation levels were more similar in whales from the same stranding group than animals of the same sex or age. Also, high levels of toxic metals (mercury, lead, cadmium) and selenium were measured in pilot whales harvested in the Faroe Island drive fishery (Nielsen *et al.* 2000). Similarly, Dam and Bloch (2000) found very high PCB levels in pilot whales in the Faroes. The population effect of the observed levels of such contaminants is unknown.

STATUS OF STOCK

The status of long-finned short-finned pilot whales relative to OSP in US the U.S. Atlantic EEZ is unknown, but stock abundance may have been affected by reduction in foreign fishing, curtailment of the Newfoundland drive fishery for pilot whales in 1971, and increased abundance of herring, mackerel, and squid stocks. There are insufficient data to determine the population trends for this species. The species is not listed under the Endangered Species Act. The total fishery-related mortality and serious injury for this stock is not less than 10% of the calculated PBR, and; therefore, cannot be considered to be insignificant and approaching zero mortality and serious injury rate. This is not a strategic stock, because the ~~1996-2000~~1999-2003 estimated average annual fishery-related mortality, ~~excluding Nova Scotia bycatches~~ to pilot whales, *Globicephala* spp., ~~exceeds PBR.~~ does not exceed PBR. The status has gone back and forth, because mortality has been close to PBR. In the last six editions of this stock assessment report, it has been designated as non-strategic in 1998 and 1999 and this year.

REFERENCES

Anon. 1991. Northeast cetacean aerial survey and interplatform study. NOAA, NMFS, SEFSC & NEFSC. Available from NEFSC, Woods Hole Laboratory, Woods Hole, MA. 4 pp.

Anon. 1993. Cruise results, NOAA ship DELAWARE II, Cruise No. DEL 93-06, Marine mammal survey, NOAA NMFS NEFSC, Woods Hole Laboratory, Woods Hole, MA 4 pp. Available from NEFSC, Woods Hole Laboratory, Woods Hole, MA. 4 pp.

Barlow, J., S. L. Swartz, T. C. Eagle and P. R. Wade. 1995. U.S. Marine Mammal Stock Assessments: Guidelines

- for Preparation, Background, and a Summary of the 1995 Assessments. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-6, 73 pp. Available from NMFS, Southwest Fisheries Science Center, 8604 La Jolla Shores Drive, La Jolla, CA, 92037-1508.
- Buckland, S. T., D. R. Anderson, K. P. Burnham and J. L. Laake. 1993. Distance 3.51. ~~D~~Borchers and L. Thomas. 2001. Introduction to distance sampling: estimating abundance of biological populations. Chapman & Hall, London, 446 pp.
- Cramer J. 1994. Large pelagic logbook newsletter - 1993. NOAA Tech. Mem. NMFS-SEFSC-352, 19 pp. NMFS, Southeast Fisheries Science Center, Miami, FL. Oxford University Press, New York, 432 pp.
- CETAP (Cetacean and Turtle Assessment Program). 1982. A Characterization of Marine Mammals and Turtles in the Mid- and North Atlantic Areas of the U.S. Outer Continental Shelf, Final Report, Contract AA551-CT8-48. U.S. NTIS PB83-215855, Bureau of Land Management, Washington, DC, 576 pp.
- Dam, M. and D. Bloch. 2000. Screening of mercury and persistent organochlorine pollutants in long-finned pilot whale (*Globicephala melas*) in the Faroe Islands. *Marine Pollution Bulletin* 40:1090-1099.
- Fairfield, C. P., G. T. Waring and M. H. Sano. 1993. Pilot whales incidentally taken during the distant water fleet Atlantic mackerel fishery in the mid-Atlantic Bight, 1984-88. Rep. int. Whal. Commn. pp. 107-116. In: G.P. Donovan, C.H. Lockyer and A. R. Martin (eds.) *Biology of Northern Hemisphere Pilot Whales*. Rep. Int. Whal. Commn. (Special Issue 14): 107-116.
- Gerrior, P., A. S. Williams and D. J. Christensen. 1994. Observations of the 1992 U.S. pelagic pair trawl fishery in the Northwest Atlantic. *Mar. Fish. Rev.* 56(3): 24-27.
- Goudey, C.A. 1995. The 1994 experimental pair trawl fishery for tuna in the northwest Atlantic, Massachusetts Institute of Technology, Sea Grant, MITSG 95-6, Cambridge, MA. 10 pp.
- Goudey, C.A. 1996. The 1995 experimental pair trawl fishery for tuna in the northwest Atlantic, Massachusetts Institute of Technology, Sea Grant, MITSG 95-6, Cambridge, MA. 13 i-x + 479 pp.
- Garrison, L.P. 2003. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2001-2002. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-SEFSC-515, 52 pp. Available from NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Road, Miami, FL 33149.
- Garrison, L.P. and P.M. Richards. 2004. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2003. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-SEFSC-527, 57 pp. Available from NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Road, Miami, FL 33149.
- Hansen, L. J., K. D. Mullin, T. A. Jefferson and G. P. Scott. 1996. Visual surveys aboard ships and aircraft. pp. 55-132. In: R. W. Davis and G. S. Fargion (eds.), *Distribution and abundance of marine mammals in the north-central and western Gulf of Mexico: Final report. Volume II: Technical report*. OCS Study MMS 96-0027. Minerals Management Service, Gulf of Mexico OCS Region, New Orleans.
- Hiby, L. 1999. The objective identification of duplicate sightings in aerial survey for porpoise. pp. 179-189. In: G.W. Garner, S. C. Amstrup, J. L. Laake, B. F. J. Manly, L. L. McDonald, and D. G. Robertson (eds.) *Marine mammal survey and assessment methods*. Balkema, Rotterdam, 287 pp.
- Hooker, S. K., R. W. Baird and M. A. Showell. 1997. Cetacean strandings and bycatches in Nova Scotia, Eastern Canada, 1991-1996. Paper SC/49/05 presented to the IWC Scientific Committee, September 1997. 11 pp.
- Johnson, D. R., C. Yeung and C. A. Brown. 1999. Estimates of marine mammal and marine turtle catch by the U.S. Atlantic pelagic longline fleet in 1992-1997. NOAA Tech. Mem. NMFS-SEFSC-418. 70 pp. NMFS, Southeast Fisheries Science Center, Miami, FL.
- Laake, J. L., S. T. Buckland, D. R. Anderson and K. P. Burnham. 1993. DISTANCE user's guide, V2.0. Colorado Cooperative Fish & Wildlife Research Unit, Colorado State University, Ft. Collins, Colorado, 72 pp.
- Kingsley, M. C. S. and R. R. Reeves. 1998. Aerial survey of cetaceans in the Gulf of St. Lawrence in 1995 and 1996. *Can. J. Zool.* 76:1529-1550.
- Leatherwood, S. and R. R. Reeves. 1983. *The Sierra Club handbook of whales and dolphins*. Sierra Club Books, San Francisco, 302 pp.
- Leatherwood, S., D. K. Caldwell and H. E. Winn. 1976. Whales, dolphins, and porpoises of the western North Atlantic. A guide to their identification. U.S. Dept. of Commerce, NOAA Tech. Rep. NMFS Circ. 396, 176 pp.
- Lee, D. W., C. J. Brown, A. J. Catalano, J. R. Grubich, T. W. Greig, R. J. Miller and M. T. Judge. 1994. SEFSC pelagic longline observer program data summary for 1992-1993. NOAA Tech. Mem. NMFS-SEFSC-347. 19 pp. Available from: NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Drive, Miami, FL 33149.
- Lens, S. 1997. Interactions between marine mammals and deep water trawlers in the NAFO regulatory area. ICES CM 1997/Q:8. 10 pp.
- Lucas, Z. N. and S. K. Hooker. 1997. Cetacean strandings on Sable Island, Nova Scotia, 1990-1996. Paper SC/49/06 presented to the IWC Scientific Committee, September 1997. 10 pp.
- Lucas, Z. N. and S. K. Hooker. 2000. Cetacean strandings on Sable Island, Nova Scotia, 1970-1998. *Can. Field Nat.* 114(4):45-61.
- Mercer, M. C. 1975. Modified Leslie-DeLury population models of the long-finned pilot whale (*Globicephala melana*) and annual production of the short-finned squid (*Illex illecebrosus*) based upon their interactions at Newfoundland. *J. Fish. Res. Bd. Can.* 32(7):1145-1154.
- Mitchell, E. 1974. Present status of northwest Atlantic fin and other whale stocks. Pages 108-169. In: W. E. Schevill (ed.) *The whale problem: A status report*. Harvard University Press, Cambridge, MA, 419 pp.

- Muir, D. C. G., R. Wagermann, N. P. Grift, R. J. Norstrom, M. Simon and J. Lien. 1998. Organochlorine chemical and heavy metal contaminants in white-beaked dolphins (*Lagenorhynchus albirostris*) and pilot whales (*Globicephala melaena*) from the coast of Newfoundland. *Canada Arch. Environ. Contam. Toxicol.* 17:613-629.
- Mullin, K. D. (in review) and G.L. Fulling. 2003. Abundance and distribution of cetaceans in the southern U.S. North Atlantic Ocean during summer 1998. *Fish. Bull., U.S.* 101:603-613.
- Mullin, K., W. Hoggard, C. Roden, R. Lohocfencer, C. Rogers and B. Taggart. 1991. Cetaceans on the upper continental slope in the north-central Gulf of Mexico. OCS Study/MMS 91-0027. U.S. Dept. of the Interior, D. and G. L. Fulling. 2004. Abundance of cetaceans in the oceanic northern Gulf of Mexico, 1996-2001. *Mar. Mamm. Sci.* 20(4):787-807.
- Mullin, K. D. and W. Hoggard. 2000. Visual surveys of cetaceans and sea turtles from aircraft and ships. Pages 111-172. In: R. W. Davis, W. E. Evans, and B. Würsig (editors), *Cetaceans, sea turtles and seabirds in the northern Gulf of Mexico: Distribution, abundance and habitat associations. Volume II: Technical report*. OCS Study MMS 96-0027. Minerals Management Service, Gulf of Mexico OCS Regional Office Region, New Orleans; Louisiana, 108 pp.
- Nielsen, J. B., F. Nielsen, P.-J. Jørgensen, and P. Grandjean. 2000. Toxic metals and selenium in blood from pilot whales (*Globicephala melas*) and sperm whales (*Physeter catodon*). *Mar. Pollut. Bull.* 40:348-351.
- Northridge, S. 1996. Estimation of cetacean mortality in the U.S. Atlantic swordfish and tuna drift gillnet and pair trawl fisheries. Final report to the Northeast Fisheries Science Center, Contract No. 40ENNF500160. 348-35.
- Palka, D. 1995. Abundance estimate of the Gulf of Maine harbor porpoise. *Rep. int. Whal. Commn.* pp. 27-50 In: A. Bjørge and G.P. Donovan (eds.) *Biology of the Phocoenids.. Rep. Int. Whal. Commn* (Special Issue 16):27-50.
- Palka, D., G. Waring and D. Potter. (in review). Abundances of cetaceans and sea turtles in the northwest Atlantic during summer 1995 and 1998. *Fish. Bull., U.S.*
- Read, A. J. 1994. Interactions between cetaceans and gillnet and trap fisheries in the northwest Atlantic. *Rep. int. Whal. Commn. Special Issue 15*: 133-147.
- Scott, G. P. and C. A. Brown. 1997. Estimates of marine mammal and marine turtle catch by the U.S. Atlantic pelagic longline fleet in 1994-1995. Miami Laboratory Contribution MIA-96/97-28.
- Wade, P. R. and R. P. Angliss. 1997. i-x + 552 pp.
- Palka, D. 1996. Update on abundance of Gulf of Maine/Bay of Fundy harbor porpoises. NOAA/NMFS/NEFSC. Ref. Doc. 96-04; 37 pp. Available from National Marine Fisheries Service, 166 Water Street, Woods Hole, MA 02543-1026.
- Palka, D. and P.S. Hammond. 2001. Accounting for responsive movement in line transect estimates of abundance. *Can. J. Fish. Aquat. Sci.* 58: 777-787.
- Taruski, A. G., C. E. Olney and H. E. Winn. 1975. Chlorinated hydrocarbons in cetaceans. *J. Fish. Res. Bd. Can.* 32(11):2205-2209.
- Thomas, L., J. L. Laake, J. F. Derry, S. T. Buckland, D. L. Borchers, D. R. Anderson, K. P. Burnham, S. Strindberg, S.L. Hedley, F. F. C. Marques, J. H. Pollard and R. M. Fewster. 1998. Distance 3.5. Research Unit for Wildlife Population Assessment, University of St. Andrews, St. Andrews, UK.
- Wade, P.R., and R.P. Angliss. 1997. Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-12, 93 pp.
- Waring, G. T., P. Gerrior, P. M. Payne, B. L. Parry and J. R. Nicolas. 1990. Incidental take of marine mammals in foreign fishery activities off the northeast United States, 1977-1988. *Fish. Bull., U.S.* 88(2): 347-360 Available from: NOAA, National Marine Fisheries Service, 166 Water St., Woods Hole, MA 02543.
- Waring, G. T., C. P. Fairfield, C. M. Ruhsam and M. Sano. 1992. Cetaceans associated with Gulf Stream features off the northeastern USA shelf. *ICES Marine Mammals Comm.* CM 1992/N:12, 29 pp.
- Waring, G. T. 1998. Results of the summer 1991 R/V Chapman marine mammal sighting survey. NOAA NMFS NEFSC, Lab. Ref. Doc. No. 98-09, 21pp. Available from: National Marine Fisheries Service, 166 Water Street, Woods Hole, MA 02543-1026.
- Waring, G. T. 1995. *Fishery and ecological interactions for selected cetaceans off the northeast USA*. Ph.D. dissertation, University of Massachusetts, Amherst, 260 pp.
- Waring, G. T., D. L. Palka, P. J. Clapham, S. Swartz, M. C. Rossman, T. V. N. Cole, L. J. Hansen, K. D. Bisack, K. D. Mullin, R. S. Wells, D. K. Odell, and N. B. Barros. 1999. U.S. Atlantic and Gulf of Mexico marine mammal stock assessments — 1999. U.S. Dep. Commer., NOAA Tech. Memo NMFS-NE-153, 196pp.
- Weisbrod, A. V., D. Shea, M. J. Moore, and J. J. Stegeman. 2000. Bioaccumulation patterns of polychlorinated biphenyls and chlorinated pesticides in northwest Atlantic pilot whales. *Environ. Toxicol. Chem.* 19:667-677.
- Yeung, C. 2001. Estimates of marine mammal and marine turtle bycatch by the U.S. Atlantic pelagic longline fleet in 1999-2000. NOAA Tech. Memo. NMFS-SEFSC-467, 43 pp. NMFS, Southeast Fisheries Science Center, Miami, FL.
- Yeung, C. 1999a. Estimates of marine mammal and marine turtle bycatch by the U.S. Atlantic pelagic longline fleet in 1998. NOAA Tech. Memo. NMFS-SEFSC-430, 26 pp. NMFS, Southeast Fisheries Science Center, Miami, FL.
- Yeung, C. 1999b. Revised Mortality Estimates of Marine Mammal Bycatch in 1992-1997 based on Serious Injury

~~Guidelines. NOAA Tech. Memo. NMFS-SEFSC-429, 23 pp. NMFS, Southeast Fisheries Science Center, Miami, FL.~~

~~Yeung, C., S. Epperly, and C. A. Brown. 2000. Preliminary revised estimates of marine mammals and marine turtle bycatch by the U.S. Atlantic pelagic longline fleet, 1992-1999. NMFS, Miami Lab. PRD Contribution Number 99/00-13. -58 pp.~~

~~Yeung, C. 2001. Estimates of marine mammal and marine turtle bycatch by the U.S. Atlantic pelagic longline fleet in 1999-2000. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-SEFSC-467, National Marine Fisheries Service, Miami, FL, 43 pp. Available from NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Road, Miami, FL 33149.~~

~~December 1998~~ February 2005

SPINNER DOLPHIN (*Stenella longirostris*): Western North Atlantic Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

Spinner dolphins are distributed in oceanic and coastal tropical waters (Leatherwood *et al.*: 1976). This is presumably an offshore, deep-water species (Schmidly 1981; Perrin and Gilpatrick 1994), and its distribution in the Atlantic is very poorly known. In the western North Atlantic, these dolphins occur in deep water along most of the U.S. coast south to the West Indies and Venezuela, including the Gulf of Mexico. Spinner dolphin sightings have occurred exclusively in deeper (>2,000 m) oceanic waters (CETAP 1982; Waring *et al.*: 1992; NMFS, unpublished data) off the northeast U.S. coast. Stranding records exist from North Carolina, South Carolina, and Florida and Puerto Rico in the Atlantic and in Texas and Florida in the Gulf of Mexico. ~~The North Carolina strandings represent the northernmost documented distribution of this species in the Atlantic.~~ Stock structure in the western North Atlantic is unknown.

POPULATION SIZE

The numbers of spinner dolphins ~~inhabiting~~off the U.S. or Canadian Atlantic Exclusive Economic Zone (EEZ) ~~is~~coast are unknown, and seasonal abundance estimates are not available for this speciesstock since it was rarely seen in any of the surveys.

Minimum Population Estimate

Present data are insufficient to calculate a minimum population estimate.

Current Population Trend

There are insufficient data to determine the population trends for this species stock.

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Current and maximum net productivity rates are unknown for this stock. For purposes of this assessment, the maximum net productivity rate was assumed to be 0.04. This value is based on theoretical modeling showing that cetacean populations may not grow at rates much greater than 4% given the constraints of their reproductive life history (Barlow *et al.*: 1995).

POTENTIAL BIOLOGICAL REMOVAL

Potential Biological Removal (PBR) is the product of minimum population size, one-half the maximum productivity rate, and a “recovery” factor (MMPA Sec. 3. 16 U.S.C. 1362; Wade and Angliss 1997). The minimum population size is unknown. The maximum productivity rate is 0.04, the default value for cetaceans. The “recovery” factor, which accounts for endangered, depleted, threatened stocks, or stocks of unknown status, relative to optimum sustainable population (OSP), is assumed to be 0.5 because this stock is of unknown status. PBR for the western North Atlantic spinner dolphin is unknown because the minimum population size is unknown.

ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

Fishery Information

Detailed fishery information is reported in Appendix III. Total ~~average~~ annual estimated average fishery-related mortality and serious injury to this stock ~~in the Atlantic~~ during ~~1992-1996~~ 1999-2003 was ~~0.38~~ zero spinner dolphin (CV = 0.35).

Fishery Information

—dolphins, as there were no reports of mortalities or serious injury to spinner dolphins (Yeung 2001; Garrison 2003; Garrison and Richards, 2004).

EARLIER INTERACTIONS

There was no documentation of spinner dolphin mortality or serious injury in distant-water fleet (DWF) activities off the northeast U.S. coast (Waring *et al.*: 1990). No takes were documented in a review of Canadian gillnet and trap fisheries (Read 1994).

Data on current incidental takes in U.S. fisheries are available from several sources. In 1986, NMFS established a mandatory self-reported Fishery information system for large pelagic fisheries. Data files are maintained at the Southeast Fisheries Science Center (SEFSC). The Northeast Fisheries Science Center (NEFSC) Fisheries Observer Program was initiated in 1989 and since that year several fisheries have been covered by the program. In late 1992 and in 1993, the SEFSC provided observer coverage of pelagic longline vessels fishing off the Grand Banks (Tail of the Banks) and provides observer coverage of vessels fishing south of Cape Hatteras.

By-catch has been observed by NMFS Sea Samplers in the now prohibited pelagic drift gillnet fishery, but no mortalities or serious injuries have been documented in the pelagic longline, pelagic pair trawl, Northeast multispecies sink gillnet, mid-Atlantic coastal gillnet, and North Atlantic bottom trawl fisheries.

The estimated total number of hauls in the pelagic drift gillnet fishery increased from 714 in 1989 to 1,144 in 1990; thereafter, with the introduction of quotas, effort was severely reduced. The estimated number of hauls in 1991, 1992, 1993, 1994, 1995, and 1996 were 233, 243, 232, 197, 164, and 149 respectively. Fifty-nine different vessels participated in this fishery at one time or another between 1989 and 1993. Since 1994, between 10-12 vessels have participated in the fishery (Table 2). Observer coverage, expressed as percent of sets observed, was 8% in 1989, 6% in 1990, 20% in 1991, 40% in 1992, 42% in 1993, 87% in 1994, 99% in 1995, and 64% in 1996. Effort was concentrated along the southern edge of Georges Bank and off Cape Hatteras. Examination of the species composition of the catch and locations of the fishery throughout the year, suggested that the pelagic drift gillnet fishery be stratified into two strata, a southern or winter stratum, and a northern or summer stratum. Estimates of the total by-catch, from 1989 to 1993, were obtained using the aggregated (pooled 1989-1993) catch rates, by strata (Northridge 1996). Estimates of total annual by-catch for 1994 and 1995 were estimated from the sum of the observed caught and the product of the average bycatch per haul and the number of unobserved hauls as recorded in self-reported Fishery information. Variances were estimated using bootstrap re-sampling techniques.

Pelagic Drift Gillnet

One spinner dolphin mortality was observed in the pelagic driftnet between 1989 and 1993 and occurred east of Cape Hatteras in March 1993. Estimated annual fishery-related mortality and serious injury attributable to this fishery (CV in parentheses) was 0.7 in 1989 (1.00), 1.7 in 1990 (1.00), 0.7 in 1991 (1.00), 1.4 in 1992 (0.31), 0.5 in 1993 (1.00), and zero from 1994-1996. Total average annual estimated average fishery-related mortality and serious injury to this stock in the Atlantic during 1992-1996 was 0.38 spinner dolphin (CV = 0.35) (Table 1). The 1992-1996 period provides a better characterization of this fishery (i.e., fewer vessels and increased observer coverage).

Table 1. Summary of the incidental mortality of spinner dolphins (*Stenella longirostris*) by commercial fishery including the years sampled (Years), the number of vessels active within the fishery (Vessels), the type of data used (Data Type), the annual observer coverage (Observer Coverage), the mortalities recorded by on-board observers (Observed Mortality), the estimated annual mortality (Estimated Mortality), the estimated CV of the annual mortality (Estimated CVs) and the mean annual mortality (CV in parentheses).

FisheryYears Vessels¹

Data Type²

Observer Coverage³

Observed MortalityEstimated Mortality⁴

Estimated CVs⁴

Mean Annual MortalityPelagic Drift Gillnet 92-961994-12

1995=11 1996=10Obs. Data Logbook.40, .42, .87, .99, .641, 0, 0, 0, 01.4, 0.5, 0, 0⁵, 0

— .31, 1.0, 0, 0, 00.31

(.35)TOTAL0.31

(.35)

- ¹ 1994 and 1995 - 1996 shown, other years not available on an annual basis.
- ² Observer data (Obs. Data) are used to measure bycatch rates, and the data are collected within the Northeast Fisheries Science Center (NEFSC) Fisheries Observer Program. Mandatory logbook (Logbook) data are used to measure total effort, and the data are collected at the Southeast Fisheries Science Center (SEFSC).
- ³ The observer coverage and unit of effort for the Pelagic Drift Gillnet is a set.
- ⁴ For 1991-1993, pooled bycatch rates were used to estimate bycatch in months that had fishing effort but did not have observer coverage. This method is described in Northridge (1996). In 1994 and 1995, observer coverage increased substantially, and bycatch rates were not pooled for this period.
- ⁵ One vessel was not observed and recorded 1 set in a 10 day trip in the SEFSC mandatory logbook. If you assume the vessel fished 1.4 sets per day as (Northridge 1996). Estimates of total annual bycatch for 1994 and 1995 were estimated from the 1995 SS data, the point estimate may increase by 0.8 animals. However, the SEFSC mandatory logbook data was taken at face value, and therefore it was assumed that 1 set was fished within this trip, and the point estimate would then increase by 0.1 animals.

sum of the observed caught and the product of the average bycatch per haul and the number of unobserved hauls as recorded in self-reported fisheries information. Variances were estimated using bootstrap re-sampling techniques. Estimated annual mortality and serious injury attributable to this fishery (CV in parentheses) was 0.7 in 1989 (1.00), 1.7 in 1990 (1.00), 0.7 in 1991 (1.00), 1.4 in 1992 (0.31), 0.5 in 1993 (1.00) and zero from 1994-1996. This fishery is no longer in operation.

Other Mortality

From 1999-2003, 9 spinner dolphins were reported stranded between Maine and Puerto Rico (Table 1). The total includes 2 animals stranded in North Carolina in 2001, 2 animals stranded in Puerto Rico in 2002, 4 mass stranded live animals in December 2003 in Flagler, Florida (all died on the scene), and 1 additional animal stranded in Florida in 2003. There were no indications of human interactions for these stranded animals.

Table 1. Spinner dolphin (*Stenella longirostris*) strandings along the U.S. Atlantic coast, 1999-2003

<u>STATE</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>TOTALS</u>
<u>North Carolina</u>	<u>0</u>	<u>0</u>	<u>2</u>	<u>0</u>	<u>0</u>	<u>2</u>
<u>South Carolina</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>
<u>Georgia</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>
<u>Florida</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>5¹</u>	<u>5</u>
<u>Puerto Rico</u>	<u>0</u>	<u>0</u>	<u>0</u>	<u>2</u>	<u>0</u>	<u>2</u>
<u>TOTALS</u>	<u>0</u>	<u>0</u>	<u>2</u>	<u>2</u>	<u>5</u>	<u>9</u>

¹Florida live mass stranding of 4 animals in Flagler, Florida on December 29, 2003

STATUS OF STOCK

The status of spinner dolphins, relative to OSPOPS, in the U.S. western North Atlantic EEZ is unknown. The species is not listed as threatened or endangered under the Endangered Species Act. There are insufficient data to determine the population size or trends for this species. and PBR cannot be calculated for this stock, but n. No fishery-related mortality and serious injury has been observed since 19929; therefore, total fishery-related mortality and serious injury rate can be considered insignificant and approaching zero mortality and serious injury rate. Population size and PBR cannot be estimated, but fishery-related mortality is very low; therefore, this stock. This is not a strategic stock.

REFERENCES

Barlow, J., S.L. Swartz, T.C. Eagle, and P.R. Wade. 1995. U.S. Marine Mammal Stock Assessments: Guidelines for

- Preparation, Background, and a Summary of the 1995 Assessments. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-6, 73 pp. Available from: NOAA, National Marine Fisheries Service, 7600 Sand Point Way NE, Seattle, WA, 98115-0070.
- CETAP. 1982. A characterization of marine mammals and turtles in the mid- and north Atlantic areas of the U.S. outer continental shelf. Cetacean and Turtle Assessment Program, University of Rhode Island. Final Report, Contract AA51-C78-48, Bureau of Land Management, Washington, DC, 538 pp.
- Garrison, L.P. 2003. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2001-2002. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-SEFSC-515, 52 pp. Available from NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Road, Miami, FL 33149.
- Garrison, L.P. and P.M. Richards. 2004. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2003. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-SEFSC-527, 57 pp. Available from NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Road, Miami, FL 33149.
- Leatherwood, S., D.K. Caldwell, and H.E. Winn. 1976. Whales, dolphins, and porpoises of the western North Atlantic. A guide to their identification. U.S. Dept. of Commerce, NOAA Tech. Rep. NMFS Circ. 396, 176 pp. Available from: NOAA, National Marine Fisheries Service, 166 Water St., Woods Hole, MA 02543.
- Northridge, S. 1996. Estimation of cetacean mortality in the U.S. Atlantic swordfish and tuna drift gillnet and pair trawl fisheries. Final report to the Northeast Fisheries Science Center, Contract No. 40ENNF500045, 18 pp. Available from: NOAA, National Marine Fisheries Service, 166 Water St., Woods Hole, MA 02543.
- Perrin, W. F. and J. W. Gilpatrick, Jr. 1994. Spinner dolphin. ~~Pages~~ pp. 99-128. ~~In:~~ In: S. H. Ridgway and R. Harrison (eds.), *Handbook of marine mammals, Volume 5: The first book of dolphins*. Academic Press, San Diego, California. 418 pp.
- Read, A. J. 1994. Interactions between cetaceans and gillnet and trap fisheries in the northwest Atlantic. ~~Rep. in~~ pp. 133-147. In: W.F. Perrin, G.P. Donovan and J. Barlow (eds.) *Gillnets and Cetaceans*. Rep. Int. Whal. Commn. (Special Issue 15: 133-147.). I-ix + 629 pp.
- ~~Reilly, S. B. and J. Barlow. 1986. Rates of increase in dolphin population size. *Fish. Bull., U.S.* 84(3): 527-533.~~
- Schmidly, D. J. 1981. Marine mammals of the southeastern United States coast and the Gulf of Mexico. Pub. No. FWS/OBS-80/41, U.S. Fish and Wildlife Service, Office of Biological Services, Washington, DC, 163 pp.
- Wade, P.R., and R.P. Angliss. 1997. Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-12, 93 pp. Available from: NOAA, National Marine Fisheries Service, 166 Water St., Woods Hole, MA 02543.
- Waring, G.T., P. Gerrior, P.M. Payne, B.L. Parry and J.R. Nicolas. 1990. Incidental take of marine mammals in foreign fishery activities off the northeastern United States 1977-1998. *Fish. Bull., U.S.* 88(2):347-360.
- Waring, G. T., C. P. Fairfield, C. M. Ruhsam and M. Sano. 1992. Cetaceans associated with Gulf Stream features off the northeastern USA shelf. ICES Marine Mammals Comm. CM 1992/N:12, 29 pp.
- Yeung, C. 2001. Estimates of marine mammal and marine turtle bycatch by the U.S. Atlantic pelagic longline fleet in 1999-2000. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-SEFSC-467, 43 pp. Available from: NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Dr., Miami, FL, 33149.
-
-